

[Click here to view linked References](#)

1
2
3 **Synthesis, leishmanicidal, trypanocidal and cytotoxic activities of**
4
5
6
7 **quinoline-chalcone and quinoline-chromone hybrids**
8
9

10
11
12
13
14 Juan C. Coa¹. Elisa García¹ . Miguel Carda² . Raúl Agut² . Iván D. Vélez³ . July A. Muñoz³ . Lina M.
15
16 Yepes³ Sara M. Robledo³ . Wilson I. Cardona¹
17
18
19
20
21
22
23
24
25

26 ¹ *Chemistry of Colombian Plants, Institute of Chemistry, Exact and Natural Sciences School,*
27
28 *University of Antioquia-UdeA, Calle 70 No. 52-21, A.A 1226, Medellín, Colombia*
29
30

31 ² *Department of Inorganic and Organic Chemistry, Jaume I University, E-12071 Castellón, Spain*
32
33

34 ³ *PECET-Medical Research Institute, School of Medicine, University of Antioquia-UdeA. Calle 70 No.*
35
36 *52-21, A.A 1226 Medellín, Colombia*
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

*Corresponding authors: *E-mail address:*

Wilson I. Cardona

wilson.cardona1@udea.edu.co

phone: +574-2195653

Fax: +57-42330120

Sara M. Robledo

sara.robledo@udea.edu.co

phone: +574-2196503

Fax: +574-2196511

1
2
3 **Abstract**
4
5
6
7
8

9 We report herein the synthesis and biological activities (cytotoxicity, leishmanicidal and trypanocidal)
10 of six quinoline-chalcone and five quinoline-chromone hybrids. The synthesized compounds were
11 evaluated against amastigotes forms of *L. (V) panamensis* which is the most prevalent *Leishmania*
12 species in Colombia and *Trypanosoma cruzi* which is the major pathogenic species to humans.
13
14 Cytotoxicity was evaluated against human U-937 macrophages. Compounds **8-12**, **20**, **23** and **24**
15 showed activity against *L. (V) panamensis* while compounds **9**, **10**, **12**, **20** and **23** had activity against *T.*
16 *cruzi* with EC₅₀ values lower than 18 mg/ml. **20** was the most active compound for both *L. (V)*
17 *panamensis* and *T. cruzi* with EC₅₀ of 6.11 ± 0.26 µg/ml (16.91 µM) and 4.09 ± 0.24 (11.32µM),
18 respectively. All hybrids compounds showed better activity than the anti-leishmanial drug meglumine
19 antimoniate. Compounds **20** and **23** showed higher actives than benznidazole, the current anti-
20 trypanosomal drug. Although these compounds showed toxicity for mammalian U-937 cells, still
21 have potential to be considered as candidates to antileishmanial or trypanocidal drug
22 development.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **Keywords:** leishmaniasis; Chagas disease; antiprotozoal activity; cytotoxicity; quinoline; chalcone;
52 chromone; hybrids
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

Neglected tropical diseases (NTD) are a cause of mortality in various developing countries of tropical and subtropical regions. These diseases are significant health problems in endemic countries, affecting more than one billion people worldwide (WHO, 2013). This situation is aggravated by increasing treatment failures with available drugs (Bhutta *et al.*, 2014). NTD include, among others, Chagas' disease (American trypanosomiasis) and leishmaniasis. These are parasitic diseases caused by the parasitic protozoan *Trypanosoma cruzi* (*T. cruzi*) and *Leishmania* species. These diseases affect more than 10 million people worldwide (Alvar *et al.*, 2012; Nouvellet *et al.*, 2015) *L. (V) panamensis* is one of the most prevalent *Leishmania* species involved in human cases of cutaneous leishmaniasis in Colombia (Alvar *et al.*, 2012). Current chemotherapies are based on old drugs, pentavalent antimonials (meglumine antimoniate and sodium stibogluconate) to treat cutaneous leishmaniasis and nitroaromatic compounds (benznidazole and nifurtimox) for treatment of Chagas disease. Unfortunately, all of these drugs are not very effective in the chronic phase and have toxicity, side effects and parasite resistance (Den Boer *et al.*, 2011; Keenan *et al.*, 2015; Chatelain *et al.*, 2011).

A quinolinic core is a structural feature of several bioactive compounds. Thus, this core is an interesting constituent for new drugs design. Anti-mycobacterial, anti-microbial, anti-convulsant, anti-inflammatory, anti-tumoral, cardiovascular but also leishmanicidal and trypanocidal, are some biological activities exhibited by compounds having this heteroaromatic ring (Suresh *et al.*, 2009; Nakayama *et al.*, 2005; Tempone *et al.*, 2005; Dietze *et al.*, 2001; Mohammed *et al.*, 2003; Vieira *et al.*, 2008; Cardona *et al.*, 2013; Palit *et al.*, 2009; Franck *et al.*, 2004; Coa *et al.*, 2015). The antileishmanial activity of several chalcones has been reported (Kayser *et al.*, 2001; Liu *et al.*, 2003; Boeck *et al.*, 2006). The most promising of this class of compounds is licochalcone A, an oxygenated chalcone

1
2
3 isolated from the roots of the Chinese plant *Glycyrrhiza* spp, which inhibits the fumarate reductase, a
4
5 selective target present in the mitochondria of the parasite (Chen *et al.*, 2001). Similarly, chromones
6
7 are important classes of compounds having versatile biological activities (Horton *et al.*, 2003;
8
9 Hadjeri *et al.*, 2003; Ellis *et al.*, 1972; Houghton *et al.*, 2000; Mallick *et al.*, 2011; Baloch *et al.*,
10
11 2012). Both moieties are well known for their antiprotozoal activity. Some synthetic chromones were
12
13 effective against *L. (L) donovani* (Mallick *et al.*, 2011) and *L. (L) major* (Baloch *et al.*, 2012) in *in*
14
15 *vivo* studies.
16
17
18
19
20
21

22 In the search for new therapeutic alternatives to treat cutaneous leishmaniasis and Chagas disease, in
23
24 this work we designed and synthesized a series of quinoline-chalcone and quinoline-chromone hybrids
25
26 and evaluated *in vitro* their cytotoxicity, leishmanicidal and trypanocidal activities.
27
28
29
30
31
32

33 **Material and Methods**

34 **Chemistry**

35 *General remarks*

36
37
38
39
40
41
42
43 Microwave reactions were carried out in a CEM Discover microwave reactor in sealed vessels
44
45 (monowave, maximum power 300 W, temperature control by IR sensor, fixed temperature). ¹H and ¹³C
46
47 NMR spectra were recorded on a Varian instrument operating at 500 and 125 MHz, respectively. The
48
49 signals of the deuterated solvent (CDCl₃) were used as reference (the singlet at $\delta = 7.27$ ppm for ¹H
50
51 NMR and the triplet centred at $\delta = 77.00$ ppm for ¹³C NMR). Carbon atom types (C, CH, CH₂, CH₃)
52
53 were determined by using the DEPT or APT pulse sequence. High resolution mass spectra were
54
55 recorded using electrospray ionization mass spectrometry (ESI-MS). A QTOF Premier instrument with
56
57 an orthogonal Z-spray-electrospray interface (Waters, Manchester, UK) was used operating in the
58
59
60
61
62
63
64
65

1
2
3 W-mode. The drying and cone gas was nitrogen set to flow rates of 300 and 30 L/h, respectively.
4
5 Methanol sample solutions (ca. 1×10^{-5} M) were directly introduced into the ESI spectrometer at a
6
7 flow rate of 10 μ L/min. A capillary voltage of 3.5 kV was used in the positive scan mode, and the cone
8
9 voltage set to $U_c = 10$ V. For accurate mass measurements, a 2 mg/L standard solution of leucine
10
11 enkephalin was introduced via the lock spray needle at a cone voltage set to 85 V and a flow rate of
12
13 30 μ L/min. IR spectra were recorded on a Spectrum RX I FT-IR system (Perkin-Elmer, Waltham, MA,
14
15 USA) in KBr disks. Silica gel 60 (0.063–0.200 mesh, Merck, Whitehouse Station, NJ, USA) was used
16
17 for column chromatography, and precoated silica gel plates (Merck 60 F254 0.2 mm) were used for
18
19 thin layer chromatography (TLC).
20
21
22
23
24
25
26
27

28 *General procedure for the synthesis of bromoalkyl derivatives*

29
30

31 Quinoline or chromone (1 mmol), potassium hydroxide (1.5 mmol, 84.2 mg) and acetonitrile (10
32
33 mL), were placed in a 25 mL flat-bottomed flask equipped with a magnetic stirring bar. The
34
35 mixture was stirred and heated to reflux for a period of 5 min, under microwave irradiation. Then, 1, ω -
36
37 dibromoalkane (1.1 mmol) was added to the reaction mixture which was refluxed for 30 minutes (70
38
39 W). The crude reaction mixture was concentrated on a rotatory evaporator and the residue was
40
41 purified by column chromatography over silica gel eluting with hexane and a mixture of
42
43 hexane-ethyl acetate (9:1 ratio) to obtain bromoalkyl derivatives in yields ranging between 42–80%.
44
45
46
47
48 Monitoring of the reaction progress and product purification was carried out by TLC.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 8-(3-bromopropoxy)quinoline (**2**): Yield 75% (0.75 mmol, 200 mg); brown oil. IR (cm⁻¹): ν_{\max} 1570
4
5 (C=C_{Ar}), 1500 (C=N), 1263 (C-O-C), 794 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 2.20-2.30 (2H, m), 4.03 (2H,
6
7 t, $J = 5.3$ Hz), 4.47 (2H, t, $J = 5.6$ Hz), 7.21 (1H, d, $J = 6.2$ Hz), 7.44-7.56 (3H, m), 8.20 (1H, dd, $J =$
8
9 8.3, 1.5 Hz), 8.94 (1H, dd, $J = 4.2, 1.5$ Hz). ¹³C-NMR (CDCl₃): δ 30.0 (CH₂), 31.9 (CH₂), 69.6 (CH₂),
10
11 111.6 (C), 120.6 (C), 121.7 (C), 126.8 (C), 129.6 (C), 136.2 (C), 140.7 (C), 149.4 (C), 154.8 (C).
12
13
14

15 8-(4-bromobutoxy)quinoline (**3**): Yield 64% (0.64 mmol, 179 mg); dark yellow solid, M.p. 46-48°C. IR
16
17 (cm⁻¹): ν_{\max} 1593 (C=C_{Ar}), 1529 (C=N), 1240 (C-O-C), 835 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 2.17-2.27
18
19 (4H, m), 3.57 (2H, t, $J = 5.8$ Hz), 4.29 (2H, t, $J = 5.5$ Hz), 7.09 (1H, d, $J = 7.5$ Hz), 7.39-7.53 (3H, m),
20
21 8.17 (1H, dd, $J = 8.3, 1.1$ Hz), 9.01 (1H, dd, $J = 4.2, 1.3$ Hz). ¹³C-NMR (CDCl₃): δ 27.7 (CH₂), 29.5
22
23 (CH₂), 33.7 (CH₂), 67.8 (CH₂), 108.8 (C), 119.7 (C), 121.7 (C), 126.8 (C), 129.6 (C), 136.3 (C), 139.9
24
25 (C), 149.2 (C), 154.5 (C).
26
27
28
29
30

31 8-((5-bromopentyl)oxy) quinoline (**4**): Yield 51% (0.51 mmol, 150 mg); brown solid, M.p. 58-62°C. IR
32
33 (cm⁻¹): ν_{\max} 1595 (C=C_{Ar}), 1529 (C=N), 1288 (C-O-C), 829 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.69-1.82
34
35 (2H, m), 1.97-2.07 (2H, m), 2.07-2.19 (2H, m), 3.50 (2H, t, $J = 6.8$ Hz), 4.29 (2H, t, $J = 6.8$ Hz), 7.10
36
37 (1H, d, $J = 7.5$ Hz), 7.40-7.53 (3H, m), 8.17 (1H, dd, $J = 8.2, 1.0$ Hz), 8.99 (1H, dd, $J = 4.2, 1.0$ Hz).
38
39 ¹³C-NMR (CDCl₃): δ 24.9 (CH₂), 28.2 (CH₂), 32.6 (CH₂), 33.6 (CH₂), 68.6 (CH₂), 108.7 (C), 119.6
40
41 (C), 121.6 (C), 126.7 (C), 129.6 (C), 136.0 (C), 140.3 (C), 149.4 (C), 154.7 (C).
42
43
44
45

46 8-((8-bromooctyl)oxy) quinoline (**5**): Yield 80% (0.80 mmol, 269 mg); light brown oil. IR (cm⁻¹): ν_{\max}
47
48 1593 (C=C_{Ar}), 1529 (C=N), 1286 (C-O-C), 831 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.33-1.50 (6H, m), 1.51-
49
50 1.64 (2H, m), 1.81-1.94 (2H, m), 1.99-2.11 (2H, m), 3.42 (2H, t, $J = 6.9$ Hz), 4.25 (2H, t, $J = 6.9$ Hz),
51
52 7.08 (1H, d, $J = 7.6$ Hz), 7.36-7.52 (3H, m), 8.15 (1H, dd, $J = 8.3, 1.5$ Hz), 8.99 (1H, dd, $J = 4.2, 1.6$
53
54 Hz). ¹³C-NMR (CDCl₃): δ 26.0 (CH₂), 28.1 (CH₂), 28.7 (CH₂), 29.0 (CH₂), 29.3 (CH₂), 32.8 (CH₂),
55
56 34.1 (CH₂), 68.9 (CH₂), 108.8 (C), 119.4 (C), 121.6 (C), 126.8 (C), 129.5 (C), 136.1 (C), 140.2 (C),
57
58 149.2 (C), 154.8 (C).
59
60
61
62
63
64
65

1
2
3 8-((9-bromononyl)oxy)quinoline (**6**): Yield 75% (0.75 mmol, 263 mg); light brown oil. IR (cm⁻¹): ν_{\max}
4
5 1595 (C=C_{Ar}), 1531 (C=N), 1286 (C-O-C), 833 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.31-1.52 (6H, m), 1.52-
6
7 1.66 (2H, m), 1.82-1.95 (2H, m), 2.00-2.13 (2H, m), 3.44 (2H, t, $J = 6.9$ Hz), 4.26 (2H, t, $J = 6.9$ Hz),
8
9 7.10 (1H, d, $J = 7.6$ Hz), 7.38-7.53 (3H, m), 8.18 (1H, dd, $J = 8.3, 1.3$ Hz), 9.02 (1H, dd, $J = 4.2, 1.3$
10
11 Hz). ¹³C-NMR (CDCl₃): δ 26.1 (CH₂), 28.2 (CH₂), 28.7 (CH₂), 29.0 (CH₂), 29.4 (CH₂) (x2), 32.8
12
13 (CH₂), 34.1 (CH₂), 69.0 (CH₂), 108.6 (C), 119.3 (C), 121.6 (C), 126.8 (C), 129.5 (C), 136.2 (C), 140.0
14
15 (C), 149.2 (C), 154.8 (C).
16
17
18
19

20 8-((12-bromododecyl)oxy)quinoline (**7**): Yield 67% (0.67 mmol, 263 mg); brown solid, M.p. 44-46°C.
21
22 IR (cm⁻¹): ν_{\max} 1597 (C=C_{Ar}), 1529 (C=N), 1259 (C-O-C), 829 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.23-
23
24 1.38 (6H, m), 1.37-1.49 (2H, m), 1.49-1.62 (2H, m), 1.79-1.93 (2H, m), 1.98-2.12 (2H, m), 3.42 (2H, t,
25
26 $J = 6.9$ Hz), 4.25 (2H, t, $J = 7.0$ Hz), 7.07 (1H, d, $J = 7.5$ Hz), 7.35-7.51 (3H, m), 8.13 (1H, dd, $J = 8.3,$
27
28 1.2 Hz), 8.98 (1H, dd, $J = 4.2, 1.2$ Hz). ¹³C-NMR (CDCl₃): δ 26.1 (CH₂), 28.2 (CH₂), 28.8 (CH₂), 29.0
29
30 (CH₂), 29.6 (CH₂) (x3), 29.5 (CH₂), 29.4 (CH₂) (x2), 32.9 (CH₂), 34.1 (CH₂), 69.0 (CH₂), 108.6 (C),
31
32 119.3 (C), 121.5 (C), 126.8 (C), 129.5 (C), 136.0 (C), 140.3 (C), 149.2 (C), 154.9 (C).
33
34
35
36
37

38 7-[(9-bromononyl)oxy]-4H-chromen-4-one (**18**): Yield 49% (0.49 mmol, 180 mg); light yellow solid,
39
40 M.p. 60-62°C. IR (cm⁻¹): ν_{\max} 1649 (C=O), 1602 (C=C), 1444 (C=C_{Ar}), 1234 (C-O-C), 813 (C-H_{Ar}).
41
42 ¹H-NMR (CDCl₃): δ 1.28-1.42 (10H, m), 1.42-1.55 (2H, m), 1.78-1.93 (2H, m), 3.43 (2H, t, $J = 6.8$
43
44 Hz), 4.05 (2H, t, $J = 6.5$ Hz), 6.29 (1H, d, $J = 6.0$ Hz), 6.83 (1H, d, $J = 2.3$ Hz), 6.97 (1H, dd, $J = 9.0,$
45
46 2.3 Hz), 7.80 (1H, d, $J = 6.1$ Hz), 8.11 (1H, d, $J = 9.0$ Hz). ¹³C-NMR (CDCl₃): δ 25.9 (CH₂), 28.1
47
48 (CH₂), 28.7 (CH₂), 28.9 (CH₂), 29.2 (CH₂), 29.3 (CH₂), 32.8 (CH₂), 34.1 (CH₂), 68.7 (CH₂), 100.8 (C),
49
50 112.9 (C), 114.9 (C), 118.6 (C), 127.1 (C), 154.9 (C), 158.3 (C), 163.7 (C), 177.1 (C=O).
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 7-[(12-bromododecyl)oxy]-4H-chromen-4-one (**19**): Yield 42% (0.42 mmol, 172 mg); light yellow
4
5 solid, M.p. 58-60°C. IR (cm⁻¹): ν_{\max} 1651 (C=O), 1605 (C=C), 1450 (C=C_{Ar}), 1238 (C-O-C), 812 (C-
6
7 H_{Ar}). ¹H-NMR (CDCl₃): δ 1.22-1.39 (12H, m), 1.40-1.52 (4H, m), 1.79-1.90 (4H, m), 3.41 (2H, t, *J* =
8
9 6.8 Hz), 4.05 (2H, t, *J* = 6.5 Hz), 6.30 (1H, d, *J* = 6.0 Hz), 6.83 (1H, d, *J* = 2.2 Hz), 6.97 (1H, dd, *J* =
10
11 8.9, 2.2 Hz), 7.77 (1H, d, *J* = 6.0 Hz), 8.11 (1H, d, *J* = 8.9 Hz). ¹³C-NMR (CDCl₃): δ 25.9 (CH₂), 28.1
12
13 (CH₂), 28.7 (CH₂), 28.9 (CH₂), 29.3 (CH₂), 29.4 (CH₂), 29.5 (CH₂), 29.5 (CH₂) (x2), 32.8 (CH₂), 33.9
14
15 (CH₂), 68.7 (CH₂), 100.9 (C), 112.9 (C), 114.8 (C), 118.6 (C), 127.1 (C), 154.8 (C), 158.3 (C), 163.7
16
17 (C), 177.1 (C=O).

22 *General procedure for the synthesis of quinoline-chalcone and quinoline-chromone hybrids*

25 Chalcone or 8-hydroxyquinoline (0.75 mmol), potassium hydroxide (1 mmol) and acetonitrile (10
26
27 mL), were placed in a 50 mL flat-bottomed flask equipped with a magnetic stirring bar. The mixture
28
29 was stirred and heated to reflux for a period of 5 minutes, under microwave irradiation. Then,
30
31 bromoalkylquinoline or bromoalkylchromenone (0.5 mmol) was added to the reaction mixture which
32
33 was then refluxed for 30 minutes (70 W). The crude reaction mixture was concentrated on a rotatory
34
35 evaporator and the residue was purified by column chromatography over silica gel eluting with hexane-
36
37 ethyl acetate to obtain quinoline-chalcone or quinoline-chromone hybrids in yields ranging 44-65% and
38
39 34-70%, respectively. Monitoring of the reaction progress and product purification was carried out by
40
41 TLC.
42
43
44
45
46
47

48 (*E*)-3-(3,4-dimethoxyphenyl)-1-(4-(3-(quinolin-8-yloxy)propoxy)phenyl)prop-2-en-1-one (**8**): Yield
49
50 65% (0.33 mmol, 155 mg); yellow solid, M.p. 62-64°C; IR (cm⁻¹): ν_{\max} 1655 (C=O), 1599 (C=C),
51
52 1510, (C=N) 1262 (C-O-C), 803 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 2.51-2.59 (2H, m), 3.94 (3H, s), 3.96
53
54 (3H, s), 4.37 (2H, t, *J* = 6.1 Hz), 4.48 (2H, t, *J* = 6.2 Hz), 6.90 (1H, d, *J* = 8.3 Hz), 7.01 (2H, d, *J* = 8.8
55
56 Hz), 7.13 (1H, dd, *J* = 7.6, 1.0 Hz), 7.16 (1H, d, *J* = 2.0 Hz), 7.23 (1H, dd, *J* = 8.3, 2.0), 7.37-7.49 (4H,
57
58 m), 7.75 (1H, d, *J* = 15.6 Hz), 8.01 (2H, d, *J* = 8.8 Hz), 8.14 (1H, dd, *J* = 8.3, 1.7 Hz), 8.96 (1H, dd, *J* =
59
60
61
62
63
64
65

1
2
3 4.4, 1.7 Hz). ¹³C-NMR (CDCl₃): δ 29.1 (CH₂), 56.0 (CH₃) (x2), 65.0 (CH₂), 65.4 (CH₂), 109.0 (C),
4
5 110.2 (C), 111.2 (C), 114.3 (C) (x2), 119.8 (C), 119.9 (C), 121.6 (C), 122.9 (C), 126.6 (C), 128.1 (C),
6
7 129.5 (C), 130.7 (C) (x2), 131.3 (C), 135.9 (C), 140.4 (C), 144.0 (C), 149.2 (C), 149.3 (C), 151.3 (C),
8
9 154.6 (C), 162.6 (C), 188.6 (C=O). ESI-MS: *m/z* 470.1967 [M + H]⁺, Calcd for C₂₉H₂₇NO₅ : 470.1959

10
11
12
13 (*E*)-3-(3,4-dimethoxyphenyl)-1-(4-(4-(quinolin-8-yloxy)butoxy)phenyl)prop-2-en-1-one (**9**): Yield 44%
14
15 (0.22 mmol, 106.4 mg); yellow solid, M.p. 130-134°C; IR (cm⁻¹): ν_{max} 1659 (C=O), 1599 (C=C), 1512
16
17 (C=N), 1262 (C-O-C), 814 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 2.08-2.17 (2H, m), 2.20-2.29 (2H, m), 3.94
18
19 (3H, s), 3.97 (3H, s), 4.21 (2H, t, *J* = 6.3 Hz), 4.36 (2H, t, *J* = 6.4 Hz), 6.91 (1H, d, *J* = 8.3 Hz), 6.99
20
21 (2H, d, *J* = 8.8 Hz), 7.09 (1H, d, *J* = 7.6 Hz), 7.17 (1H, d, *J* = 1.8 Hz), 7.24 (1H, dd, *J* = 8.3, 1.7), 7.37-
22
23 7.50 (4H, m), 7.76 (1H, d, *J* = 15.6 Hz), 8.01 (2H, d, *J* = 8.9 Hz), 8.13 (1H, dd, *J* = 8.3, 1.9 Hz), 8.95
24
25 (1H, dd, *J* = 4.2, 1.9 Hz). ¹³C-NMR (CDCl₃): δ 25.7 (CH₂), 26.1 (CH₂), 56.0 (CH₃) (x2), 67.9 (CH₂),
26
27 68.5 (CH₂), 108.8 (C), 110.2 (C), 111.2 (C), 114.3 (C) (x2), 119.6 (C), 120.0 (C), 121.6 (C), 122.9 (C),
28
29 126.6 (C), 128.1 (C), 129.5 (C), 130.7 (C) (x2), 131.2 (C), 135.9 (C), 140.5 (C), 144.0 (C), 149.2 (C),
30
31 149.3 (C), 151.3 (C), 154.7 (C), 162.8 (C), 188.7 (C=O). ESI-MS: *m/z* 484.2124 [M + H]⁺, Calcd for
32
33 C₃₀H₂₉NO₅ : 484.2119
34
35
36
37
38
39
40

41 (*E*)-3-(3,4-dimethoxyphenyl)-1-(4-((5-(quinolin-8-yloxy)pentyl)oxy)phenyl)prop-2-en-1-one (**10**): Yield
42
43 56% (0.28 mmol, 139.3 mg); yellow solid, M.p. 220-224°C; IR (cm⁻¹): ν_{max} 1641 (C=O), 1593 (C=C),
44
45 1513 (C=N), 1221 (C-O-C), 812 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.73-1.88 (2H, m), 1.92-2.05 (2H, m),
46
47 2.11-2.24 (2H, m), 3.94 (3H, s), 4.01 (3H, s), 4.13 (2H, t, *J* = 6.4 Hz), 4.32 (2H, t, *J* = 6.8 Hz), 6.94
48
49 (1H, d, *J* = 8.5 Hz), 7.01 (2H, d, *J* = 8.7 Hz), 7.11 (1H, d, *J* = 7.5 Hz), 7.21 (1H, d, *J* = 1.5 Hz), 7.27
50
51 (1H, dd, *J* = 8.5, 1.5), 7.37-7.53 (4H, m), 7.80 (1H, d, *J* = 15.5 Hz), 8.06 (2H, d, *J* = 8.7 Hz), 8.17 (1H,
52
53 dd, *J* = 8.2, 1.1 Hz), 8.89 (1H, dd, *J* = 4.0, 1.1 Hz). ¹³C-NMR (CDCl₃): δ 22.8 (CH₂), 28.8 (CH₂), 29.0
54
55 (CH₂), 56.0 (CH₃) (x2), 68.0 (CH₂), 68.7 (CH₂), 108.7 (C), 110.1 (C), 111.2 (C), 114.3 (C) (x2), 119.6
56
57 (C), 119.9 (C), 121.6 (C), 123.0 (C), 126.7 (C), 128.1 (C), 129.6 (C), 130.8 (C) (x2), 131.1 (C), 136.0
58
59 (C), 151.3 (C), 154.7 (C), 162.8 (C), 188.7 (C=O). ESI-MS: *m/z* 484.2124 [M + H]⁺, Calcd for
60
61 C₃₀H₂₉NO₅ : 484.2119
62
63
64
65

(C), 140.4 (C), 144.1 (C), 149.2 (C), 149.4 (C), 151.3 (C), 154.8 (C), 162.9 (C), 188.9 (C=O). ESI-MS: m/z 498.2280 $[M + H]^+$, Calcd for $C_{31}H_{31}NO_5$: 498.2287

(E)-3-(3,4-dimethoxyphenyl)-1-(4-((8-(quinolin-8-yloxy)octyl)oxy)phenyl)prop-2-en-1-one (**11**): Yield 58% (0.29 mmol, 156.5 mg); yellow solid, M.p. 104-108°C; IR (cm^{-1}): ν_{max} 1657 (C=O), 1599 (C=C), 1513 (C=N), 1258 (C-O-C), 816 (C-H_{Ar}). 1H -NMR ($CDCl_3$): δ 1.40-1.48 (2H, m), 1.48-1.54 (2H, m), 1.54-1.62 (2H, m), 1.79-1.86 (2H, m), 2.01-2.09 (2H, m), 3.94 (3H, s), 3.96 (3H, s), 4.04 (2H, t, $J = 6.5$ Hz), 4.26 (2H, t, $J = 7.0$ Hz), 6.90 (1H, d, $J = 8.4$ Hz), 6.97 (2H, d, $J = 8.8$ Hz), 7.07 (1H, d, $J = 7.6$ Hz), 7.17 (1H, d, $J = 1.9$ Hz), 7.24 (1H, dd, $J = 8.4, 2.0$), 7.36-7.48 (4H, m), 7.76 (1H, d, $J = 15.6$ Hz), 8.03 (2H, d, $J = 8.9$ Hz), 8.12 (1H, dd, $J = 8.3, 1.7$ Hz), 8.96 (1H, dd, $J = 4.1, 2.0$ Hz). ^{13}C -NMR ($CDCl_3$): δ 25.9 (CH₂), 26.0 (CH₂), 29.0 (CH₂), 29.1 (CH₂), 29.2 (CH₂), 29.3 (CH₂), 56.0 (CH₃) (x2), 68.2 (CH₂), 69.0 (CH₂), 108.6 (C), 110.2 (C), 111.2 (C), 114.3 (C) (x2), 119.4 (C), 119.9 (C), 121.5 (C), 122.9 (C), 126.6 (C), 128.1 (C), 129.5 (C), 130.7 (C) (x2), 131.1 (C), 135.8 (C), 140.5 (C), 144.0 (C), 149.2 (C), 149.3 (C), 151.3 (C), 154.7 (C), 162.9 (C), 188.7 (C=O). ESI-MS: m/z 540.2750 $[M + H]^+$, Calcd for $C_{34}H_{37}NO_5$: 540.2742.

(E)-3-(3,4-dimethoxyphenyl)-1-(4-((9-(quinolin-8-yloxy)nonyl)oxy)phenyl)prop-2-en-1-one (**12**): Yield 49% (0.23 mmol, 127.4 mg); yellow solid, M.p. 84-88°C; IR (cm^{-1}): ν_{max} 1728 (C=O), 1655 (C=C), 1601 (C=N), 1262 (C-O-C), 797 (C-H_{Ar}). 1H -NMR ($CDCl_3$): δ 1.35-1.51 (8H, m), 1.52-1.65 (2H, m), 1.76-1.90 (2H, m), 2.01-2.13 (2H, m), 3.95 (3H, s), 3.98 (3H, s), 4.05 (2H, t, $J = 6.5$ Hz), 4.26 (2H, t, $J = 7.1$ Hz), 6.91 (1H, d, $J = 8.3$ Hz), 6.99 (2H, d, $J = 8.8$ Hz), 7.08 (1H, d, $J = 7.5$ Hz), 7.19 (1H, d, $J = 1.5$ Hz), 7.26 (1H, dd, $J = 8.3, 1.6$), 7.36-7.51 (4H, m), 7.79 (1H, d, $J = 15.5$ Hz), 8.06 (2H, d, $J = 8.8$ Hz), 8.14 (1H, dd, $J = 8.3, 1.4$ Hz), 8.97 (1H, dd, $J = 4.2, 1.5$ Hz). ^{13}C -NMR ($CDCl_3$): δ 26.0 (CH₂), 26.1 (CH₂), 29.0 (CH₂), 29.1 (CH₂), 29.3 (CH₂), 29.4 (CH₂), 29.5 (CH₂), 56.0 (CH₃) (x2), 68.3 (CH₂), 69.0 (CH₂), 108.6 (C), 110.1 (C), 111.1 (C), 114.3 (C) (x2), 119.4 (C), 119.9 (C), 121.6 (C), 123.0 (C), 126.7 (C), 128.1 (C), 129.5 (C), 130.8 (C) (x2), 131.6 (C), 135.9 (C), 140.4 (C), 144.0 (C), 149.2 (C),

1
2
3 149.3 (C), 151.2 (C), 154.9 (C), 163.0 (C), 188.8 (C=O). ESI-MS: m/z 554.2987 [M + H]⁺, Calcd for
4
5 C₃₅H₃₉NO₅ : 554.2991.
6

7
8
9 (*E*)-3-(3,4-dimethoxyphenyl)-1-(4-((12-(quinolin-8-yloxy)dodecyl)oxy)phenyl)prop-2-en-1-one (13):

10 Yield 51% (0.26 mmol, 155 mg); yellow solid, M.p. 98-102°C; IR (cm⁻¹): ν_{\max} 1656 (C=O), 1599
11 (C=C), 1503 (C=N), 1258 (C-O-C), 801 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.27-1.49 (14H, m), 1.50-1.64
12 (2H, m), 1.75-1.93 (2H, m), 2.01-2.15 (2H, m), 3.98 (3H, s), 4.01 (3H, s), 4.08 (2H, t, J = 6.7 Hz), 4.28
13 (2H, t, J = 7.1 Hz), 6.94 (1H, d, J = 8.4 Hz), 7.02 (2H, d, J = 8.8 Hz), 7.10 (1H, d, J = 7.5 Hz), 7.21
14 (1H, d, J = 1.5 Hz), 7.28 (1H, dd, J = 8.3, 1.7), 7.38-7.53 (4H, m), 7.80 (1H, d, J = 15.7 Hz), 8.07 (2H,
15 d, J = 8.8 Hz), 8.16 (1H, dd, J = 8.3, 1.6 Hz), 8.99 (1H, dd, J = 4.2, 1.6 Hz). ¹³C-NMR (CDCl₃): δ 26.0
16 (CH₂), 26.1 (CH₂), 29.0 (CH₂), 29.1 (CH₂), 29.4 (CH₂), 29.5 (CH₂), 29.6 (CH₂) (x4), 56.0 (CH₃) (x2),
17 68.3 (CH₂), 69.0 (CH₂), 108.6 (C), 110.1 (C), 111.1 (C), 114.3 (C) (x2), 119.4 (C), 119.9 (C), 121.6
18 (C), 123.0 (C), 126.7 (C), 128.1 (C), 129.5 (C), 130.8 (C) (x2), 131.1 (C), 135.9 (C), 140.5 (C), 144.1
19 (C), 149.2 (C), 149.3 (C), 151.2 (C), 154.9 (C), 163.0 (C), 188.8 (C=O). ESI-MS: m/z 596.3410 [M +
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 H]⁺, Calcd for C₃₈H₄₅NO₅ : 596.3404.
36
37

38
39 7-[4-(quinolin-8-yloxy)butoxy]-4H-chromen-4-one (20): Yield 34% (0.17 mmol, 61.4 mg); yellow
40
41 solid, M.p. 160-164°C; IR (cm⁻¹): ν_{\max} 1641 (C=O), 1593 (C=C), 1565 (C=N), 1437 (C=C_{Ar}), 1267 (C-
42
43 O-C), 820 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 2.10-2.19 (2H, m), 2.20-2.29 (2H, m), 4.23 (2H, t, J = 6.2 Hz),
44
45 4.36 (2H, t, J = 6.2 Hz), 6.27 (1H, d, J = 6.0 Hz), 6.85 (1H, d, J = 2.3 Hz), 6.94 (1H, dd, J = 9.0, 2.3
46
47 Hz), 7.08 (1H, d, J = 7.6 Hz), 7.36-7.48 (3H, m), 7.76 (1H, d, J = 6.0 Hz), 8.07 (1H, d, J = 9.0 Hz),
48
49 8.12 (1H, dd, J = 8.3, 1.5 Hz), 8.94 (1H, dd, J = 4.1, 1.5 Hz). ¹³C-NMR (CDCl₃): δ 25.6 (CH₂), 26.2
50
51 (CH₂), 68.4 (CH₂), 68.5 (CH₂), 100.1 (C), 108.7 (C), 112.9 (C), 114.8 (C), 118.6 (C), 119.7 (C), 121.5
52
53 (C₁), 126.6 (C), 127.1 (C), 129.5 (C), 135.8 (C), 140.4 (C), 149.3 (C), 154.6 (C), 154.7 (C), 158.2 (C),
54
55
56
57 163.5 (C), 176.9 (C=O). ESI-MS: m/z 362.1392 [M + H]⁺, Calcd for C₂₂H₁₉NO₄: 362.1390
58
59
60
61
62
63
64
65

1
2
3 7-*{[5-(quinolin-8-yloxy)pentyl]oxy}*-4*H-chromen-4-one* (**21**): Yield 61% (0.31 mmol, 116.4 mg);
4
5 yellow solid, M.p. 134-138°C; IR (cm⁻¹): ν_{\max} 1651 (C=O), 1597 (C=C), 1564 (C=N), 1443 (C=C_{Ar}),
6
7 1263 (C-O-C), 824 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.73-1.81 (2H, m), 1.92-2.01 (2H, m), 2.08-2.17 (2H,
8
9 m), 4.02 (2H, t, *J* = 6.4 Hz), 4.29 (2H, t, *J* = 6.7 Hz), 6.26 (1H, d, *J* = 6.1 Hz), 6.81 (1H, d, *J* = 2.4 Hz),
10
11 6.94 (1H, dd, *J* = 8.8, 2.4 Hz), 7.07 (1H, d, *J* = 7.8 Hz), 7.36-7.47 (3H, m), 7.76 (1H, d, *J* = 6.1 Hz),
12
13 8.09 (1H, d, *J* = 8.8 Hz), 8.12 (1H, dd, *J* = 8.3, 1.6 Hz), 8.94 (1H, dd, *J* = 4.2, 1.6 Hz). ¹³C-NMR
14
15 (CDCl₃): δ 22.8 (CH₂), 28.7 (CH₂), 28.8 (CH₂), 68.4 (CH₂), 68.7 (CH₂), 100.9 (C), 108.7 (C), 112.9
16
17 (C), 114.8 (C), 118.6 (C), 119.6 (C), 121.5 (C), 126.6 (C), 127.1 (C), 129.5 (C), 135.8 (C), 140.5 (C),
18
19 149.3 (C), 154.7 (C), 154.8 (C), 158.2 (C), 163.6 (C), 177.0 (C=O). ESI-MS: *m/z* 376.1549 [M + H]⁺,
20
21 Calcd for C₂₃H₂₁NO₄: 376.1542.
22
23
24
25
26

27 7-*{[8-(quinolin-8-yloxy)octyl]oxy}*-4*H-chromen-4-one* (**22**): Yield 70% (0.35 mmol, 146.1 mg); yellow
28
29 solid, M.p. 110-114°C; IR (cm⁻¹): ν_{\max} 1655 (C=O), 1593 (C=C), 1593 (C=N), 1443 (C=C_{Ar}), 1263 (C-
30
31 O-C), 818 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.42-1.54 (2H, m), 1.55-1.67 (2H, m), 1.80-1.93 (2H, m),
32
33 2.02-2.19 (2H, m), 4.07 (2H, t, *J* = 6.6 Hz), 4.28 (2H, t, *J* = 7.0 Hz), 6.31 (1H, d, *J* = 6.1 Hz), 6.85 (1H,
34
35 d, *J* = 2.3 Hz), 7.01 (1H, dd, *J* = 8.9, 2.3 Hz), 7.10 (1H, d, *J* = 7.5 Hz), 7.38-7.53 (3H, m), 7.80 (1H, d,
36
37 *J* = 6.0 Hz), 8.13 (1H, d, *J* = 8.7 Hz), 8.16 (1H, dd, *J* = 8.0, 1.4 Hz), 8.99 (1H, dd, *J* = 4.2, 1.4 Hz). ¹³C-
38
39 NMR (CDCl₃): δ 25.9 (CH₂), 26.0 (CH₂), 28.9 (CH₂), 29.0 (CH₂), 29.2 (CH₂), 29.3 (CH₂), 68.7 (CH₂),
40
41 68.9 (CH₂), 100.8 (C), 108.6 (C), 112.9 (C), 114.9 (C), 118.6 (C), 119.4 (C), 121.6 (C), 126.7 (C),
42
43 127.2 (C), 129.5 (C), 136.0 (C), 140.4 (C), 149.3 (C), 154.7 (C), 154.9 (C), 158.3 (C), 163.7 (C), 177.1
44
45 (C=O). ESI-MS: *m/z* 418.2127 [M + H]⁺, Calcd for C₂₆H₂₇NO₄: 418.2123.
46
47
48
49
50

51 7-*{[9-(quinolin-8-yloxy)nonyl]oxy}*-4*H-chromen-4-one* (**23**): Yield 36% (0.18 mmol, 77.7 mg); yellow
52
53 solid, M.p. 93-95°C; IR (cm⁻¹): ν_{\max} 1659 (C=O), 1626 (C=C), 1594 (C=N), 1381 (C=C_{Ar}), 1265 (C-
54
55 O-C), 818 (C-H_{Ar}). ¹H-NMR (CDCl₃): δ 1.31-1.45 (6H, m), 1.45-1.51 (2H, m), 1.51-1.60 (2H, m),
56
57 1.78-1.86 (2H, m), 2.01-2.08 (2H, m), 4.03 (2H, t, *J* = 6.5 Hz), 4.25 (2H, t, *J* = 6.9 Hz), 6.27 (1H, d, *J* =
58
59
60
61
62
63
64
65

1
2
3 6.0 Hz), 6.82 (1H, d, $J = 2.3$ Hz), 6.96 (1H, dd, $J = 8.9, 2.3$ Hz), 7.07 (1H, d, $J = 7.8$ Hz), 7.36-7.48
4
5 (3H, m), 7.76 (1H, d, $J = 6.0$ Hz), 8.11 (1H, dd, $J = 8.1, 1.0$ Hz), 8.13 (1H, d, $J = 8.5$ Hz), 8.96 (1H, dd,
6
7 $J = 4.2, 1.4$ Hz). ^{13}C -NMR (CDCl_3): δ 25.9 (CH_2), 26.0 (CH_2), 28.9 (CH_2), 29.0 (CH_2), 29.2 (CH_2),
8
9 29.3 (CH_2), 29.4 (CH_2), 68.7 (CH_2), 69.0 (CH_2), 100.8 (C), 108.6 (C), 112.9 (C), 114.8 (C), 118.6 (C),
10
11 119.3 (C), 121.5 (C), 126.6 (C), 127.1 (C), 129.5 (C), 135.9 (C), 140.4 (C), 149.2 (C), 154.7 (C), 154.9
12
13 (C), 158.2 (C), 163.7 (C), 177.0 (C=O). ESI-MS: m/z 432.2175 $[\text{M} + \text{H}]^+$, Calcd for $\text{C}_{27}\text{H}_{29}\text{NO}_4$:
14
15 432.2173.
16
17

18
19
20 7-[[12-(quinolin-8-yloxy)dodecyl]oxy]-4H-chromen-4-one (**24**): Yield 61% (0.31 mmol, 146.8 mg);
21
22 yellow solid, M.p. 98-100°C; IR (cm^{-1}): ν_{max} 1651 (C=O), 1622 (C=C), 1596 (C=N), 1441 (C=C_{Ar}),
23
24 1263 (C-O-C), 817 (C-H_{Ar}). ^1H -NMR (CDCl_3): δ 1.24-1.43 (12H, m), 1.44-1.50 (2H, m), 1.51-1.60
25
26 (2H, m), 1.78-1.87 (2H, m), 1.99-2.08 (2H, m), 4.04 (2H, t, $J = 6.6$ Hz), 4.24 (2H, t, $J = 7.1$ Hz), 6.27
27
28 (1H, d, $J = 6.1$ Hz), 6.82 (1H, d, $J = 2.1$ Hz), 6.96 (1H, dd, $J = 8.8, 2.0$ Hz), 7.06 (1H, d, $J = 7.7$ Hz),
29
30 7.34-7.48 (3H, m), 7.76 (1H, d, $J = 6.1$ Hz), 8.10 (1H, dd, $J = 8.1, 1.2$ Hz), 8.12 (1H, d, $J = 8.2$ Hz),
31
32 8.95 (1H, dd, $J = 4.2, 1.2$ Hz). ^{13}C -NMR (CDCl_3): δ 25.9 (CH_2), 26.0 (CH_2), 28.9 (CH_2), 29.0 (CH_2),
33
34 29.3 (CH_2), 29.41 (CH_2), 29.47 (CH_2), 29.48 (CH_2), 29.49 (CH_2), 29.5 (CH_2), 68.7 (CH_2), 69.0 (CH_2),
35
36 100.9 (C), 108.6 (C), 112.9 (C), 114.8 (C), 118.6 (C), 119.3 (C), 121.5 (C), 126.6 (C), 127.1 (C), 129.5
37
38 (C), 135.8 (C), 140.5 (C), 149.2 (C), 154.7 (C), 154.9 (C), 158.3 (C), 163.7 (C), 177.0 (C=O). ESI-MS:
39
40 m/z 474.2644 $[\text{M} + \text{H}]^+$, Calcd for $\text{C}_{30}\text{H}_{35}\text{NO}_4$: 474.2642.
41
42
43
44
45
46
47
48

49 Biological activity assays

50
51 The compounds were subjected to evaluation of *in vitro* cytotoxicity on U937 human cells and
52
53 leishmanicidal and trypanocidal activities on intracellular amastigotes of *L. (V) panamensis* and *T.*
54
55 *cruzi*.
56
57
58
59
60
61
62
63
64
65

In vitro Cytotoxicity

The cytotoxic activity of the compounds was assessed based on the viability of the human promonocytic cell line U-937 (ATCC CRL-1593.2TM) evaluated by the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay following the methodology described previously (Taylor *et al.*, 2011). Briefly, cells grown in tissue flasks were harvested and washed with phosphate buffered saline (PBS) by centrifuging. Cells were counted and adjusted at 1×10^6 cells/mL of RPMI-1640 supplemented with complete 10% Fetal Bovine Serum (FBS) and 1% antibiotics (100 U/mL penicillin and 0.1 mg/mL streptomycin). One hundred μ L were dispensed into each well of a 96-well cell-culture plate and then 100 μ L of RPMI-1640 and the corresponding concentrations of the compounds were added, starting at 200 μ g/mL in duplicate. Plates were incubated at 37 °C, 5% CO₂ during 72 h in the presence of extracts. The effect of compounds was determined by measuring the activity of the mitochondrial dehydrogenase by adding 10 μ L/well of MTT solution (0.5 mg/mL) and incubation at 37 °C for 3h. The reaction was stopped by adding 100 μ L/well of 50% isopropanol solution with 10% sodium dodecyl sulfate and 30 min incubation. Cell viability was determined based on the quantity of formazan produced according to the intensity of color (absorbance) registered as optical densities (O.D) obtained at 570 nm in a spectrophotometer (VarioskanTM Flash Multimode Reader - Thermo Scientific, USA). Cells cultured in absence of compounds were used as control of viability (negative control), while meglumine antimoniate (Sbv) and amphotericin B (AmB) were used as control for cytotoxicity (non-cytotoxic and cytotoxic drugs, respectively). Assays were conducted in two independent runs with three replicates per each concentration tested.

In vitro leishmanicidal activity

The activity of compounds was evaluated on intracellular amastigotes of *L. (V) panamensis* transfected with the green fluorescent protein gene (MHOM/CO/87/UA140pIR-GFP) (Pulido *et al.*,

1
2
3 2012). The effect of each extract was determined according to the inhibition of the infection evidenced
4
5 by both decrease of the infected cells and decrease of intracellular parasite load. Briefly, U-937 human
6
7 cells at a concentration of 3×10^5 cells/mL in RPMI 1640 and 0.1 $\mu\text{g/mL}$ of phorbol-12-myristate-13-
8
9 acetate (PMA) were dispensed into each well of a 24-well cell culture plate and then infected with 5
10
11 days-old promastigotes in a 15:1 parasites per cell ratio. Plates were incubated at 34 °C, 5% CO₂ during
12
13 3h and cells were washed two times with PBS to eliminate not internalized parasites. One mL of fresh
14
15 RPMI 1640 supplemented with 10% FBS and 1% antibiotics was added into each well, cells were
16
17 incubated again to guarantee multiplication of intracellular parasites. After 24 h of infection, culture
18
19 medium was replaced by fresh culture medium containing each compound at 20 $\mu\text{g/mL}$ or lower (based
20
21 on the cytotoxicity showed previously by each compound) and plates were incubated at 37 °C, 5%
22
23 CO₂. After 72 h, inhibition of the infection was determined. For this, cells were removed from the
24
25 bottom plate with a trypsin/EDTA (250 mg) solution; recovered cells were centrifuged at 1100 rpm
26
27 during 10 min at 4 °C, the supernatant was discarded and cells were washed with 1 mL of cold PBS
28
29 and centrifuged at 1100 rpm during 10 min at 4 °C. The supernatant was discarded and cells were
30
31 suspended in 500 μL of PBS and analyzed by flow cytometry (FC 500MPL, Cytomics, Brea, CA, US.
32
33 All determinations for each extract and standard drugs were carried out in triplicate, in two independent
34
35 experiments (Buckner *et al.*, 1996; Pulido *et al.*, 2012). Activity of tested extracts was carried out in
36
37 parallel with infection progress in culture medium alone and in culture medium with AmB and SbV as
38
39 antileishmanial drugs (positive controls). Compounds that showed percentages of inhibition higher than
40
41 50% to 20 or fewer $\mu\text{g/mL}$ were then evaluated at four additional concentrations to determine the
42
43 effective concentration 50 (EC₅₀). Here, infected cells were exposed against each concentration of
44
45 compounds during 72 h; then, cells were removed and tested by flow cytometry as described before.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In vitro Trypanocidal Activity

Compounds were tested on intracellular amastigotes of *T. cruzi*, Tulahuen strain transfected with β -galactosidase gene (donated by Dr. F. S. Buckner, University of Washington) (Buckner et al., 1996).

The activity was determined according to the ability of the extract to reduce the infection of U-937 cells by *T. cruzi* as described elsewhere (Insuasty *et al.*, 2015). Following the procedure described above, anti-*T. cruzi* activity was initially screened at a single concentration of 20 mg/mL. In this case, 100 μ L of U-937 human cells at a concentration of 2.5×10^5 cells/mL in RPMI-1640, 10% SFB and 0.1 μ g/mL of PMA were placed in each well of 96-well plates and then infected with phase growth epimastigotes in 5:1 (parasites per cell) ratio and incubated at 34 °C, 5% CO₂. After 24 hours of incubation, 20 μ g/mL of each extract were added to infected cells. After 72 h of incubation, the effect of all extracts on viability of intracellular amastigotes was determined by measuring the β -galactosidase activity by spectrophotometry adding 100 μ M CPRG and 0.1% nonidet P-40 to each well. After 3 h of incubation, plates were read at 570 nm in a spectrophotometer (Varioskan™ Flash Multimode Reader - Thermo Scientific, USA) and intensity of color (absorbance) was registered as O.D. Extracts that showed inhibition percentages higher than 50% were evaluated again at four concentrations selected according to the LC₅₀ previously obtained for each compound. Infected cells exposed to benznidazol (BNZ) were used as control for antitrypanosomal activity (positive control) while infected cells incubated in culture medium alone were used as control for infection (negative control). Non-specific absorbance was corrected by subtracting the O.D of the blank. Determinations were done by triplicate in at least two independent experiments (Insuasty *et al.*, 2015).

Statistical Analysis

Cytotoxicity was determined according to the percentages of viability and mortality registered to each compound an concentration, including amphotericin B, meglumine antimoniate and culture medium alone. Percentage of viability was calculated by Equation 1, where the O.D of control, corresponds to 100% of viability.

$$\% \text{ Viability} = (\text{O.D Exposed cells}) / (\text{O.D Control cells}) \times 100 \quad (1)$$

In turn, mortality percentage corresponds to 100%–% viability.

Results were expressed as 50 lethal concentrations (LC₅₀) that corresponds to the concentration necessary to eliminate 50% of cells and calculated by Probit analysis (Finney, 1978). The degree of toxicity was graded according to the LC₅₀ value using the following scale: high cytotoxicity: LC₅₀ < 100 µg/mL; moderate cytotoxicity: LC₅₀ > 100 to < 200 µg/mL and potentially non-cytotoxicity: LC₅₀ > 200 µg/mL.

On the other hand, anti-leishmanial activity was determined according to the percentage of infected cells and parasite load, obtained for each experimental condition by flow cytometry. The percentage of infected cells was determined as the number of positive events evidenced by green fluorescence (parasites) and Forward Scatter (FSC) using dotplot analysis, while, the parasitic load was determined by analysis of mean fluorescence intensity (MFI) of fluorescent parasites (Pulido *et al.*, 2012). The parasite inhibition was calculated by equation 2, where the MFI of control, corresponds to 100% of parasites.

$$\% \text{ Parasite} = (\text{MFI Exposed parasites}) / (\text{MFI Control parasites}) \times 100 \quad (2)$$

In turn, inhibition percentage corresponds to 100% – % Parasites. Results of leishmanicidal activity were expressed as EC₅₀ determined by the Probit analysis (Finney, 1978).

1
2
3 Similarly, trypanocidal activity was determined according to the percentage of infected cells and
4
5 parasite load obtained for each experimental condition by colorimetry. Parasite inhibition was
6
7 calculated by equation 3, where the O.D of control corresponds to 100% of infection.
8
9

$$\% \text{ Infection} = (\text{O.D Exposed parasites}) / (\text{O.D Control parasites}) \times 100 \quad (3)$$

10
11
12 In turn, percentage of inhibition of infection corresponds to $100\% - \% \text{ of Infection}$.
13

14
15 Results of anti-leishmanial and anti-trypanocidal activities were expressed as EC_{50} determined by the
16
17 Probit analysis (Finney, 1978). The leishmanicidal or trypanocidal activities were graded according to
18
19 the EC_{50} value using the following scale: High activity: $EC_{50} < 25 \mu\text{g/mL}$, moderate activity: $EC_{50} > 25$
20
21 to $< 50 \mu\text{g/mL}$; potentially non activity: $EC_{50} > 50 \mu\text{g/mL}$.
22
23
24


25
26 The selectivity index (SI), was calculated by dividing the cytotoxic activity and the leishmanicidal
27
28 or trypanocidal activity using the following formula: $SI = CL_{50}/CE_{50}$. Cytotoxic compound: $LC_{50} < 100$
29
30 $\mu\text{g/mL}$.
31
32

33 34 35 **Results and discussion**

36 37 38 **Chemistry**

39
40
41
42 Quinoline-chalcone hybrids **8-13** were obtained via microwave assisted Williamson etherification
43
44 (Peng *et al.*, 2002; Otero *et al.*, 2014) between bromoalkylquinoline **2-7** and 3,4-dimethoxy-4'-
45
46 hydroxychalcone. Reaction yields ranged between 44% and 65%. Chalcone was prepared using a
47
48 previously described method (Peyman *et al.*, 2004) (Scheme 1). Compounds **2-7** were obtained using
49
50 the same method from 8-hydroxyquinoline and dibromoalkanes with different numbers of carbon
51
52 atoms with yields between 51% and 80%.
53
54
55
56
57
58
59
60
61
62
63
64
65

Quinoline-chromone hybrids were obtained following the same synthetic strategy (Scheme 1). Initially, 7-hydroxychromone was treated with potassium hydroxide and 1, ω -dibromoalkanes ($\omega = 3, 4, 5, 7, 9$ and 12) to obtain the respective bromoalkyl derivatives **15-19** in yields similar to previous reports (Otero *et al.*, 2014; Li *et al.*, 2013) but in significantly shorter times. These compounds were coupled with quinoline to produce compounds **20-24** in 34-70% yields. Remarkably, low yields were obtained when bromoalkylquinoline analogues were used as tactical variants.



Scheme 1 Synthetic pathway to quinoline-chalcone and quinoline-chromone hybrids.

1
2
3 Leishmanicidal, trypanocidal and cytotoxic activities
4

5
6 The effect of quinoline-chalcone and quinolone-chromone hybrids on cell growth and viability was
7
8 assessed in human macrophages (U-937 cells) (Pulido *et al.*, 2012) which are the host cells for *L. (V)*
9
10 *panamensis* and *T. cruzi* parasites. On the other hand, the antiparasite activity of these compounds was
11
12 tested on intracellular amastigotes of *L. (V.) panamensis* (Taylor *et al.*, 2011) and *T. cruzi* (Buckner *et*
13
14 *al.*, 1996; Insuasty *et al.*, 2015) according to the ability of these compounds to reduce the amount of
15
16 parasite inside infected macrophages. Results are summarized in the [Table 1](#).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 *In vitro* cytotoxicity and antiprotozoal activity of quinoline-chalcone and quinoline-chromone hybrids

Compound	Cytotoxicity (U-937 cells)	Leishmanicidal activity		Trypanocidal activity	
	LC ₅₀ ^a	EC ₅₀ ^b	SI ^c	EC ₅₀	SI
8	39.6 ± 1.5, 84.34	11.79 ± 0.34, 25.11	3.36	35.08 ± 4.17, 74.71	1.13
9	16.1 ± 0.9, 33.29	6.24 ± 0.14, 12.90	2.58	17.62 ± 1.56, 36.44	0.91
10	23.5 ± 2.4, 47.23	12.37 ± 0.93, 24.86	1.90	15.79 ± 1.47, 31.73	1.49
11	16.7 ± 1.4, 30.94	8.53 ± 0.69, 15.81	1.96	37.61 ± 4.07, 69.69	0.44
12	>40, 72.24	16.41 ± 2.47, 29.64	>2.43	15.12 ± 1.75, 27.30	>2.64
13	69.2 ± 9.3, 116.15	22.0 ± 4.47, 36.93	3.15	54.95 ± 5.47, 92.23	1.26
20	5.4 ± 0.9, 14.94	6.11 ± 0.26, 16.91	0.89	4.09 ± 0.24, 11.32	1.33
21	113.5 ± 16.3, 302.33	48.29 ± 8.18, 128.63	2.35	40.70 ± 7.40, 108.41	2.79
22	121.4 ± 9.3, 290.77	21.54 ± 6.47, 51.59	5.64	28.43 ± 2.77, 68.09	4.27
23	4.1 ± 0.2, 9.50	7.35 ± 1.15, 17.03	0.56	>2, >4.63	<2
24	71.9 ± 8.9, 151.81	16.18 ± 1.45, 34.16	4.89	>20, >42.23	<3.95
3,4-dimethoxy-4'-hydroxychalcone	4.1 ± 0.3, 14.42	2.36 ± 0.37, 8.30	1.75	>2, >7.03	<2
8-hydroxyquinoline (1)	0.2 ± 0.01, 1.38	0.36 ± 0.02, 2.48	0.62	0.34 ± 0.07, 2.34	0.66
7-hydroxychromone (14)	132.5 ± 25.7, 814.11	116.49 ± 13.27, 718.45	1.14	18.23 ± 2.62, 112.43	7.27
Meglumine antimoniate	416.4 ± 66.6	9.4 ± 2.1	44.3	NA ^d	NA ^d
Amphotericin B	42.1 ± 2.0, 45.6	0.05 ± 0.01, 0.054	842	NA ^d	NA ^d
Benznidazole	179.0 ± 4.2, 687.8	NA ^d	NA ^d	10.5 ± 1.8, 40.3	17.0

Data represent mean value +/- standard deviation; ^a LC₅₀: Lethal Concentration 50 in µg/mL, µM; ^b EC₅₀: Effective Concentration 50 in µg/mL, µM; ^c SI: Selectivity Index = LC₅₀/ EC₅₀; ^dNA: Not applicable. Active compounds: EC₅₀ < 25 µg/mL

1
2
3 All compounds, amphotericin B and benznidazole with exception of **21**, **22** and chromone, were
4
5 highly cytotoxic to U-937 cells showing $LC_{50} < 100.0 \mu\text{g/mL}$ (Table 1). Compounds **21**, **22** and
6
7 chromone, showed moderate cytotoxicity evidenced by LC_{50} values higher than $100.0 \mu\text{g/mL}$. In turn,
8
9 meglumine antimoniate showed no cytotoxicity ($LC_{50} > 200.0 \mu\text{g/mL}$).
10

11
12
13 The anti-leishmanial and anti-trypanocidal activities were measured by determining the effective
14
15 concentration 50 (EC_{50}) that corresponds to the concentration of drug that gives the half-maximal
16
17 reduction of the amount of intracellular parasites (Table 1). Dose-response relationship showed that
18
19 compounds **8-12**, **20**, **23**, **24**, chalcone and quinoline were highly active against intracellular
20
21 amastigotes of *L. (V) panamensis* with $EC_{50} < 20 \mu\text{g/mL}$. The most actives hybrids compounds were
22
23 **20**, **9**, **23** and **11** with an EC_{50} of $6.11 \pm 0.26 \mu\text{g/mL}$ ($16.91 \mu\text{M}$), $6.24 \pm 0.14 \mu\text{g/mL}$ ($12.90 \mu\text{M}$), 7.35
24
25 $\pm 1.15 \mu\text{g/mL}$ ($17.03 \mu\text{M}$) and $8.53 \pm 0.69 \mu\text{g/mL}$ ($15.81 \mu\text{M}$) respectively, followed by **8**, **10**, **12** and
26
27 **24** with an EC_{50} of $11.79 \pm 0.34 \mu\text{g/mL}$ ($25.11 \mu\text{M}$), $12.37 \pm 0.93 \mu\text{g/mL}$ ($24.86 \mu\text{M}$), 16.41 ± 2.47
28
29 $\mu\text{g/mL}$, $29.64 \mu\text{M}$ $16.18 \pm 1.45 \mu\text{g/mL}$ ($34.16 \mu\text{M}$). As expected, the leishmanicidal drugs amphotericin
30
31 B and meglumine antimoniate showed activity with low EC_{50} values.
32
33
34
35
36
37
38
39

40 In turn, compounds **20**, **23**, chalcone and quinoline were highly active against intracellular
41
42 amastigotes of *T. cruzi* with EC_{50} of $4.09 \pm 0.24 \mu\text{g/mL}$ ($11.32 \mu\text{M}$), $>2 \mu\text{g/mL}$ ($>4.63 \mu\text{M}$), $>2 \mu\text{g/mL}$
43
44 ($>7.03 \mu\text{M}$) $0.34 \pm 0.07 \mu\text{g/mL}$ ($2.34 \mu\text{M}$) respectively, followed by compounds **9**, **10**, **12** and
45
46 chromone with an EC_{50} of $17.62 \pm 1.56 \mu\text{g/mL}$ ($36.44 \mu\text{M}$), $15.79 \pm 1.47 \mu\text{g/mL}$ ($31.73 \mu\text{M}$), $15.12 \pm$
47
48 $1.75 \mu\text{g/mL}$ ($27.30 \mu\text{M}$) and $18.23 \pm 2.62 \mu\text{g/mL}$ ($112.43 \mu\text{M}$) respectively. In this case, benznidazole
49
50 showed activity with an EC_{50} of $10.5 \pm 1.8 \mu\text{g/mL}$ ($40.3 \mu\text{M}$).
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 The anti-leishmanial activity of compounds **8-13**, **21**, **22**, **24**, chalcone and chromone and anti-
4
5 trypanocidal activity of compounds **8**, **10**, **12**, **13**, **20-24** and chromone were higher than their
6
7 cytotoxicity. Thus, the SI (Selectivity Index) values calculated for these compounds were >1 (Table 1).
8
9

10
11
12 As demonstrated elsewhere, amphotericin B and meglumine antimoniate have very high SI values.
13
14 Although all hybrid compounds showed better activity than meglumine antimoniate and the anti-
15
16 trypanocidal activity of compounds **20** and **23** were higher than benznidazole, the SI of these
17
18 compounds is affected by their high cytotoxicity. These results suggest that biological activity of the
19
20 quinoline derivatives reported here, with exception of **20** and **23**, is selective, being more active against
21
22 *L. (V) panamensis* than U-937 cells. On the other hand, compounds **8**, **10**, **12**, **13**, **20-24** were more
23
24 actives against *T. cruzi* parasites than U-937 cells.
25
26
27
28
29
30
31

32
33 There is not a clear relationship between the antiprotozoal activity and the length of the alkyl linker,
34
35 because as the chain length increase (increased lipophilicity), the activity does not show a definite
36
37 tendency. All quinoline-chalcone hybrids were less cytotoxic and less active than the parent subunits.
38
39 However, a synergistic effect of the parent subunits was observed in the quinoline-chromone hybrids in
40
41 comparison with the unlinked cases, due to decreased cytotoxicity based on quinoline and increased
42
43 activity based chromone. One possible mechanism of action for these compounds may be formulated in
44
45 terms of conjugated addition of nucleophilic amino acid residues present in target enzymes of
46
47 *Leishmania* e.g. such cysteine proteases (Mottram *et al.*, 2004) in a Michael addition (Cardona *et al.*,
48
49 2014).
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 **Conclusion**
4
5

6 The synthesis, anti-leishmanial and anti-trypanocidal screening of eleven quinoline derivatives are
7 reported. Several of the synthetic compounds have potential as templates for drugs development. Eight
8 of them were active against *L. (V) panamensis* (**8-12, 20, 23** and **24**) and five of them against *T. cruzi*
9 (**9, 10, 12, 20** and **23**) with EC₅₀ values lower than 18 µg/mL, **20** being the most active compound for
10 both *L. (V) panamensis* and *T. cruzi*. All hybrid compounds showed better activity than meglumine
11 antimoniate and compounds **20** and **23** showed higher actives than benznidazole. Studies on an animal
12 model of leishmaniasis are needed to confirm the results observed *in vitro*. These compounds were
13 toxic for mammalian U-937 cells, however they may still have potential to be considered as
14 candidates to antileishmanial drug development. More studies on toxicity using other cell lines
15 are needed to discriminate whether the toxicity shown by these compounds is specific against
16 tumor or non-tumor cells. Moreover, Michael acceptor moieties may modify the parasitic activity
17 and cytotoxicity. The mechanism of action of these promising compounds also needs to be addressed.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33
34
35
36
37
38 **Acknowledgments**
39

40 The authors thank Universidad de Antioquia (grant CODI IN656CE and CIDEPRO) for financial
41 support.
42
43
44
45
46
47

48 **Conflict of interest**
49

50 The authors declare that they have no conflict of interest.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J, den Boer M (2012) Leishmaniasis Worldwide and Global Estimates of Its Incidence. PLOS One 7: e35671
- Baloch N, Alkahraman Y, Zaidi M, Madkour H (2012) Evaluation of 6, 8-Dichloro-2-Methyl-4H-Chromen-4-One Derivatives as Antileishmanial Agents. Global J Sci Front Res Chem 12:1-7
- Bhutta ZA, Sommerfeld J, Lassi ZS, Salam RA, Das JK (2014) Tackling the existing burden of infection diseases in the developing world: existing gaps and the way forward. Infect Dis Poverty 3: 1-6
- Boeck P, Bandeira Falcão CA, Leal PC, Yones RA, Filho VC, Torres-Santos EC, Rossi-Bergmann B (2006) Synthesis of chalcone analogues with increased antileishmanial activity. Bioorg Med Chem 14:1538-1545
- Buckner FS, Verlinde CL, La Flamme AC, Van Voorhis WC (1996) Efficient technique for screening drugs for activity against *Trypanosoma cruzi* using parasites expressing beta-galactosidase. Antimicrob Agents Chemother 40: 2592-2597
- Cardona W, Arango V, Domínguez J, Robledo S, Muñoz D, Figadère B, Velez ID, Sáez, J (2013) Synthesis and leishmanicidal activity of new bis-alkylquinolines. J Chil Chem Soc 58:1709-1712
- Cardona W, Guerra D, Restrepo A (2014) Reactivity of δ -substituted α,β -unsaturated cyclic lactones with antileishmanial activity. Mol Simul 40:477-484
- Chatelain E, Ioset JR (2011) Drug discovery and development for neglected diseases: the DNDi model. Drug Des Devel Ther 16:175-181
- Chen M, Zhai L, Christensen SB, Theander TG, Kharazmi A (2001) Inhibition of fumarate reductase in *Leishmania major* and *L. donovani* by chalcones. Antimicrobiol. Agents Chemother. 45:2023-2029
- Coa JC, Castrillón W, Cardona W, Carda M, Ospina V, Muñoz JA, Vélez ID, Robledo SM (2015)

1
2
3 Synthesis, leishmanicidal, trypanocidal and cytotoxic activity of quinoline-hydrazone hybrids. *Eur J*
4
5 *Med Chem* 101:746-753
6
7
8 Den Boer M, Argaw D, Jannin J, Alvar J (2011) Leishmaniasis impact and treatment access. *Clin*
9
10 *Microbiol Infect* 17:1471-1477
11
12
13 Dietze R, Carvalho SF, Valli LC, Berman J, Brewer T, Milhous W, Sanchez J, Schuster B, Grogl M
14
15 (2001) Phase 2 trial of WR6026, an orally administered 8-aminoquinoline, in the treatment of
16
17 visceral leishmaniasis caused by *Leishmania chagasi*. *Am J Trop Med Hyg* 65:685-689
18
19
20
21
22 Ellis GP, Barker G (1972) Chromone-2- and -3-carboxylic acids and their derivatives. *Progr Med*
23
24 *Chem* 9:65-116
25
26
27 Finney JD (1978) Probit Analysis: Statistical Treatment of the Sigmoid Response Curve, 3rd
28
29 ed.; Cambridge University Press: Cambridge, UK, p. 550.
30
31
32 Franck X, Fournet A, Prina E, Mahieux R, Hocquemiller R, Figadère B (2004) Biological evaluation of
33
34 substituted quinolones. *Bioorg Med Chem Lett* 14:3635-3638
35
36
37 Hadjeri M, Barbier M, Ronot X, Mariotte AM, Boumendjel A, Boutonnat J (2003) Modulation of P-
38
39 glycoprotein-mediated multidrug resistance by flavonoid derivatives and analogues. *J Med Chem*
40
41 46:2125-2131
42
43
44 Horton DA, Boume GT, Smythe ML (2003). The combinatorial synthesis of bicyclic privileged
45
46 structures or privileged substructures. *Chem Rev* 103:893-930
47
48
49 Houghton PJ (2000) Chemistry and biological activity of natural and semi-synthetic 450 chromone
50
51 alkaloids. *Stud Nat Prod Chem* 21:123-155
52
53
54 Insuasty B, Ramirez J, Becerra D, Echeverry C, Quiroga J, Abonia R, Robledo SM, Velez ID, Upegui
55
56 Y, Muñoz JA, Ospina V, Noguera M, Cobo J (2015). An efficient synthesis of a new caffeine-based
57
58 chalcones, pyrazolines and pyrazolo[3-4-b][1-4]diazepines as potential antimalarial,
59
60
61
62
63
64
65

- 1
2
3 antitrypanosomal and antileishmanial agents. *Eur J Chem Med* 93:401-413
4
5 Kayser O, Kiderlen AF (2001) In vitro leishmanicidal activity of naturally occurring chalcones.
6
7 *Phytother Res* 15:148-152
8
9
10 Keenan M, Chaplin JH (2015) A new era for chagas disease drug discovery?, *Prog Med Chem* 54:
11
12 185-230
13
14
15 Li SY, Wang XB, Xie SS, Jian N, Wang KD, Yao HQ, Sun HB, Kong LY (2013)
16
17 Multifunctional tacrine-flavonoid hybrids with cholinergic, β -amyloid-reducing, and metal chelating
18
19 properties for the treatment of Alzheimer's disease. *Eur J Med Chem* 69:632-646
20
21
22 Liu M, Wilairat P, Croft SL, Tand AL, Go ML (2003) Structure-activity relationships of
23
24 antileishmanial and antimalarial chalcones. *Bioorg Med Chem* 11: 2729-2738
25
26
27 Mallick S, Dutta A, Ghosh J, Maiti S, Mandal AK, Banerjee R, Bandyopadhyay C, Pal C (2011)
28
29 Protective therapy with novel chromone derivative against *Leishmania donovani* infection induces
30
31 Th1 response in vivo. *Chemotherapy* 57:388-393
32
33
34 Mohammed AF, Fournet A, Prina E, Mouscadet JF, Franck X, Hocquemiller R, Figadère B, Fakhfakh
35
36 MA (2003) Synthesis and biological evaluation of substituted quinolines: potential treatment of
37
38 protozoal and retroviral co-infections. *Bioorg Med Chem* 11:5013-5023
39
40
41 Mottram JC, Coombs GH, Alexander J (2004) Cysteine peptidases as virulence factors of *Leishmania*.
42
43 *Curr Opin Microbiol* 7:375-381
44
45
46 Nakayama H, Loiseau PM, Bories C, De Ortiz ST, Schinini A, Serna E, Rojas de Arias A, Fakhfakh
47
48 MA, Franck X, Figadère B, Hocquemiller R, Fournet A (2005) Efficacy of orally administered 2-
49
50 substituted quinolines in experimental murine cutaneous and visceral leishmaniases. *Antimicrob.*
51
52 *Agents Chemother* 49 :4950-4956
53
54
55
56 Nouvellet P, Cucunubá ZM, Gourbière S (2015) Ecology, evolution and control of Chagas disease: a
57
58 century of neglected modelling and promising future. *Adv Parasitol* 87:135-191
59
60
61
62
63
64
65

- 1
2
3 Otero E, Vergara S, Robledo SM, Cardona W, Carda M, Vélez ID, Rojas C, Otálvaro F (2014)
4
5 Synthesis, Leishmanicidal and Cytotoxic Activity of Triclosan-Chalcone, Triclosan-Chromone and
6
7 Triclosan-Coumarin Hybrids. *Molecules* 19:13251-13266
8
9
- 10 Palit P, Paira P, Hazra A, Banerjee S, Das Gupta A, Dastidar S, Mondal N (2009) Phase transfer
11
12 catalyzed synthesis of bis-quinolines: antileishmanial activity in experimental visceral leishmaniasis
13
14 and in vitro antibacterial evaluation. *Eur J Med Chem* 44:845-853
15
16
- 17 Peng Y, Song G (2002) Combined microwave and ultrasound assisted Williamson ether synthesis in
18
19 the absence of phase-transfer catalysts. *Green Chem* 4:349-351.
20
21
- 22 Peyman S, Mino D, Mohammad AZ, Mohammad AB (2004) Silica Sulfuric Acid as an Efficient and
23
24 Reusable Reagent for Crossed-Aldol Condensation of Ketones with Aromatic Aldehydes under
25
26 Solvent-Free Conditions. *J Braz Chem Soc* 15:773-776
27
28
- 29 Pulido SA, Muñoz DL, Restrepo AM, Mesa CV, Alzate JF, Vélez ID, Robledo SM (2012)
30
31 Improvement of the green fluorescent protein reporter system in *Leishmania* spp. for the *in vitro* and
32
33 *in vivo* screening of antileishmanial drugs. *Acta Trop* 122:36-45
34
35
36
- 37 Suresh K, Sandhya B, Himanshu G (2009) Biological activities of quinoline derivatives. *Mini-Rev*
38
39
40 *Med Chem* 9: 1648-1654
41
42
- 43 Taylor VM, Cedeño DL, Muñoz DL, Jones MA, Lash TD, Young AM, Constantino MH, Esposito N,
44
45 Vélez ID, Robledo SM (2011) *In vitro* and *in vivo* studies of the utility of dimethyl and diethyl
46
47 carbaporphyrin ketals in treatment of cutaneous leishmaniasis. *Antimicrob Agents Chemother*
48
49
50
51
52 55:4755-4764
- 53 Tempone A, Melo A, Da Silva P, Brandt C, Martinez F, Borborema A (2005) Synthesis and
54
55 antileishmanial activities of novel 3-substituted quinolones. *Agents chemother* 49:1076-1080
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Vieira NC, Herrenknecht C, Vacus J, Fournet A, Bories C, Figadère B, Espindola LS, Loiseau PM
(2008) Selection of the most promising 2-substituted quinoline as antileishmanial candidate for
clinical trials. *Biomed Pharmacother* 62:684-689

World Health Organization. Why are some tropical diseases called ‘neglected’?
<http://www.who.int/features/qa/58/en/>, 2013 [accessed 04.12.13].