

Portland State University

PDXScholar

Mathematics and Statistics Faculty
Publications and Presentations

Fariborz Maseeh Department of Mathematics
and Statistics

2014

Regularity of Mediatrices in Surfaces


Pilar Herreros

Mario Ponce

J. J. P. Veerman

Portland State University, veerman@pdx.edu

Follow this and additional works at: https://pdxscholar.library.pdx.edu/mth_fac

 Part of the [Applied Mathematics Commons](#), and the [Discrete Mathematics and Combinatorics Commons](#)

Let us know how access to this document benefits you.

Citation Details

Herreros, Pilar; Ponce, Mario; and Veerman, J. J. P., "Regularity of Mediatrices in Surfaces" (2014).
Mathematics and Statistics Faculty Publications and Presentations. 123.
https://pdxscholar.library.pdx.edu/mth_fac/123

This Pre-Print is brought to you for free and open access. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications and Presentations by an authorized administrator of PDXScholar. For more information, please contact pdxscholar@pdx.edu.

REGULARITY OF MEDIATRICES IN SURFACES

PILAR HERREROS, MARIO PONCE AND J.J.P. VEERMAN

ABSTRACT. For distinct points p and q in a two-dimensional Riemannian manifold, one defines their mediatrix L_{pq} as the set of equidistant points to p and q . It is known that mediatrices have a cell decomposition consisting of a finite number of branch points connected by Lipschitz curves. This paper establishes additional geometric regularity properties of mediatrices. We show that mediatrices have the radial linearizability property, which implies that at each point they have a geometrically defined derivative in the branching directions. Also, we study the particular case of mediatrices on spheres, by showing that they are Lipschitz simple closed curves exhibiting at most countably many singularities, with finite total angular deficiency.

1. INTRODUCTION

Let M be a compact, connected Riemannian manifold. For any distinct points $p, q \in M$ the *mediatrix* L_{pq} is defined as the set of points with equal distance to p and q , that is

$$L_{pq} = \{x \in M \mid d(x, p) = d(x, q)\},$$

where $d(\cdot, \cdot)$ is the Riemannian distance in M . These geometrical objects can be found in the very beginning of the foundations of geometry. As an illustration, Leibniz proposed to define *planes* as mediatrices in the space. Mediatrices appear in the literature with multiple names, and have been developed by many communities, often separately. The first non-trivial approach to mediatrices appears in the book [5] by Busemann, under the name of *bisectors*. There, the last chapter is mainly devoted to characterize metric spaces with flat bisectors. The remarkable result by Busemann is that metric spaces having flat bisectors are (up to a cover) isometric to a finite dimensional euclidian, hyperbolic, or spherical space. In a slightly more general framework, where one replaces points p, q by disjoint compact sets $P, Q \subset M$, the locus of points with equal distance to P and Q have been studied using the names *conflict sets* (see for instance [17], [3]), *midsets* (see [10]), *equidistant sets* (see [16], [12]), and *skeletal or medial structures* (see [7]).

The study of mediatrices came about to answer some questions that came up in the study of *Focal Decomposition* pioneered by [11]. This point of view was taken in [15]. In spite of the work done in [11], there still are many open problems related to even the simplest cases such as that of the “mathematical pendulum”. The simple observation of that mediatrices are level sets of the difference of the distance functions to the corresponding points, allows to consider the relationship between the cut locus and mediatrices. Namely, the distance function to a point is differentiable outside the cut locus and hence, the Implicit Function Theorem yields that a mediatrix is differentiable outside the union of the cut locus of the underlying points. It is well-known (Theorems by Singer-Gluck and Itoh, see for instance [8], [6]) that the cut locus even for a C^∞ surface of revolution can be a pretty awful set. In contrast the mediatrix on any C^r , $r \geq 3$ surface is locally really surprisingly well-behaved (as we show below).

For $p \neq q \in M$, the mediatrix is the boundary between the two disjoint connected components $\{d(x, p) < d(x, q)\}$ and $\{d(x, q) < d(x, p)\}$ of $M \setminus L_{pq}$. Notice that by removing any single point of

the mediatrix these sets stop being separated. Hence one says that mediatrices are *Minimally Separating Sets*. Separating Sets and especially Minimally Separating Sets are interesting since they occur in a variety of places throughout mathematics. In the first place they occur as boundaries between the regions where different types of Focalization occurs as explained in [11] and [15]. The classical ‘Lakes of Wada’ construction is another example (see [9]). Basin boundaries may in some cases be another instance of where these sets naturally occur. The latter are related to Julia sets, Birkhoff attractor and so on. Recently other topological properties of this kind of sets have gained some attention, namely the fact that some are indecomposable continua (see for example [13]).

In [15], Bernhard and Veerman show that mediatrices in surfaces have a cell decomposition consisting of a finite number of branch points connected by Lipschitz curves. This paper establishes additional geometric regularity properties of mediatrices in surfaces (see definitions below).

Main Theorem (short version). *Mediatrices in surfaces have the radial linearizability property.*

This implies that at each point mediatrices have a geometrically defined derivative in the branching directions. We devote Section 4 to the study of the particular case of mediatrices on spheres, by showing that they are Lipschitz simple closed curves exhibiting at most countably many singularities, with finite total angular deficiency.

2. PRELIMINARIES

In what follows, M is a two-dimensional compact Riemannian manifold of class C^r , $r \geq 3$.

Minimizing Geodesics. Let $x \in M$ and θ in the unit tangent sphere $S_x M$. We write $\gamma_{x,\theta} : \mathbb{R} \rightarrow M$ the unit speed geodesic satisfying $\gamma_{x,\theta}(0) = x$ and $\gamma'_{x,\theta}(0) = \theta$. For $t^* > 0$ we say that $\gamma_{x,\theta}$ is a minimizing geodesic between x and $\gamma_{x,\theta}(t^*)$ if the length of $\gamma_{x,\theta}$ equals the infimum of lengths among absolutely continuous curves joining x to $\gamma_{x,\theta}(t^*)$, that is, $length(\gamma_{x,\theta}|_{[0,t^*]}) = d(x, \gamma_{x,\theta}(t^*))$. The triangle inequality yields that $\gamma_{x,\theta}$ is minimizing for every $0 < t < t^*$.

Polar Coordinates. Recall that the exponential map exp_x at $x \in M$ is defined as

$$exp_x(v) = \gamma_{x,\hat{v}}(|v|), \text{ where } v \in T_x M, \hat{v} = \frac{v}{|v|}, \text{ and } exp_x(0) = x.$$

For small $\rho > 0$ (less than the *injectivity radius* $injr(M) > 0$), one has that $exp_x|_{\{|v| < \rho\}}$ is a diffeomorphism. By fixing a polar coordinates system in $T_x M$, one can define (via exp_x) a local polar coordinates system centered at x on $B_{x,\rho} = \{z \in M \mid d(x,z) < \rho\}$. In that way, a point $z \in B_{x,\rho}$, $z \neq x$, is identified with a pair $(\theta, d) \in \mathbb{S}^1 \times (0, \rho)$, such that $z = \gamma_{x,\theta}(d)$, $d = d(x, z)$ (recall that geodesics are locally minimizing curves, see for instance [6]).

Minimizing Directions. For $p \in M$, $\Theta_{x,p} \subset S_x M$ is the set of directions θ such that $\gamma_{x,\theta}$ is a minimizing curve from x to p . We endow $S_x M$ with the usual arc-distance in the circle. Since M is complete one has

Lemma 1 (Hopf-Rinow Theorem, see [6]). *For $x \neq p$ the set $\Theta_{x,p}$ is non-empty and compact.* \square

Lemma 2 (see [2]). *Let $p \neq q \in M$ and $x \in L_{pq}$. Then $\Theta_{x,p} \cap \Theta_{x,q} = \emptyset$.*

Proof. Let $\theta \in \Theta_{x,p} \cap \Theta_{x,q}$ and $d = d(x, p) = d(x, q)$. Then $\gamma_{x,\theta}(d) = q = p$, which is impossible. \square

Lemma 3. *Let $x \neq p$, and two convergent sequences $x_n \rightarrow x$, $\theta_n \rightarrow \theta$, with $\theta_n \in \Theta_{x_n, p}$ for every n (we identify $S_{x_n}M$ with S_xM via a local chart). Then one has $\theta \in \Theta_{x, p}$.*

Proof. Let $d_n = d(x_n, p)$. Continuity of the distance function implies $d_n \rightarrow d(x, p)$. We have $\gamma_{x_n, \theta_n}(d_n) = p$ for every n . Taking the limit we obtain $\gamma_{x, \theta}(d(x, p)) = p$, and hence $\theta \in \Theta_{x, p}$. \square

Corollary 4. *If $x \in L_{pq}$ then there exist $\rho_x > 0$ and $\beta_x \in (0, \pi]$ such that*

$$d_{S_x M}(\Theta_{z, p}, \Theta_{z, q}) \geq \beta_x$$

for all $z \in B_{x, \rho_x}$. \square

Proof. Given x we know from Lemma 2 that $\Theta_{x, p} \cap \Theta_{x, q} = \emptyset$. Assume by contradiction that there exist sequences $z_n \rightarrow x$, $\theta_{p, n} \in \Theta_{z_n, p}$, $\theta_{q, n} \in \Theta_{z_n, q}$ with $|\theta_{p, n} - \theta_{q, n}| \rightarrow 0$. Choosing subsequences we can assume that there exists $\theta \in S_x M$ such that $\theta_{p, n} \rightarrow \theta$, $\theta_{q, n} \rightarrow \theta$. Lemma 3 implies that $\theta \in \Theta_{x, p} \cap \Theta_{x, q}$, which is impossible. \square

For $p \neq q \in M$ and $x \in L_{pq}$, fix a positive orientation in $S_x M$. A closed connected set P in $S_x M$ is called a *pre-wedge at x* if its end-points are minimizing directions to p and q respectively, and there are no other minimizing directions to p or q in its interior. Such a set is clearly an interval in the circle, and with a slight abuse of notation we denote it as: $P = [\theta_p, \theta_q] \subset S_x M$ (or $[\theta_q, \theta_p]$). Thus P satisfies:

- $\theta_p \in \Theta_{x, p}$, $\theta_q \in \Theta_{x, q}$.
- $[\theta_p, \theta_q] \cap (\Theta_{x, p} \cup \Theta_{x, q}) = \{\theta_p, \theta_q\}$.

The *midpoint* $\hat{\theta}$ of a wedge $P = [\theta_p, \theta_q]$ is the unique point *contained in the pre-wedge* that equidistant from its endpoints.

Lemma 5 (see [2]). *There are finitely many pre-wedges at $x \in L_{pq}$.*

Proof. Assume $([\theta_p^n, \theta_q^n])_{n \in \mathbb{N}}$ is an infinite sequence of different pre-wedges at x . We must have $\lim_n |\theta_p^n - \theta_q^n| = 0$, since pre-wedges have disjoint interiors. We conclude that the sequences of end-points $(\theta_p^n)_{n \in \mathbb{N}}$, $(\theta_q^n)_{n \in \mathbb{N}}$ have a common accumulation point $\hat{\theta}$. Compactness of $\Theta_{x, p}$ and $\Theta_{x, q}$ yields that $\hat{\theta}$ belongs to $\Theta_{x, p} \cap \Theta_{x, q}$ which is impossible. \square

For $0 < \rho < \text{in}jrad(M)$ and a pre-wedge $[\theta_p, \theta_q] \subset S_x M$, we define the corresponding *wedge at x of radius ρ* as

$$W_{x, \rho}[\theta_p, \theta_q] = \{\text{exp}_x(r, \theta) \mid 0 \leq r < \rho, \theta \in [\theta_p, \theta_q]\} \subset M.$$

In [2] the authors show how the mediatrix L_{pq} is located in M with respect to the wedges at $x \in L_{pq}$ in the following sense:

Theorem 6 (Bernhard and Veerman [2]). *Let M be a 2-dimensional compact Riemannian manifold of class C^r , $r \geq 3$. For every pair of distinct points $p, q \in M$ the mediatrix $L_{pq} = \{x \in M \mid d(x, p) = d(x, q)\}$ verifies*

- (1) *Let $x \in L_{pq}$. If the cardinalities $\#\Theta_{x, p}$ and $\#\Theta_{x, q}$ are both 1, then for small ρ the intersection $L_{pq} \cap B_{x, \rho}$ is a continuous simple curve passing through x and differentiable at x .*
- (2) *Let $x \in L_{pq}$ and $0 < \rho < \text{in}jrad(M)$. The intersection $L_{pq} \cap B_{x, \rho}$ is contained in the finite union of the wedges of radius ρ at x .*
- (3) *Let $W_{x, \rho}[\theta_p, \theta_q]$ be a wedge at $x \in L_{pq}$. The intersection $L_{pq} \cap W_{x, \rho}[\theta_p, \theta_q]$ is a Lipschitz simple curve connecting x to a boundary point of $W_{x, \rho}[\theta_p, \theta_q]$. This curve is called a *spoke* at x .*
- (4) *There are finitely many points $x \in L_{pq}$ so that there exists more than two pre-wedges at x .*

(5) *The mediatrice L_{pq} is homeomorphic to a finite and closed simplicial 1-complex.*

Notice that 1. above is a direct consequence of the fact that the distance function to a fixed point $p \in M$ is differentiable outside the cut locus of p and a direct application of the Implicit Function Theorem. The Theorem above says that $L_{pq} \cap B_{x,\rho}$ is the finite union of two or more Lipschitz spokes emanating from $x \in L_{pq}$. The goal of this article is to study the behavior of a single spoke as it approaches $x \in L_{pq}$. We show that spokes are *radially linearizable* in the following sense:

Definition. Let $\phi : [0, 1] \rightarrow M$ be a continuous simple curve. We say that η is *radially linearizable* at $x = \phi(0) \in M$ if there exists a direction $\hat{\theta} \in S_x M$ such that

$$\lim_{t \rightarrow 0^+} \theta(t) = \hat{\theta},$$

where $\phi(t) = (\theta(t), d(t))$ in polar coordinates centered at x .

Main Theorem [Radial Linearizability of Mediatrices]. *Let M be a 2-dimensional compact Riemannian manifold of class C^r , $r \geq 3$. Let $p \neq q \in M$, $x \in L_{pq}$, and $W_{x,\rho}[\theta_p, \theta_q]$ a wedge at x . The spoke $L_{pq} \cap W_{x,\rho}[\theta_p, \theta_q]$ is radially linearizable at x in the direction of the midpoint $\hat{\theta}$ of $[\theta_p, \theta_q]$.*

Notice that this Theorem asserts that $L_{pq} \cap W_{x,\rho}[\theta_p, \theta_q]$ is tangent to the bisector of the angle $\angle \theta_p \theta_q$, and hence, L_{pq} is tangent to the directions of the equidistant set to $\Theta_{x,p}$ and $\Theta_{x,q}$ in $S_x M$. A similar result was conjectured in [2] (compare also with [3]).

Corollary. *Mediatrices in surfaces have no cusp like points.*

3. PROOF OF THE MAIN THEOREM

The distance function has one-side directional differential. For $p \in M$, the triangle inequality implies that the function $x \mapsto d(x, p)$ is 1-Lipschitz. Even though $d(x, p)$ is not necessarily differentiable in M (in fact, it is not differentiable at the cut locus of p , see for instance [6]), for every $x \in M$ this function has a one-sided derivative in every direction. More precisely, let $\theta \in S_x M$. The limit

$$\lim_{t \rightarrow 0^+} \frac{d(\gamma_{x,\theta}(t), p) - d(\gamma_{x,\theta}(0), p)}{t}$$

does exist. This fact, and the explicit formula for the limit, is apparently a well known fact that can be deduced from the first variation formula. Nevertheless, just recently, and due to the absence of any unambiguous literature in the subject, I. Adelstein [1] has produced an explicit proof (see also Section 4.5 in [4]).

Proposition 7. *Using the above notation, one has*

$$\lim_{t \rightarrow 0^+} \frac{d(\gamma_{x,\theta}(t), p) - d(\gamma_{x,\theta}(0), p)}{t} = -\cos(d_{S_x M}(\theta, \Theta_{x,p}))$$

where $d_{S_x M}(\cdot, \cdot)$ is the Riemannian distance on the unit circle between θ and the compact set of minimizing directions from x to p . \square

Lemma 8. *Let $g : [0, 1] \rightarrow \mathbb{R}$ be a continuous function and $c \in \mathbb{R}$. Assume that the one-side derivative exists and verifies*

$$g'_+(t) \equiv \lim_{s \rightarrow 0^+} \frac{g(t+s) - g(t)}{s} > c$$

for every $t \in [0, 1)$. Then $g(t) > g(0) + ct$ for every $t \in (0, 1)$.

Proof. By considering the function $g(t) - ct$ we can reduce us to the case $c = 0$. For every $t \in [0, 1)$ there exists $\delta_t > 0$ such that $g(t + s) - g(t) > 0$ for $0 < s < \delta_t$. Hence g can not have local maximal points. As g is continuous, the only possibility is that g is strictly increasing. \square

In what follows we use the notation of the statement of the Main Theorem, that is, we fix a pre-wedge $[\theta_p, \theta_q]$ at $x \in L_{pq}$, write $\hat{\theta}$ the midpoint of $[\theta_p, \theta_q]$, and $\rho_x > 0$ given by Corollary 4.

Lemma 9. *If $L_{pq} \cap W_{x,\rho}[\theta_p, \theta_q] \cap \gamma_{x,\theta}([0, \rho_x])$ contains a sequence of points converging to x then $\theta = \hat{\theta}$.*

Proof. Define $f_{pq}(z) = d(z, q) - d(z, p)$. Hence $L_{pq} = f_{pq}^{-1}(0)$. Proposition 7 allows to deduce that f_{pq} has one-side differential at x in the direction of θ equals to $-\cos|\theta - \tilde{\theta}_p| + \cos|\theta - \tilde{\theta}_q|$ for some $\tilde{\theta}_p \in \Theta_{x,p}$, $\tilde{\theta}_q \in \Theta_{x,q}$. If the restriction of f_{pq} to the ray $\gamma_{x,\theta}$ has roots accumulating at x then this one-side differential must be equal to zero. The only possible case is $\tilde{\theta}_p = \theta_p, \tilde{\theta}_q = \theta_q$ (otherwise minimizing geodesics will cross) and it follows that the only solution is $\theta = \hat{\theta}$. \square

This Lemma says that whenever a spoke crosses infinitely many times a geodesic ray emanating from x , and the crossing points approach x , then the geodesic ray necessarily points in the direction of the bisector of the corresponding pre-wedge. The next result estimates for how much the points in the geodesic ray $\gamma_{x,\hat{\theta}}$ fail to belong to the mediatrice L_{pq} .

Lemma 10. *Given p and q , then for each $x \in L_{pq}$ there exists a positive function η such that*

$$\left| d\left(\gamma_{x,\hat{\theta}}(t), q\right) - d\left(\gamma_{x,\hat{\theta}}(t), p\right) \right| \leq \eta(t)t \text{ for } 0 \leq t < \rho,$$

and $\lim_{t \rightarrow 0} \eta(t) = 0$.

Proof. The function $t \mapsto f_{pq}\left(\gamma_{x,\hat{\theta}}(t)\right)$ together with its first (right directional) derivative vanish at $t = 0$ and the result follows. \square

The next result tells us that close to points that fail to belong to L_{pq} by few, we can find a genuine point in the mediatrice. This result is inspired by Section 4.2 of [12].

Lemma 11. *Let $x \in L_{pq}$ and $\rho_x > 0, \beta_x \in (0, \pi)$ given by Corollary 4. There exists $\varepsilon_x > 0$ such that for all $z \in B_{x,\rho_x/2}$ that verifies $|d(z, p) - d(z, q)| < \varepsilon_x$ there exists $z^* \in L_{pq}$, and*

$$d(z, z^*) < \frac{|d(z, q) - d(z, p)|}{1 - \cos(\beta_x)}.$$

Proof. Let's assume that $d(z, q) > d(z, p)$. Pick $\theta \in \Theta_{z,q}$ (this also gives $\theta \notin \Theta_{z,p}$). By moving continuously along the geodesic ray $z(t) \equiv \gamma_{z,\theta}(t)$ we must encounter a point $z^* \equiv z(t^*)$ in the mediatrice L_{pq} (since $\gamma_{z,\theta}$ reaches q). We estimate the distance t^* between z and z^* . Note that since $d(x, z) < \rho_x/2$ then by construction $d(x, z(t)) < \rho_x$ at least for $t \leq \rho_x/2$.

By Proposition 7 we have


$$\lim_{t \rightarrow 0^+} \frac{d(z(t), p) - d(z(0), p)}{t} = -\cos(d_{S_{z(t)}M}(z'(t), \Theta_{z(t),p}))$$

For every $t \in [0, \rho_x/2]$ one has $z'(t) \in \Theta_{z(t),q}$ and so Corollary 4 implies that there is a $\beta_x \in (0, \pi)$ so that

$$\cos(d_{S_{z(t)}M}(z'(t), \Theta_{z(t),p})) \leq \cos \beta_x$$

Now Lemma 8 implies that for every $t \in (0, \rho_x/2]$

$$d(z(t), p) \geq d(z, p) - t \cos \beta_x.$$



We also have that

$$d(z(t), q) = d(z, q) - t.$$

Putting these two together we have that

$$\begin{aligned} d(z(0), q) - d(z(0), p) &> 0 \\ d(z(t), q) - d(z(t), p) &\leq d(z(0), q) - d(z(0), p) - t(1 - \cos(\beta_x)) \end{aligned}$$

The latter is less than or equal to zero at $t_0 = \frac{d(z, q) - d(z, p)}{1 - \cos(\beta_x)}$. By continuity of the distance function, we see that

$$t^* \leq \frac{d(z, q) - d(z, p)}{1 - \cos(\beta_x)}.$$

If we choose $\varepsilon_x \equiv \frac{1}{2}\rho_x(1 - \cos(\beta_x)) > 0$, so that $t_0 < \rho_x/2$, we stay inside B_{x, ρ_x} and the Lemma is verified. \square


Proof of the Main Theorem. Pick $x \in L_{pq}$ and a wedge $W_{x, \rho}[\theta_p, \theta_q]$. Reasoning by contradiction let's suppose that there exists $\tau > 0$ such that the spoke contains a sequence of points x_n converging to x outside the cone $[\hat{\theta} - \tau, \hat{\theta} + \tau] \times (0, \rho)$ (in polar coordinates centered at x). Lemmas 10 and 11 assert that the spoke contains a sequence of points inside the cone as it approaches x , since the spoke passes through a point at a distance less than $(1 - \cos \beta_x)^{-1} \eta(t)t$ from $\gamma_{x, \hat{\theta}}(t)$ and thus inside the cone for t sufficiently small. By the continuity of the spoke, it must cross $\gamma_{x, \hat{\theta} + \tau}$ (or $\gamma_{x, \hat{\theta} - \tau}$) an infinite number of times, and Lemma 9 says that this is impossible unless $\tau = 0$. \blacksquare

4. MEDIATRICES IN SPHERES, THE PUGH-TANGERMANN EXAMPLE

Let M be a simply connected 2-dimensional Riemannian manifold of class C^r , $r \geq 3$ (a sphere). For $p \neq q \in M$ we already know that there are at most finitely many points $x \in L_{pq}$ such that L_{pq} is not locally homeomorphic to a real interval at x (that is, there are more than two wedges at x). Let us call such a point a *branching point* (and a *simple point* to a non-branching point). For a simple point x the Main Theorem asserts that L_{pq} has two well defined directions $\hat{\theta}_1, \hat{\theta}_2 \in S_x M$ at x (the bisectors of the two pre-wedges at x , or in other words, the equidistant set in $S_x M$ of the sets of minimizing directions from x to p and q respectively). We define the *deficiency angle at x* as

$$\text{def}_{L_{pq}}(x) = \left| \pi - |\hat{\theta}_1 - \hat{\theta}_2| \right|.$$

If $\text{def}_{L_{pq}}(x) \neq 0$ we say that x is a *Lipschitz singularity* of the mediatrix L_{pq} .



Pugh-Tangerman example. A very illustrative example is what has come to be known as *Pugh's Cigar C* embedded in \mathbb{R}^3 . It consists of the cylinder given by $x^2 + y^2 = 1$ plus two (unit) hemispheres glued to each of the boundaries. Let p be its North Pole $(0, 0, 1)$ and q its South Pole. The mediatrice is given by the equator at $z = 0$.

We can obtain an arbitrarily smooth cigar by smoothing around $|z| = \pm 1$. If we add a “bump” in the flat “northern” part of the cigar, keeping C as a smooth manifold, is easy to see that the mediatrice will develop a Lipschitz singularity near the closest point to the bump. Adding a sequence of bumps in the northern part that do not intersect each other we can create a cigar whose mediatrice will have countably many Lipschitz singularities.

In what follows we want to show that this is, in some sense, the worst possible behavior for mediatrices on spheres. This proposition is inspired on conversations in the 1990's by Pugh and Tangerman with Veerman at Rockefeller University.

Proposition 12. *Let M be a sphere of class C^r , $r \geq 3$ and $p \neq q \in M$. Then the mediatrice L_{pq} verifies*

- (1) L_{pq} is a Lipschitz simple closed curve (there are no branching points).
- (2) There are at most countably many Lipschitz singularities in L_{pq} . Moreover the deficiency angles form an absolutely summable sequence, that is

$$\sum_{x \in L_{pq}} \text{def}_{L_{pq}}(x) < \infty.$$

Proof. Simple closed follows from earlier results ([2]). The novelty here is the Lipschitz property. The fact that the mediatrice is a simple closed curve implies that at each $x \in L_{pq}$ there are exactly two spokes, and thus exactly two wedges. By our Main Theorem, the mediatrice is tangent to the bi-sectors of these two wedges, and thus is Lipschitz. This proves (1).

We saw that at a given point $x \in L_{pq}$ there are exactly two wedges each of which is bounded by two minimizing geodesics $\gamma_{x,p}$ and $\gamma_{x,q}$. Take the complement of the closure of these wedges. If it is the empty set, the mediatrice is differentiable at x . If not, then (a) that complement is a maximal wedge shaped set bounded by two minimizing geodesics from x to p , or (b) bounded two minimizing geodesics from x to q , or (c) both of these occur. In case (a) let us denote by $\mathcal{J}_{x,p}$ the open domain bounded by the indicated geodesic rays from x to p and by $\mu_{x,p}$ the opening

angle at x of that wedge-shaped figure. Similarly we have $\mathcal{J}_{x,q}$ and $\mu_{x,q}$ for case (b). Case (c) means that both $\mathcal{J}_{x,p}$ and $\mathcal{J}_{x,q}$ are non-empty. Notice that $\mathcal{J}_{x,p} = \emptyset$ if and only if $\#\Theta_{x,p} = 1$.

If we have two minimizing geodesics from $x_1, x_2 \in L_{pq}$ to p or q , by triangle inequality the intersection point would have to be strictly closer to the first endpoint than the second, thus by symmetry these geodesics do not intersect. Therefore the collection (as x runs on L_{pq}) of Jordan domains $\mathcal{J}_{x,p}, \mathcal{J}_{x,q}$ are pairwise disjoint. We conclude that there are at most countably many points $x \in L_{pq}$ such that $\#\Theta_{x,p}$ or $\#\Theta_{x,q}$ are different from 1. A direct examination yields that

$$2\text{def}_{L_{pq}}(x) = |\mu_{x,p} - \mu_{x,q}|$$

The Gauss-Bonnet Theorem applied to $\mathcal{J}_{x,p}$ gives

$$\int_{\mathcal{J}_{x,p}} k(w)dw = \mu_{x,p} + \alpha_{x,p},$$

where $\alpha_{x,p}$ is the angle of the landing geodesic rays make at p (note that $\mu_{x,p}$ is the angle they make at x), and $k(w)$ is the curvature at $w \in M$. We estimate

$$\begin{aligned} \sum_{x \in L_{pq}} \text{def}_{L_{pq}}(x) &< \frac{1}{2} \sum_{x \in L_{pq}} \left(\int_{\mathcal{J}_{x,p}} |k(w)|dw + \int_{\mathcal{J}_{x,q}} |k(w)|dw + |\alpha_{x,p}| + |\alpha_{x,q}| \right) \\ &\leq \frac{1}{2} \left(\int_M |k(w)|dw + 2\pi \right). \square \end{aligned}$$

Acknowledgments. Pilar Herreros acknowledges the financial support of the research project FONDECYT 11121125. Mario Ponce and J. J. P. Veerman acknowledge the financial support of the research project FONDECYT 1140988. Mario Ponce received partial financial support from the project PIA-CONICYT ACT-1103. J.J.P. Veerman's research was also partially supported by the European Union's Seventh Framework Program (FP7-BEGPOT-2012-2013-1) under grant agreement n316165.

REFERENCES

- [1] I. Adelstein, *Minimizing Closed Geodesics via Critical Points of the Uniform Energy*. Preprint arXiv:1406.0372 [math.DG], 2014.
- [2] J. Bernhard and J.J.P. Veerman, *The topology of surface mediatrices*. Topology Appl. **154** (2007), no. 1, 54-68.
- [3] L. Birbrair and D. Siersma, *Metric properties of conflict sets*. Houston J. Math. **35** (2009), no. 1, 73-80.
- [4] D. Burago, Y. Burago and S. Ivanov, *A course in metric geometry*. Graduate Studies in Mathematics, **33**. American Mathematical Society, Providence, RI, 2001.
- [5] H. Busemann, *The geometry of Geodesics*. Academic Press, Inc., New York, 1955.
- [6] M. do Carmo, *Riemannian geometry*. Birkhäuser Boston, Inc., Boston, MA, 1992.
- [7] J. Damon, *Smoothness and geometry of boundaries associated to skeletal structures. I. Sufficient conditions for smoothness*. Ann. Inst. Fourier (Grenoble) **53** (2003), no. 6, 1941-1985.
- [8] D. Gromoll, W. Klingenberg, and W. Meyer, *Riemannsche Geometrie im Grossen*. Lecture Notes in Mathematics **55**, Springer-Verlag, New York, 1968.
- [9] Hockin and Young, *Topology*, Addison-Wesley, 1961.
- [10] L. D. Loveland, *When midsets are manifolds*. Proc. Amer. Math. Soc. **61** (1976), no. 2, 353-360.
- [11] M. M. Peixoto and R. Thom, *Le point de vue énumératif dans les problèmes aux limites pour les équations différentielles ordinaires. I: Quelques exemples*. C. R. Acad. de Sc. Paris **I 303** (1986), 629-632. Also: Ibid: CR **I 307** (1988), 197-198.
- [12] M. Ponce and P. Santibáñez, *On equidistant sets and generalized conics: the old and the new*. Amer. Math. Monthly **121** (2014), no.1, 18-32.
- [13] M. A. F. Sanjuan, J. Kennedy, C. Grebogi and J. A. Yorke, *Indecomposable continua in dynamical systems with noise: fluid flow past an array of cylinders*, Chaos **7** (1997), no.1, 125-138.

- [14] J.J.P. Veerman and J. Bernhard, *Minimally separating sets, mediatrices, and Brillouin spaces*. Topology Appl. **153** (2006), no. 9, 1421-1433.
- [15] J.J.P. Veerman, M.M. Peixoto, A.C. Rocha and S. Sutherland, *Brillouin Zones*. Comm. Math. Phys. **212/3** (2000), 725-744.
- [16] J.B. Wilker, *Equidistant sets and their connectivity properties*. Proc. Amer. Math. Soc. **47** (1975), 446-452.
- [17] Y. Yomdin, *On the local structure of a generic central set*. Compositio Math. **43** (1981), no. 2, 225-238.

Pilar Herreros

Facultad de Matemáticas, Pontificia Universidad Católica de Chile

e-mail: pherrero@mat.puc.cl

Mario Ponce

Facultad de Matemáticas, Pontificia Universidad Católica de Chile

e-mail: mponcea@mat.puc.cl

J.J.P. Veerman

Department of Mathematics and Statistics, Portland State University, and

CCQCN, Dept. of Physics, University of Crete, 71003 Heraklion, Greece

e-mail: veerman@pdx.edu