

**Portland State University**  
**PDXScholar**

---

Regional Research Institute

Regional Research Institute

---

2016

## Family Aided Community Treatment For The Treatment Of Early Psychosis: A Proof Of Concept Study

Ryan P. Melton

*Portland State University*, [rymelton@pdx.edu](mailto:rymelton@pdx.edu)

Cass Dykeman

*Oregon State University*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://pdxscholar.library.pdx.edu/rri\\_facpubs](https://pdxscholar.library.pdx.edu/rri_facpubs)

 Part of the [Community Health Commons](#), [Other Mental and Social Health Commons](#), and the [Psychiatric and Mental Health Commons](#)

---

### Citation Details

Melton, R. P., & Dykeman, C. (2016). Family Aided Community Treatment for the Treatment of Early Psychosis: A Proof of Concept Study. *Community Mental Health Journal*, 52(6), 623–630. <http://doi.org/10.1007/s10597-015-9984-z>

This Post-Print is brought to you for free and open access. It has been accepted for inclusion in Regional Research Institute by an authorized administrator of PDXScholar. For more information, please contact [pdxscholar@pdx.edu](mailto:pdxscholar@pdx.edu).

# Community Mental Health Journal

## Family Aided Community Treatment For The Treatment Of Early Psychosis: A Proof Of Concept Study --Manuscript Draft--

<b>Manuscript Number:</b>	COMH-D-13-00156R1	
<b>Full Title:</b>	Family Aided Community Treatment For The Treatment Of Early Psychosis: A Proof Of Concept Study	
<b>Article Type:</b>	Article	
<b>Keywords:</b>	major psychotic disorders; prodromal syndromes; family aided community treatment; early psychosis	
<b>Corresponding Author:</b>	Cass Dykeman, PhD Oregon State University Corvallis, UNITED STATES	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	Oregon State University	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Ryan P. Melton, PhD	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Ryan P. Melton, PhD	
	Cass Dykeman, PhD	
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	unfunded research project	Ryan P. Melton
<b>Abstract:</b>	Major psychotic disorders are one of the leading causes of disability worldwide. If these conditions are identified early and treatment promptly implemented, the prognosis is improved. This study examined the impact of a yearlong family aided community treatment (FACT) intervention upon psychiatric symptoms. Psychiatric symptom scores improved with the FACT intervention. Improved training on early recognition for mental health clinicians, implementation of a specific treatment model in community settings and policy around treatment funding allocation are implications of this study.	
<b>Response to Reviewers:</b>	Please thank the blind reviewers for their excellent edits. Dr. Melton and I added them all in toto.	

Family Aided Community Treatment For The Treatment Of Early Psychosis:

A Proof Of Concept Study

Ryan P. Melton

Portland State University

Cass Dykeman

Oregon State University

Author Note

Ryan P. Melton, Regional Research Institute, Portland State University; Cass Dykeman, Counseling Academic Unit, Oregon State University.

This study was completed as part of the first author's dissertation.

Correspondence concerning this article should be addressed to Ryan P. Melton, Regional Research Institute (RRI), Portland State University, Suite 918  
1600 SW 4th Ave., Portland OR. 97201. Email: [rymelton@pdx.edu](mailto:rymelton@pdx.edu)

## FAMILY AIDED COMMUNITY TREATMENT

1

## Abstract

Major psychotic disorders are one of the leading causes of disability worldwide. If these conditions are identified early and treatment promptly implemented, the prognosis is improved. This study examined the impact of a yearlong family aided community treatment (FACT) intervention upon psychiatric symptoms. Psychiatric symptom scores improved with the FACT intervention. Improved training on early recognition for mental health clinicians, implementation of a specific treatment model in community settings and policy around treatment funding allocation are implications of this study.

*Keywords:* major psychotic disorders, prodromal syndromes

Family Aided Community Treatment for The Treatment of Early Psychosis:  
A Proof of Concept Study

Schizophrenia has a prevalence rate of 1 in 100 people. Onset usually begins in late adolescence or early adulthood (APA, 2000). Symptoms include hallucinations, delusions, confused thinking, and a range of cognitive deficits (APA, 2000). In its acute phase, a person afflicted with schizophrenia becomes unable to discern what is real from what is not and may act on incorrect or inaccurate information about the environment. The young person often loses the ability to participate in school and work or even to take care of basic needs. Families may mistakenly attribute the cause of these changes in a loved one to antisocial behavior or drug use. Friends drop away quickly as these youth isolate or behave strangely.

The fact that schizophrenia begins during teenage and young adult years makes early intervention critical. The illness interferes with key developmental tasks, including forming an identity, finishing school and beginning employment, taking on adult roles and responsibilities, and forming intimate relationships. Psychosis can impede these developmental processes in ways that will affect a person for life. The stress can lead to family dissolution and even homelessness. Unable to tell what is safe from what is not or what is real from what is not, the person with schizophrenia is at risk of accidental death, injury, or suicide. It is common for psychosis to lead to arrest and legal charges (Sale & Melton, 2010).

Unfortunately, these experiences are common because many people with schizophrenia do not get the right help in the early stages of psychosis (Harrigan, McGorry, & Krstev, 2003). Treatment programs and providers are often untrained and

1  
2  
3  
4 unprepared to provide appropriate support. Too often, a series of traumatizing crises and  
5  
6 inappropriate care lead to an involuntary hospital commitment, entry into the federal  
7  
8 disability system, or both (Edwards & McGorry, 2002). The longer it takes for a person  
9  
10 to get appropriate help, the more challenging recovery becomes.  
11  
12

13  
14 To address the problem that many people with schizophrenia do not get the right  
15  
16 help in the early stages of psychosis, the State of Oregon implemented Early Assessment  
17  
18 and Support Alliance (EASA) programs in 11 counties. These counties cover 60% of  
19  
20 Oregon's population. The first year case rate for this program ran between \$9,000 to  
21  
22 \$12,000 per annum. The year two case rate ran at two-thirds of the year one rate. These  
23  
24 case rates were consistent with those reported by Mihalopoulos, Harris, Henry, Harrigan,  
25  
26 and McGorry (2009).  
27  
28  
29

30  
31 A key thrust of EASA programming was providing a family aided community  
32  
33 treatment (FACT) for this at-risk population. FACT is a combination of four distinct  
34  
35 elements. These elements are: (a) psychoeducational multifamily groups, (b) assertive  
36  
37 community treatment, (c) psychotropic medications, and (d) supported employment and  
38  
39 education (McFarlane, Stastny, & Deakins, 1992; McFarlane, 2002). While theory and  
40  
41 research backs each FACT element, the question whether FACT treatment package is  
42  
43 effective is an open one. The present proof of concept study is a response to this open  
44  
45 question.  
46  
47  
48  
49

50  
51 The specific research question for this study was: "What is the impact of a family  
52  
53 aided community treatment upon the psychiatric symptoms of people aged 12-25 with  
54  
55 early psychosis?" Based upon the extant research on this topic, the null hypothesis ( $H_0$ )  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

was directional: A family aided community treatment will not improve psychiatric symptoms of people aged 12-25 with early psychosis.

## Methods

### Research Design

Archival data sets were analyzed using a single-group pre/post test design (Harris, et al. 2006). The analysis compared baseline and 12-month scores for the purpose of proof of concept of the FACT intervention (Eli Lilly and Company, n.d.).

### Participants

Archival data were analyzed from 8 participants who received and completed 12 months of the FACT intervention. The mean age was 19.6 ( $SD = 3.28$ ). Males comprised 75% of the participants. The sex difference is consistent with Hafner et al. (1993). The racial demographics included 75% White and 25% Hispanic, which is consistent with data from Brekke and Barrio (1997).

Participants were selected from referrals made to the Early Assessment and Support Alliance. Eligible referrals included residents, aged 12-25, of the Mid-Willamette Valley Region of Oregon. At initial referral the participants appeared to have early, low-intensity or low-frequency psychotic symptoms. If participants met the inclusion criteria (i.e., age, location, and level of symptoms), they and their families were asked if they would be interested in participating in a research study. If they agreed, they were asked to attend an orientation session, which covered the FACT intervention, informed consent, description of financial incentives provided, and scheduling of the Structured Interview for Prodromal Syndromes (SIPS). The final inclusion and exclusion criteria were determined by scores and syndromes of the SIPS. Inclusion criteria

1  
2  
3  
4 consisted of participants who met criteria for a first episode psychosis. These participants  
5  
6 presented with psychotic symptoms (at least one score of 6 on one of the positive  
7  
8 symptom scales from the Structured Interview for Prodromal Syndromes (SIPS), with  
9  
10 loss of insight, and treatment with anti-psychotic medication) for less than 30 days.  
11  
12

13  
14 Included participants were offered the FACT intervention. At baseline the data set  
15  
16 included 11 individuals who met criteria for inclusion. At the end of 12 months, 8  
17  
18 participants remained for final analysis (see Figure 1). There were no clinical or  
19  
20 demographics differences between the participants who remained in the research and  
21  
22 those who were not in the final analysis. The final age range for the participants who  
23  
24 remained was 16-25. A two-tailed t-distribution test did not find significant differences  
25  
26 in the Structured Interview for Prodromal Syndromes (SIPS) Positive, Negative,  
27  
28 Disorganized, or General symptoms baseline scores between participants who dropped  
29  
30 out and the participants who remained in the study.  
31  
32  
33  
34  
35

36 Some potential participants were excluded because the purpose of this research  
37  
38 project was to prevent the disabling effects of the *onset* of psychotic disorder. Those  
39  
40 excluded met at least one of the following criteria: (a) a current diagnosis of a psychotic  
41  
42 disorder at baseline, (b) a positive symptom score of 6 for over 30 days, (c) receiving  
43  
44 anti-psychotic medication at a dosage appropriate to treat a psychotic illness for over 30  
45  
46 days, (d) a score lower than a 6 on any of the Positive symptoms scales, (e) aged less than  
47  
48 12 years or over 25, (f) an IQ less than 70 based on school records, (g) living outside the  
49  
50 Mid-Willamette Valley region, (h) not an English speaker, and/or neither parent is an  
51  
52 English speaker, (i) incarcerated, (j) psychotic symptoms due solely to acute  
53  
54 psychoactive substance toxicity, or (k) clear organic etiology of psychotic symptoms.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## Measures

**Structured Interview for Prodromal Syndromes (SIPS).** Participants received the SIPS at baseline and again at 12 months. The SIPS is a semi-structured diagnostic interview including five components: (a) the 19-point Scale of Prodromal Symptoms, (b) a version of the Global Assessment of Functioning with well-defined anchor points, (c) a Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR, 2000) schizotypal personality disorder checklist, (d) a family history of mental illness, and (e) a checklist for the Criteria of Prodromal Syndromes (Miller et al., 2002). The five components are used to determine if the person meets criteria of a Psychosis Risk Syndrome, indicating lower risk or a presence of psychotic symptoms (greater risk). SIPS questions include questions determined valid for predicting conversion to schizophrenia. For example, an item exploring unusual thought content asks the question, “Have you felt that you are not in control of your own ideas or thoughts?”

People who meet criteria for a prodromal syndrome have a positive, predictive value of conversion to psychosis related to schizophrenia: 43% at 6 months, 50% at 12 months, 62% at 18 months and 67% at 24 months. There was a significant relationship of diagnostic status at baseline to outcomes at 6 months and 12 months (Miller et al., 2002).

Independent research assistants who were blind to which participants received the FACT intervention conducted all interviews. Research assistants received regular inter-rater reliability checks by a Yale University certified rater expert in scoring the SIPS. SIPS raters who received training from a certified rater yielded an inter-rater reliability score of  $k = .81$  (Miller et al., 2002). To maintain inter-rater reliability, research assistants were asked to score 10-videotaped SIPS interviews. Reliability was measured

1  
2  
3  
4 by computing the intraclass correlation of raters' scores with criterion scores of  
5  
6 experienced researcher trained by Dr. Tandy Miller, one of the developers of the  
7  
8 Structured Interview for Prodromal Syndromes instrument. In a prior study using the  
9  
10 SIPS, this format was used to establish reliability (McFarlane et al., 2010). The intraclass  
11  
12 correlation was .81 for the positive symptom scale and .81 for the negative symptom  
13  
14  
15  
16  
17 scale.

18  
19 The Structured Interview for Prodromal Syndromes 19 item instrument broken  
20  
21 into four domains; positive (5 items), negative (6 items), disorganized (4 items) and  
22  
23 general symptoms (4 items). Each item is rated on a seven point fully-anchored Likert  
24  
25 scale. The end anchors are Absent (0) to Extreme (6). Any score in the positive  
26  
27 symptoms domain between a 3 and 5 is indicative of sub-clinical psychosis, whereas a 6  
28  
29 is considered psychotic (Miller et al., 1999). For the purposes of this study, an average  
30  
31 score of each domain was calculated at baseline and at the end of the study to determine  
32  
33  
34  
35  
36 improvement over the course of the treatment.

37  
38 **Positive and Negative Syndrome Scale (PANSS).** Participants also received the  
39  
40 Positive and Negative Syndrome Scale (PANSS). The PANSS is a 30-item researcher  
41  
42 rated scale used to determine severity of positive and negative symptoms of psychosis, in  
43  
44 addition to other excitatory, depressive, and cognitive psychiatric symptoms (PANSS  
45  
46 Institute, 2015). Higher scores indicate a greater severity of psychiatric symptoms.  
47  
48 PANSS construct, discriminative, convergent, and predictive validity is supported by  
49  
50  
51  
52 longitudinal, multidimensional, and psycho-pharmacological research in people with  
53  
54 acute and chronic schizophrenia. In two studies of 37 acute and 47 chronic people with  
55  
56  
57 schizophrenia, the criterion-related validity of the PANSS used for typological distinction  
58  
59  
60  
61  
62  
63  
64  
65

was supported (Kay, Opler, & Lindenmayer, 1988; Lindenmayer, Kay, & Opler, 1984). Studies specific to young people with schizophrenia found that high negative scores in the early presentation co-varied with: (a) lesser incidence of psychosis related to schizophrenia and (b) greater incidence of a psychosis related to a mood disorder (Kay & Opler, 1987; Lindenmayer, Kay, & Friedman, 1986).

For the present study, the Positive Symptom Subscale and Negative Symptom Subscale were used. Both subscales have seven items. Each item is scored on a seven-point scale. The range of the scores are as followed; 1 = absent, 2 = minimal, 3 = mild, 4 = moderate, 5 = moderate- severe, 6 = severe, and 7 = extreme (Kay, Fliszbein, & Opler, 1987). Similar to how the Structured Interview for Prodromal Symptoms, an average score of the positive and negative symptom subscale was calculated at baseline and at the end of the study to determine improvement over the course of the treatment.

**Global Functioning Scale (GFS).** Participants also received the Global Functioning Scale (GFS). The GFS produces a 10-item score measuring social and role functioning. Higher scores indicate higher functioning in each domain.

GFS studies on construct validity have found statistical significant correlations with the Strauss-Carpenter Outcome Scale. The Global Functioning Social Scale (Auther, Smith, & Cornblatt, 2006) of the GFS was correlated with the Strauss-Carpenter Outcome Social Contacts Scale ( $r = .70, p < .001$ ). The Global Functioning Role Scale (Niendam, Bearden, Johnson, & Cannon, 2006) was correlated with the Strauss-Carpenter Outcome Work/School Functioning Scale ( $r = .57, p < .001$ ). Researchers on the Global Functioning Scales also have found statistical significant correlations with the

Premorbid Adjustment Scale on the Global Functioning Social Scale with the Premorbid Adjustment Social Scale ( $r = .49, p < .001$ ) and on the Global Functioning Role Scale with the Premorbid Adjustment Role Scale ( $r = .68, p < .001$ ) (Cornblatt et al., 2007).

One study using 100 research clinicians at two different mental health research clinics found that GFS inter-rater reliability yielded .84 and .92 for the Social and Role functioning scores respectively (Cornblatt et al., 2007). The researcher who collected PANSS and GFS data did not undergo regular inter-rater reliability check as they did with the Structured Interview for Prodromal Syndromes. However, the positive and negative symptoms from the SIPS were used to score the PANSS so reliability is inferred.

For the purposes of this study, the scores from the Global Functioning: Social (GF: Social) and the Global Functioning: Role (GF: Role) subscales were employed. Each scale is a 0-10 scale, with higher scores indicating higher levels of functioning (Cornblatt et al., 2007). Each scale is scored via clinical opinion, similar to how the DSM-IV Global Assessment of Functioning Scale (GAF) is scored.

## **Treatment**

**Providers.** Participants who were assigned to the FACT intervention were given appointments with a master's level clinician who was responsible for providing and coordinating the FACT intervention elements. These clinicians had Master's degrees in counseling, psychology, social work or occupational therapy.

**Components.** FACT is a manualized treatment comprised of four components (see McFarlane et al., 1992; McFarlane, 2002; McFarlane et al., 2010). These

components are: (a) psychoeducational multifamily groups, (b) assertive community treatment, (c) psychotropic medications, and (d) supported employment and education.

***Psychoeducational multifamily groups.*** The multifamily group intervention had three part. These parts were: (1) joining with people and their families, (2) educational workshops, and (3) twice-monthly problem solving groups involving the person and their family (McFarlane, 2002).

***Assertive community treatment.*** Assertive Community Treatment is a comprehensive, biopsychosocial intervention with the intended goal of improving community functioning with the mentally ill. Components of Assertive Community Treatment include in vivo teaching of coping and problem-solving skills, 24-hour availability to respond to crisis, a small caseload (1–10 people), and a multidisciplinary team approach. As described, the original intention of the Assertive Community Treatment model was to serve high-risk mentally ill populations. Recently, there has been an increased awareness of its potential with first-episode and at-risk populations (McFarlane et al., 2010).

***Psychotropic medications.*** A board certified psychiatrist managed psychotropic medications. Persons were seen every one to three months depending on level of symptom severity. Medications were typically prescribed if symptoms indicated a conversion to psychosis or reached a level at which medications were indicated (McFarlane, 2002).

***Supported employment and education.*** Supported Employment is an evidenced-based, manualized treatment to support people with symptoms of mental illness return to or gain new employment. It involves an integration of employment and traditional

1  
2  
3  
4 mental health services. Clinicians using the supported employment model must offer  
5  
6 rapid job search of competitive employment emphasizing the person's preferences for  
7  
8 work goals. The model also involves clinicians offering support while the person is  
9  
10 employed and assisting with benefits planning. The model is implemented regardless of  
11  
12 the person's level of symptoms or substance use. The only inclusion is a person's  
13  
14 statement they want to work (Swanson & Becker, 2011).  
15  
16  
17

18  
19 Supported Education involves collaboration between clinician and educational  
20  
21 institutions, and teaching skills relevant to the person's current level of education. A  
22  
23 mental health clinician in collaboration with an occupational therapist completes this  
24  
25 intervention. The occupational therapist is responsible for evaluating the person's  
26  
27 functional and cognitive abilities and using that information to guide intervention.  
28  
29  
30

### 31 **Treatment Fidelity**

32  
33 All clinicians involved in implementing the FACT intervention received a  
34  
35 weeklong training on the elements of the model. The delivery of each clinical service  
36  
37 provided was documented using a secure web-based tracking program. Using  
38  
39 measurements of services people received was found to be a reliable and valid predictor  
40  
41 of outcomes with populations similar to the one used for this study (Cook et al., 2005).  
42  
43 Treatment fidelity scales exist for FACT elements of multi-family group treatment and  
44  
45 supported employment. Certified trainers monitored fidelity of these treatments on a  
46  
47 monthly basis in collaboration with the clinicians that implemented these FACT  
48  
49 interventions.  
50  
51  
52

### 53 **Clinical Significance**

54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

The term “clinically significant change” has typically been defines as the extent to which an intervention moves a person outside the range of the dysfunctional population or within the range of the functional population (Jacobson & Truax, 1991). For the SIPS, if any of the Positive Symptoms scales dropped from a baseline score of 6, the participant would no longer be considered psychotic, hence producing a clinically significant change (Miller et al., 2002).

### **Data Analysis Plan and Procedures**

The data were obtained from the Early Detection and Intervention for the Prevention of Psychosis Program (EDIPPP) National Database (Robert Wood Johnson Foundation, 2014). To determine impact of the Family Aided Community Treatment (FACT) intervention, multiple repeated measures t-tests were used to compare the outcome variables of pre-test and post-test average positive and negative symptoms subscales scores of the Structured Interview for Prodromal Syndromes, The Positive and Negative Symptoms Scale, and the Global Functioning Scales. The post-tests were given at 12 months post baseline testing. An intent-to-treat approach was used to avoid the effects of treatment crossover inherent in the FACT model. This research design provided information about the potential effects of the FACT model rather than the effects of specific treatment within the model. The assumptions of a repeated measures t-test were met given the wide distributions of scores at baseline in all outcome variables.

### **Results**

The hypothesis posited there would be an improvement in psychiatric symptoms of participants with early psychosis receiving the FACT intervention. The alternative hypothesis ( $H_1$ ) was supported for all outcome variables except role functioning (see

Table 1). There was a change in average role functioning score indicating improvement but the change was not statistically significant. In almost every scale, the standard deviation was smaller post-intervention. This smaller standard deviation was another indicator that the FACT intervention produced change.

### Discussion

Schizophrenia and major psychotic disorders are conditions with historically poor outcomes, both at a person and system level. These outcomes have produced a sense of entrenched pessimism among clinicians treating people with schizophrenia and major psychotic disorders. The present study indicates that using an intensive, multi-discipline treatment model with people in the very early stages of a major psychotic disorder can improve psychiatric symptoms and functioning.

Using a proof of concept design with multiple repeated measures t-test, eight first-episode psychotic disorder participants engaged in the Family Aided Community Treatment (FACT) protocol in order to examine if psychiatric symptoms scale scores improved post-intervention. The alternative hypothesis was supported. The FACT intervention decreased psychiatric symptom scores in this population.

Beyond statistical significance, the intervention was clinically efficacious. The SIPS measure defines a person as psychotic when scored as a six on any item of the Positive Symptoms Scale. At baseline each individual had at least one score of six on at least one of the five Positive Symptoms Scale. At 12 months no participants had a single score of 6. This result, in combination with the statistical significant changes, implies that FACT intervention resulted in the participants moving from inside the diagnostic range of dysfunctional and psychotic to outside of that range.



1  
2  
3  
4 This study had eight potential limitations that should be noted. First, the sample  
5  
6 was composed of participants who were help-seeking and consented to a major research  
7  
8 study. People with major psychotic disorders, specifically schizophrenia, often do not  
9  
10 seek help (Amador, 2007; van Os, 2003) raising the question as to whether this study  
11  
12 reflects a representative sample of people with the condition, who generally do not seek  
13  
14 help. However, given this specific associated feature of the condition, the design and  
15  
16 selection of participants for this study is an effective way to determine if the FACT  
17  
18 model offers predictive validity. An essential component of the FACT model is assertive  
19  
20 community treatment (ACT), which has proven outcomes indicating improved  
21  
22 engagement in treatment among individuals with severe mental illness including those  
23  
24 who do not seek help (Substance Abuse and Mental Health Services Administration,  
25  
26  
27  
28  
29  
30  
31 2008).

32  
33 The second limitation involves the history threat to internal validity. The use of  
34  
35 multiple comparisons itself creates the chances of a Type I error. The history threat was  
36  
37 present given it was possible an event outside of the treatment that occurred during the  
38  
39 course of the participants' treatment resulted in the achieved results. Participants in this  
40  
41 study did not enroll simultaneously, they were enrolled as they were referred and met  
42  
43 criteria for the research. This reduces the likelihood of a history threat.  
44  
45  
46  
47

48 A third limitation centered on the possibility that the participants' outcomes  
49  
50 would have been achieved as a natural course of their development (i.e., maturation).  
51  
52 Baseline analysis of the participants found that they were a homogeneous group in terms  
53  
54 of symptomology, age, gender and race. Also, the baseline demographics are consistent  
55  
56 with persons who are diagnosed with psychosis. If maturation were present, persons  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 outside of this study with early psychosis would also have experienced a reduction in  
5  
6 symptoms that has not been the case.  
7

8  
9 A fourth limitation involved a pre-posttest design threat to internal validity is the  
10 testing threat. Consideration of this threat involve the question: "Did the participants  
11 experience of being administered the SIPS at baseline prepare them for improved  
12 responses at 12-month administration of the SIPS?" To address this threat, the PANSS  
13 and SIPS were used in tandem. The PANSS is researcher rated and hence not a risk for a  
14 testing threat. The PANSS scores were similar in terms of level of psychosis to the SIPS  
15 both at baseline and at 12 months. The advantage of using the same two instruments  
16 measuring the same construct also reduces the risk of the instrumentation threat, which  
17 would imply the participants change in symptomology would be due to the alternative  
18 forms of the same tests intended to be equivalent but are not.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 This study had a small number of participants lost to attrition ( $n = 3$ ). No  
34 statistically significant clinical or demographics differences were found between  
35 participants who remained in the research and those who dropped out; this reduced the  
36 threat of attrition.  
37  
38  
39  
40  
41  
42

43 A seventh potential limitation could be the sample size ( $n = 8$ ). Yet, it is  
44 important to note that the incidence of schizophrenia is 7 to 8 of 100,000 (Drake, Haley,  
45 Akhtar, & Lewis, 2000). The sample was pulled from a population of 630,000. The  
46 number of participants is consistent with an expected incidence rate. To investigate this  
47 limitation, power analysis was conducted to assess the importance of sample size on the  
48 t-test statistics. For the majority of scales, post-hoc power analysis supported the t-test  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 results. In all these analyses an alpha of .05 and a power of .90 were used. The only  
5  
6 analysis that did not support the sample size was the Global Functioning Role Scale.  
7  
8

9         The eighth and final potential limitation that needs to be noted is the regression  
10 threat. Given that the all the participants had extreme scores at baseline and the sample  
11 was not randomized or large, regression cannot be ruled out. However, several  
12 participants' scores improved drastically. Four participants with average SIPS P scores of  
13 four or higher reduced their scores to less than one.  
14  
15  
16  
17  
18  
19  
20

21         The results suggest some potential modifications in practice. It demonstrates that  
22 there are tools that can easily be used in clinical practice for identifying early symptoms  
23 of psychotic disorders and that if the conditions are caught early there is a treatment  
24 model that is effective in reducing psychiatric symptoms and improving functioning. The  
25 positive results of this study may encourage practitioners of all disciplines to engage  
26 people with early psychotic conditions in treatment with a sense of hope and optimism as  
27 opposed to the historical beliefs of hopelessness and pessimism. It may also improve  
28 social justice efforts to reduce discrimination and stigma associated with psychotic  
29 disorders, and advocate for appropriate and evidenced base treatment that reduces  
30 symptoms to the person, reduces costs to systems and improve overall delivery of  
31 services.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48         There are two specific implications of this study for researchers. First, given the  
49 overall strengths of the findings after 12 months a follow-up study would be to explore  
50 the effects of the intervention for longer periods would add to the nomothetical net in this  
51 area . It may be possible that role functioning requires a longer treatment dose for the  
52 impact to be at a significant level.  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4           The second implication arises from the intent-to-treat approach to avoid the  
5  
6 effects of treatment crossover inherent in the FACT model. This research design provided  
7  
8 information about the potential effects of the FACT model rather than the effects of  
9  
10 specific treatment within the model. Implications for further research would be to  
11  
12 identify the active ingredients within the FACT model that could be randomized, such as  
13  
14 participation in multifamily psychoeducation groups, supported employment and  
15  
16 education and occupational therapy. Also, using the proof of concept model this study  
17  
18 has established a dose response relationship that could be used as the basis for a decision  
19  
20 to move forward with additional trials.  
21  
22  
23  
24

25  
26           This study examined the impact of a yearlong family aided community treatment  
27  
28 intervention upon psychiatric symptoms. Psychiatric symptom scores improved with the  
29  
30 intervention. The positive findings from the study demonstrated that there are reliable and  
31  
32 valid tools for identifying early symptoms of psychotic disorders and that if the  
33  
34 conditions are caught early there is a treatment model that is potentially effective in  
35  
36 reducing psychiatric symptoms and improving functioning. Hence, further research on  
37  
38 this intervention package is warranted.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- Amador, X. (2007). *I am not sick, I don't need help*. New York: Vida Press.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., text rev.). Washington, DC: Author.
- Auther, A. M., Smith, C. W. & Cornblatt, B. A. (2006). *Global Functioning: Social Scale (GF: Social)*. Glen Oaks, NY: Zucker Hillside Hospital.
- Brekke, B. J., & Barrio, C. (1997). Cross-ethnic symptom differences in schizophrenia: The influence of culture and minority status. *Schizophrenia Bulletin*, 23, 305-316.
- Cook, J. A., Leff, H. S., Blyler, C. R., Gold, P. B., Goldberg, R. W., Mueser, K. T., ... Burke-Miller, J. (2005). Results of a multisite randomized trial of supported employment interventions for individuals with severe mental illness. *Archives of General Psychology*, 62, 505-512. doi: 10.1192/apt.bp.107.003509
- Cornblatt, B. A., Auther, A. M., Niendam, T., Smith, C. W., Zinberg, J., Bearden, C. E., & Cannon, T. (2007). Preliminary findings for two new measures of social and role functioning in the prodromal phase of schizophrenia. *Schizophrenia Bulletin*, 33, 688-702. doi: 10.1093/schbul/sbm029
- Drake, R. J., Haley, C. J., Akhtar, S., & Lewis, S. W. (2000). Causes and consequences of duration of untreated psychosis in schizophrenia. *British Journal of Psychiatry*, 177, 511-515.
- Edwards, J., & McGorry, P. D. (2002). *Implementing early intervention in psychosis: A guide to establishing early psychosis services*. London: Martin Dunitz Ltd.
- Eli Lilly and Company. (n.d.). Terminology. Retrieved from

- <http://www.lillytrials.com/docs/terminology.html>
- Hafner, H., Riecher-Rossler, A., An Der Heiden, W., Maurer, K., Fatkenheuer, B., & Loffler, W. (1993). Generating and testing a causal explanation of the gender difference in age at first onset of schizophrenia. *Psychological Medicine, 23*, 925-940. doi: 10.1017/S0033291700026398
- Harrigan, S., McGorry, P., & Krstev, H. (2003). Does treatment delay in first-episode really matter? *Psychological Medicine, 33*, 97–110. doi: 10.1017/S003329170200675X
- Harris, A. D., McGregor, J. C., Perencevich, E. N., Furuno, J. P., Zhu, J., Peterson, D. E., & Finkelstein, J. (2006). The use and interpretation of quasi-experimental studies in medical informatics. *Journal of the American Medical Informatics Association, 13*, 16-23. doi:10.1197/jamia.M1749
- Jacobson, N., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology, 59*, 12-19. doi: 10.1037//0022-006X.59.1.12
- Kay, S. R., , A., & Opfer, L. A. (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin, 13*, 261-276.
- Kay, S. R., & Opler, L. A. (1987). The positive-negative dimension in schizophrenia: Its validity and significance. *Psychiatric Developments, 5*, 79-103.
- Kay, S. R., Opler, L. A., & Lindemayer, J. P. (1988). Reliability and validity of the positive and negative syndrome scale for schizophrenics. *Psychiatry Research, 23*, 99-110.

- 1  
2  
3  
4 Lindenmayer, J. P., Kay, S. R., & Friedman, C. (1986). Negative and positive  
5  
6 schizophrenic syndromes after the acute phase: A prospective follow-up.  
7  
8  
9 *Comprehensive Psychiatry*, 27, 276-286. doi: 10.1016/0010-440X(86)90003-9  
10  
11 Lindenmayer, J. P., Kay, S.R., & Opler, L. A. (1984). Positive and negative subtypes in  
12  
13 acute schizophrenia. *Comprehensive Psychiatry*, 25, 455-464.  
14  
15  
16 doi.org/10.1016/0010-440X(84)90080-4  
17  
18  
19 McFarlane, W. R., Cook, W. L., Downing, D., Verdi, M. B., Woodberry, K. A., & Ruff,  
20  
21 A. (2010). Portland identification and early referral: A community-based system  
22  
23 for identifying and treating youths at high risk of psychosis. *Psychiatric Services*,  
24  
25 61, 512–515. doi: 10.1176/appi.ps.61.5.512  
26  
27  
28 McFarlane, W. R. (2002). *Multifamily groups in the treatment of severe psychiatric*  
29  
30 *disorders*. New York, NY: Guilford.  
31  
32  
33 McFarlane, W. R., Stastny, P., & Deakins, S. (1992). Family-aided assertive community  
34  
35 treatment: A comprehensive rehabilitation and intensive case management  
36  
37 approach for persons with schizophrenic disorders. *New Directions in Mental*  
38  
39 *Health Services*, 53, 43–54. doi: 10.1002/yd.23319925306  
40  
41  
42  
43 McGlashan, T. H., Miller, T., Woods, S., Rosen, J., Hoffman, R., & Davidson, L. (2003).  
44  
45 *The structured interview for prodromal syndromes*. New Haven, CT: PRIME  
46  
47 Research Clinic.  
48  
49  
50 Mihalopoulos, C., Harris, M., Henry, L., Harrigan, S., & McGorry, P. (2009). Is early  
51  
52 intervention in psychosis cost-effective over the long term?. *Schizophrenia*  
53  
54 *bulletin*, 35, 909-918. doi: 10.1093/schbul/sbp054  
55  
56  
57  
58 Miller, T. J., McGlashan, T. H., Rosen, J. L., Cadenhead, K., Ventura, J., McFarlane, W.,  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 et al.... Woods, S. W. (2003). Prodromal assessment with the Structured  
5  
6 Interview for Prodromal Syndromes and the Scale of Prodromal Symptoms:  
7  
8 Predictive validity, interrater reliability, and training to reliability. *Schizophrenia*  
9  
10 *Bulletin, 29*, 703–715.  
11  
12  
13  
14 Miller, T. J., McGlashan, T. H., Rosen, J. L., Somjee, L., Marcovitch, P. J., Stein, K., &  
15  
16 Woods, S. W. (2002). Prospective diagnosis of the initial prodrome for  
17  
18 schizophrenia based on the Structured Interview for Prodromal Syndromes:  
19  
20 Preliminary evidence of interrater reliability and predictive validity. *American*  
21  
22 *Journal of Psychiatry, 159*, 863–865. doi: 10.1176/appi.ajp.159.5.863  
23  
24  
25  
26 Miller, T. J., McGlashan, T. H., Woods, S. W., Stein, K., Driesen, N., Corcoran, C. M., ...  
27  
28 & Davidson, L. (1999). Symptom assessment in schizophrenic prodromal states.  
29  
30 *Psychiatric Quarterly, 70*, 273-287. doi: 10.1023/A:1022034115078  
31  
32  
33  
34 Niendam, T. A., Bearden, C. E., Johnson, J. K. & Cannon, T. D. (2006). *Global*  
35  
36 *Functioning: Role Scale (GF: Role)*. Los Angeles, CA: University of California,  
37  
38 Los Angeles.  
39  
40  
41 PANSS Institute. (2015). *The PANSS and other scales*. NY: Author. Retrieved from:  
42  
43 <http://www.panss.org/content/panss-and-other-scales>  
44  
45  
46 Robert Wood Johnson Foundation. (2014). *Early detection and intervention for the*  
47  
48 *prevention of psychosis in adolescents and young adults*. Princeton, NJ: Author.  
49  
50 Retrieved from  
51  
52 [http://www.rwjf.org/content/dam/farm/reports/program\\_results\\_reports/2014/rwjf](http://www.rwjf.org/content/dam/farm/reports/program_results_reports/2014/rwjf)  
53  
54 408484  
55  
56  
57  
58 Sale, T., & Melton, R. (2010). Early psychosis intervention in Oregon: Building a  
59  
60  
61  
62  
63  
64  
65



positive future for this generation. *Focal Point*, 24, 25–28.

Substance Abuse and Mental Health Services Administration. (2008). *Assertive Community Treatment (ACT) evidence-based practices (EBP) Kit*. Rockville, MD: Author.

Swanson, S. J., & Becker, D. R. (2011). *Updated and expanded supported employment: Applying the individual placement and support (IPS) model to help clients compete in the workforce*. Center City, MN: Dartmouth PRC-Hazelden.

Van Os, J. (2003). Is there a continuum of psychotic experiences in the general population? *Epidemiologia e Psichiatria Sociale*, 12, 242-252. doi: <http://dx.doi.org/10.1017/S1121189X00003067>

Table 1

*FACT intervention outcome variables: Inferential Statistics (N = 8)*

Variables	M	SD	t	H <sub>0</sub> (one-tail)
SIPS P Scale	3.60	0.70	14.53	Reject
Pre	3.93	0.59		$p < .05$
Post	0.33	0.41		
SIPS N Scale	1.27	1.04	3.46	Reject
Pre	2.52	1.21		$p < .05$
Post	1.25	0.71		
SIPS D Scale	1.81	1.22	4.22	Reject
Pre	2.09	1.21		$p < .05$
Post	0.28	0.34		
SIPS G Scale	2.22	0.96	6.55	Reject
Pre	3.28	0.66		$p < .05$
Post	1.06	1.19		
PANSS P Scale	1.77	0.47	10.63	Reject
Pre	2.96	0.56		$p < .05$
Post	1.20	0.34		
PANSS N Scale	1.00	0.59	4.82	Reject
Pre	2.27	0.73		$p < .05$
Post	1.27	0.30		
GFS-Social	1.13	0.84	3.81	Reject
Pre	6.00	1.60		$p < .05$
Post	7.13	1.36		
GFS-Role	0.50	4.00	0.35	Do not Reject
Pre	4.13	2.26		$p > .05$
Post	4.63	3.29		