

**HHS PUBLIC ACCESS**

Author manuscript

*Clin Pharmacol Ther.* Author manuscript; available in PMC 2017 January 01.

Published in final edited form as:

*Clin Pharmacol Ther.* 2016 January ; 99(1): 36–37. doi:10.1002/cpt.161.**Clinical Pharmacogenetics Implementation Consortium (CPIC) Guidelines for Human Leukocyte Antigen B (HLA-B) Genotype and Allopurinol Dosing: 2015 update****Yoshiro Saito<sup>1</sup>, Lisa K Stamp<sup>2</sup>, Kelly E. Caudle<sup>3</sup>, Michael Hershfield<sup>4</sup>, Ellen M. McDonagh<sup>5</sup>, John T. Callaghan<sup>6,7,8</sup>, Wichittra Tassaneeyakul<sup>9</sup>, Taisei Mushiroda<sup>10</sup>, Naoyuki Kamatani<sup>11</sup>, Barry R. Goldspiel<sup>12</sup>, Elizabeth J. Phillips<sup>13,14</sup>, Teri E. Klein<sup>5</sup>, and Ming Ta Michael Lee<sup>15,16,17</sup>**<sup>1</sup>Division of Medicinal Safety Science, National Institute of Health Sciences, Kamiyoga, Setagaya, Tokyo, Japan <sup>2</sup>Department of Medicine, University of Otago, Christchurch, New Zealand<sup>3</sup>Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN<sup>4</sup>Departments of Medicine and Biochemistry, Duke University School of Medicine, Durham, NC, USA <sup>5</sup>Department of Genetics, Stanford University Medical Center, Stanford, California, USA<sup>6</sup>ACOS for Research, Department of Veterans Affairs Medical Center, Indianapolis, Indiana, USA<sup>7</sup>Department of Medicine, Indiana University School of Medicine, Indianapolis, Indiana, USA<sup>8</sup>Department of Pharmacology/Toxicology, Indiana University School of Medicine, Indianapolis, Indiana, USA<sup>9</sup>Department of Pharmacology, Research and Diagnostic Center for Emerging Infectious Diseases, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand<sup>10</sup>Laboratory for Pharmacogenetics, RIKEN, Center for Genomic Medicine, Yokohama, Japan<sup>11</sup>Institute of Data Analysis, StaGenCo.Ltd., Tokyo, Japan <sup>12</sup>Pharmacy Department, National Institutes of Health Clinical Center, Bethesda, Maryland, USA <sup>13</sup>Division of infectious diseases, Institute of Immunology and Infectious Disease, Murdoch University, Murdoch, Western Australia<sup>14</sup>Division of Infectious Diseases, Vanderbilt University Medical Center, Nashville, TN<sup>15</sup>Laboratory for International Alliance on Genomic Research, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan <sup>16</sup>National Center for Genome Medicine, Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan <sup>17</sup>School of Chinese Medicine, China Medical University, Taichung, Taiwan

Corresponding Author: Ming Ta Michael Lee, RIKEN Center for Genomic Medicine, 1-7-22, Suehiro-cho, Tsurumi-ku, Yokohama, Japan 14101, Phone number: +81-45-503-9589, mikelee@src.riken.jp.

**Conflict of Interest:** T.E.K. is a stockholder for Personalis Inc. L.K.S has been a consultant for Astra Zeneca. As of January 2015, E.M.M. is Lead Scientific Curator of Genomics England, Queen Mary University. M.S.H. is a co-inventor of Pegloticase (Krystexxa®) and receives royalties from Crealta Pharmaceuticals. The other authors declare no conflicts of interest.

**Disclaimer:** Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines reflect expert consensus based on clinical evidence and peer-reviewed literature available at the time they are written, and are intended only to assist clinicians in decision-making, as well as to identify questions for further research. New evidence may have emerged since the time a guideline was submitted for publication. Guidelines are limited in scope and are not applicable to interventions or diseases not specifically identified. Guidelines do not account for all individual variation among patients and cannot be considered inclusive of all proper methods of care or exclusive of other treatments. It remains the responsibility of the health care provider to determine the best course of treatment for the patient. Adherence to any guideline is voluntary, with the ultimate determination regarding its application to be solely made by the clinician and the patient. CPIC assumes no responsibility for any injury to persons or damage to property related to any use of CPIC's guidelines, or for any errors or omissions.

## Abstract

The Clinical Pharmacogenetics Implementation Consortium (CPIC) Guidelines for *HLA-B\*58:01* Genotype and Allopurinol Dosing was originally published in February 2013. We reviewed recent literature and concluded that none of the evidence would change the therapeutic recommendations in the original guideline; therefore, the original publication remains clinically current. However, we have updated the supplemental material and included additional resources for applying CPIC guidelines into the electronic health record. Up-to-date information can be found at PharmGKB (<http://www.pharmgkb.org>).

## Keywords

allopurinol; hypersensitivity; pharmacogenetics; HLA-B; CPIC

---

The Clinical Pharmacogenetics Implementation Consortium (CPIC) of the Pharmacogenetics Research Network (<http://www.pgrn.org>) and the Pharmacogenomics Knowledge Base (PharmGKB, <http://www.pharmgkb.org>) provides peer-reviewed, evidence-based, freely accessible genotype-based drug guidelines to help clinicians understand how available genetic test results could be used to optimize drug therapy(1). CPIC guidelines undergo continuous peer review and information that would modify prescribing recommendations pertaining to gene specific alleles and nomenclature are updated periodically on the PharmGKB website. Furthermore, published CPIC guidelines are currently systematically reviewed for updates periodically.

The CPIC Guideline for *HLA-B* Genotype and Allopurinol Dosing was originally published in February 2013 (2). To update this guideline, we conducted a focused review of the literature published between 1966 to October 2014 on *HLA* genotype and allopurinol (see Supplemental material on-line). Our inclusion criteria for this guideline update also included other HLA variants besides HLA-B. The *HLA-B\*58:01* allele frequency tables (Supplemental Tables S1 and S2) have also been updated. The literature review yielded 26 relevant primary studies showing association between *HLA-B\*58:01* and allopurinol SCAR (Supplemental Table S3). In addition, 12 studies showed associations for *HLA-A\*33:03* (7 studies) or *HLA-C\*03:02* (5 studies) (Supplemental Table S3). However, the strength of the evidence for *HLA-A\*33:03* and *HLA-C\*03:02* did not warrant inclusion in this update (please see “other considerations” in the supplementary materials). We found no new evidence that would change our original recommendations for *HLA-B\*58:01* and allopurinol dosing; therefore, the original guideline publication and recommendation remains current (2). The 2012 American College of Rheumatology Guidelines for Management of Gout (3) recommends testing for the *HLA-B\*58:01* allele in selected subpopulations with elevated risk for allopurinol hypersensitivity syndrome (individuals of Korean decent with stage 3 or worse chronic kidney disease, and those of Han-Chinese or Thai decent) prior to initiation of the drug.

All new and updated CPIC guidelines will address dosing in pediatrics. Although none of the evidence linking *HLA-B\*58:01* to allopurinol hypersensitivity was conducted in

children, there is no reason to suspect that children positive for *HLA-B\*58:01* would be at less risk of allopurinol hypersensitivity reactions than adults positive for *HLA-B\*58:01*.

CPIC guidelines are designed to help clinicians use genetic information to optimize drug therapy, and to do this effectively, pharmacogenetic information must be incorporated into electronic health records (EHRs) with clinical decision support (CDS) (4-6). To provide additional resources for applying CPIC guidelines into the EHR, CPIC created an informatics working group focused on supporting the adoption of CPIC guidelines within a clinical electronic environment. This guideline provides these clinical implementation resources as part of the supplementary material, which include workflow diagrams that illustrate the storage of a pharmacogenetic result in an EHR and the design of CDS alerts (Supplemental Figures S1 and S2). New tables that correspond to these workflow diagrams are provided that translate genotype test results into an interpreted phenotype (Supplemental Tables S4-S8). These tables provide summary genotype/phenotype terms, example text for documentation in the EHR and point-of-care alerts, and cross-references for drug and gene names to widely-used terminologies and standardized nomenclature systems.

We recognize that each organization and EHR vendor may have different requirements and preferences for implementing pharmacogenetics within a given electronic environment. The intent of these resources is to synthesize foundational knowledge that provides a common starting point for clinical implementation so that individual organizations do not have to create a similar knowledge base for each new gene/drug pair that is implemented. Future CPIC guidelines and guideline updates will provide similar resources to guide the implementation of gene-drug pairs into the EHR.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We acknowledge the critical input of members of CPIC of the Pharmacogenomics Research Network (PGRN) particularly Dr. Mary V. Relling (St Jude Children's Research Hospital) and the CPIC informatics working group. This work was funded by the National Institutes of Health (NIH)/National Institute of General Medical Science (NIH/NIGMS) (PAAR4Kids (U01 GM92666), PharmGKB (R24 GM61374) and U01 HL0105198), the Health Labor Sciences Research Grants from MHLW of Japan and the National Science and Technology Development Agency, Thailand.

## References

1. Relling MV, Klein TE. CPIC: Clinical Pharmacogenetics Implementation Consortium of the Pharmacogenomics Research Network. *Clinical pharmacology and therapeutics*. 2011; 89:464–7. [PubMed: 21270786]
2. Hershfield MS, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for human leukocyte antigen-B genotype and allopurinol dosing. *Clinical pharmacology and therapeutics*. 2013; 93:153–8. [PubMed: 23232549]
3. Khanna D, et al. 2012 American College of Rheumatology guidelines for management of gout. Part 2: therapy and antiinflammatory prophylaxis of acute gouty arthritis. *Arthritis care & research*. 2012; 64:1447–61. [PubMed: 23024029]

4. Shuldiner AR, et al. The Pharmacogenomics Research Network Translational Pharmacogenetics Program: overcoming challenges of real-world implementation. *Clinical pharmacology and therapeutics*. 2013; 94:207–10. [PubMed: 23588301]
5. Wilke RA, et al. The emerging role of electronic medical records in pharmacogenomics. *Clinical pharmacology and therapeutics*. 2011; 89:379–86. [PubMed: 21248726]
6. Kullo IJ, Jarvik GP, Manolio TA, Williams MS, Roden DM. Leveraging the electronic health record to implement genomic medicine. *Genetics in medicine : official journal of the American College of Medical Genetics*. 2013; 15:270–1. [PubMed: 23018749]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript