



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers

Faculty of Science, Medicine and Health

2017

Antioxidant, Cytotoxic and α -Glucosidase Inhibitory Activities of Compounds isolated from the Twig Extracts of *Maclura fruticosa*

Isaraporn Polbuppha
Mae Fah Luang University

Wisanu Maneerat
Mae Fah Luang University

Tawanun Sripisut
Mae Fah Luang University

Thunwadee Ritthiwigrom
Chiang Mai University

Sarot Cheenpracha
University of Phayao

See next page for additional authors

Publication Details

Polbuppha, I., Maneerat, W., Sripisut, T., Ritthiwigrom, T., Cheenpracha, S., Pyne, S. G., Muanprasat, C., Seemakhan, S., Borwornpinyo, S. & Laphookhieo, S. (2017). Antioxidant, Cytotoxic and α -Glucosidase Inhibitory Activities of Compounds isolated from the Twig Extracts of *Maclura fruticosa*. *Natural Product Communications: an international journal for communications and reviews*, 12 (7), 1073-1076.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Antioxidant, Cytotoxic and α -Glucosidase Inhibitory Activities of Compounds isolated from the Twig Extracts of *Maclura fruticosa*

Abstract

The first phytochemical investigation of the twig extracts of *Maclura fruticosa* led to the isolation and identification of a new xanthone, maclurafruticosone (1), together with 14 known compounds (2-15). All compounds were elucidated using spectroscopic methods as well as through comparisons made with data reported in the literature. Some isolated compounds were evaluated for their antioxidant, α -glucosidase inhibitory and cytotoxic activities. Compound 4, 6 and 7 showed significant antioxidant activity against DPPH radicals with IC_{50} values ranging from 7.45-16.12 μ M. Compound 4 also exhibited potent activity against ABTS^{•+} scavenging activity with an IC_{50} value of 0.55 ± 0.01 μ M which was better than positive control (ascorbic acid, IC_{50} 2.35 ± 0.17 μ M). Compound 12 showed significant α -glucosidase inhibitory activity with an IC_{50} value of 0.02 ± 0.37 mM. Compounds 5, 6 and 12 showed weak cytotoxic activities against a colon cancer cell line with IC_{50} values ranging from 22.35-47.62 μ M.

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Polbuppha, I., Maneerat, W., Sripisut, T., Ritthiwigrom, T., Cheenpracha, S., Pyne, S. G., Muanprasat, C., Seemakhan, S., Borwornpinyo, S. & Laphookhieo, S. (2017). Antioxidant, Cytotoxic and α -Glucosidase Inhibitory Activities of Compounds isolated from the Twig Extracts of *Maclura fruticosa*. *Natural Product Communications: an international journal for communications and reviews*, 12 (7), 1073-1076.

Authors

Isaraporn Polbuppha, Wisanu Maneerat, Tawanun Sripisut, Thunwadee Ritthiwigrom, Sarot Cheenpracha, Stephen G. Pyne, Chatchai Muanprasat, Sawinee Seemakhan, Suparek Borwornpinyo, and Surat Laphookhieo

Antioxidant, Cytotoxic and α -Glucosidase Inhibitory Activities of Compounds isolated from the Twig Extracts of *Maclura fruticosa*

Isaraporn Polbuppha^a, Wisanu Maneerat^a, Tawanun Sripisut^{b,c}, Thunwadee Limtharakul^d (née Ritthiwigrom), Sarot Cheenpracha^e, Stephen G. Pyne^f, Chatchai Muanprasat^g, Sawinee Seemakha^h, Suparek Borwornpinyo^h and Surat Laphookhieo^{a,*}

^aNatural Products Research Laboratory, School of Science, Mae Fah Luang University, Tasud, Muang, Chaing Rai 57100, Thailand

^bSchool of Cosmetic Science, Mae Fah Luang University, Chaing Rai 57100, Thailand

^cPhytocosmetics and Cosmeceuticals Research Group, Mae Fah Luang University, Chiang Rai 57100, Thailand

^dDepartment of Chemistry, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

^eSchool of Science, University of Phayao, Phayao 56000, Thailand

^fSchool of Chemistry, University of Wollongong, Wollongong, New South Wales 2522, Australia

^gDepartment of Physiology, Faculty of Science, Mahidol University, Rajathevi, Bangkok 10400, Thailand

^hExcellent Center of Drug Discovery, Faculty of Science, Mahidol University, Rajathevi, Bangkok 10400, Thailand

surat.lap@mfu.ac.th

Received: May XX, 2017; Accepted: XX, 2017

The first phytochemical investigation of the twig extracts of *Maclura fruticosa* led to the isolation and identification of a new xanthone, maclurafruticosone (1), together with 14 known compounds (2-15). All compounds were elucidated using spectroscopic methods as well as through comparisons made with data reported in the literature. Some isolated compounds were evaluated for their antioxidant, α -glucosidase inhibitory and cytotoxic activities. Compound 4, 6, and 7 showed significant antioxidant activity against DPPH radicals with IC₅₀ values ranging from 7.45–16.12 μ M. Compound 4 also exhibited potent activity against ABTS^{•+} scavenging activity with an IC₅₀ value of 0.55 \pm 0.01 μ M which was better than positive control (ascorbic acid, IC₅₀ 2.35 \pm 0.17 μ M). Compound 12 showed significant α -glucosidase inhibitory activity with an IC₅₀ value of 0.02 \pm 0.37 mM. Compounds 5, 6 and 12 showed weak cytotoxic activities against a colon cancer cell line with IC₅₀ values ranging from 22.35–47.62 μ M.

Keywords: *Maclura fruticosa*, Moraceae, xanthone, isoflavone, antioxidant activity, α -glucosidase inhibitory, cytotoxicity.

The *Maclura* genus belongs to the Moraceae family which is widely distributed in Asia, America, Africa, Australia, and Europe. Many species of this genus have been used as traditional medicines as well as food, pesticides, and dyes [1-4]. For examples, the resin of *M. tinctoria* is used to treat toothache and also used as colorant and its fruit is edible [3]. The decoction of the root of *M. pomifera* had been used for the treatment of sore eyes [4]; the bark had been used for uterine haemorrhage treatment [4] and the fruit is used as an insect repellent [1]. *M. aurantiaca* had been recorded to use for its use in the treatment of cardiovascular ailments [4]. Many types of compounds have been isolated from this genus including xanthenes, triterpenes, stilbenes and flavonoids [2-11]. The latter compounds were found as major phytochemicals in this genus and many of these compounds had broad spectrum biological activities such as anti-inflammatory [4] and anti-HIV activities [7], cholinesterase [5] and PDE5 inhibitory activities [8], and their antimicrobial [9], cytotoxic [10], and antioxidant [2] activities. In the course of our ongoing search for bioactive compounds from natural resources from the northern part of Thailand [12-16], we report here the first phytochemical investigation of the twig extracts of *M. fruticosa* (Roxb.) Corner (Figure 1) collected from Doi Tung, Chiang Rai Province, Thailand resulting in the isolation and identification of a new xanthone (1) together with 14 known compounds (2-15) (Figure 2). In addition, anti-oxidant, α -glucosidase inhibitory and cytotoxic activities of some of the isolated compounds were evaluated. To the best of our knowledge, this is the first report of the phytochemical investigation of this plant.



Figure 1: *Maclura fruticosa* (these photos were taken by Sarot Cheenpracha).




Figure 1: Chemical structures of isolated compounds from *M. fruticosa*. The crude extraction of *M. fruticosa* (twigs) were separated by repeated silica gel column chromatography and Sephadex LH-20

column chromatography to obtain a new xanthone (**1**) together with 14 known compounds (**2-15**) were identified as ananixanthone (**2**) [17], 8-desoxygatanin (**3**) [18], parvifolixanthone A (**4**) [19], 1,3,7-trihydroxy-2,8-di-(3-methylbut-2-enyl)xanthone (**5**) [20], xanthone V₁ (**6**) [21], xanthone V_{1a} (**7**) [21], 1,3,5-trihydroxy-2-prenylxanthone (**8**) [22], scandenone (**9**) [23], alpinumisflavone (**10**) [24], cudraiflavone G (**11**) [25], wightone (**12**) [26], erpoeigin C (**13**) [27], genistein (**14**) [28], and artonin Y (**15**) [29], from their NMR spectroscopic data as well as comparisons made with literature data.

Table 1: NMR Spectroscopic Data of Maclurafruticosone (**1**) in CDCl₃.

Position	δ_C	δ_H [J in Hz]	HMBC
1	156.1		
2	104.6		
3	158.2		
4	107.7		
4a	154.3		
5	133.5		
5a	144.6		
6	151.1		
7	107.5	6.96 d (8.9)	C-5, C-6, C-8a
8	117.1	7.79 d (8.9)	C-5a, C-6, C-9
8a	115.3		
9	181.0		
9a	103.1		
1'	115.9	6.75 d (10.0)	C-3, C-3"
2'	127.3	5.60 d (10.0)	C-2, C-3"
3'	78.2		
4'	28.5	1.48 s	C-2', C-3', C-5'
5'	28.5	1.48 s	C-2', C-3', C-4'
1"	21.7	3.53 d (7.3)	C-3, C-4, C-4a, C-2', C-3'
2"	122.5	5.30 m	C-5"
3"	131.7		
4"	25.9	1.69 d (1.0)	C-2', C-3', C-5'
5"	18.0	1.87 d (0.8)	C-2', C-3', C-4'
OH-1	-	13.20 s	C-1, C-2, C-9a
OCH ₃ -6	56.7	4.04 s	C-6

Maclurafruticosone (**1**), a yellow solid, had a 1,3,5,6-tetraoxygenated xanthone core structure which showed a [M-H]⁻ ion at *m/z* 407.1480 (calcd, 407.1495) in HRESIMS, indicating a molecular formula of C₂₄H₂₄O₆. The UV spectrum showed the maxima absorptions at λ_{max} 286 and 337 nm while the IR spectrum showed absorption bands at 3198 cm⁻¹ for hydroxy group and at 1652 cm⁻¹ for a conjugated carbonyl group. The latter functional group was also supported by the ¹³C-NMR resonance at δ_C 181.0 (C-9). This carbonyl group formed an H-bond with the hydroxy group at C-1 as evidenced by the resonance for an H-bonded hydroxy proton at δ_H 13.20 (1H, s, 1-OH) in the ¹H NMR spectrum (Table 1). The HMBC correlations of this H-bonded hydroxy proton (δ_H 13.20) with δ_C 156.1 (C-1), 104.6 (C-2) and 103.1 (C-9a) supported the structure of **1**. The *ortho*-coupled aromatic protons at δ_H 6.96 (1H, d, *J* = 8.9 Hz) and δ_H 7.79 (1H, d, *J* = 8.9 Hz) were assigned to protons H-7 and H-8, respectively, by the ²*J* and ³*J* HMBC correlations of H-7 (δ_H 6.96) to δ_C 133.5 (C-5), 151.1 (C-6), and 115.3 (C-8a); and H-8 (δ_H 7.79) to δ_C 144.6 (C-5a), 151.1 (C-6), and 181.0 (C-9). The ¹H and ¹³C NMR spectra of **1** also displayed resonances for a 2,2-dimethylpyrano unit at δ_H 6.75 (1H, d, *J* = 10.0 Hz, H-1')/ δ_C 115.9 (C-1'), δ_H 5.60 (1H, d, *J* = 10.0 Hz, H-2')/ δ_C 127.3 (C-2'), δ_H 1.48 (6H, s, H-4' and H-5')/ δ_C 28.5 (C-4' and C-5'), and δ_C 78.2 (C-3') and an prenyl unit at δ_H 5.30 (1H, m, H-2'')/ δ_C 122.5 (C-2''), δ_H 3.53 (2H, d, *J* = 7.3 Hz, H-1'')/ δ_C 21.7 (C-1''), δ_H 1.69 (3H, s, H-4'')/ δ_C 25.9 (C-4''), 1.87 (3H, s, H-5'')/ δ_C 18.0 (C-5''), and δ_C 131.7 (C-3''). The latter moiety was located at C-4 supported by the ²*J* and ³*J* HMBC correlations of H-1'' (δ_H 3.53) with C-3 (δ_C 158.2), C-4 (δ_C 107.7), and C-4a (δ_C 154.3); and H-1' (δ_H 6.75) with C-3 (δ_C 158.2). The 2,2-dimethylpyrano moiety was placed at C-2/C-3 supported with the following HMBC correlations: H-1' (δ_H 6.75) with C-3 (δ_C 158.2); H-2' (δ_H 5.60) with C-2 (δ_C 104.6); and OH-1 (δ_H 13.20) with C-2 (δ_C 104.6). The remaining methoxy group with a resonance at δ_H 4.04 (3H, s) was located at C-6 from the ³*J* HMBC correlations of its methoxy protons (δ_H 4.04) and H-8 (δ_H 7.79) with C-6 (δ_C 151.1) as well as the cross peak between H-7

(δ_H 6.96) and the methoxy protons (δ_H 4.04) in the NOESY experiment. Thus, maclurafruticosone was assigned the structure **1**.

Most of isolated compounds were evaluated for their antioxidant activities against DPPH radical and ABTS^{•+} (Table 2). The EtOAc extract of the twigs of *M. fruticososa* showed significant antioxidant activities against DPPH radical and ABTS^{•+} with high % inhibition of 95.14 % (at 500 μ g/mL) and 99.36 % (at 200 μ g/mL), respectively. Compounds **4**, **6**, and **7** showed significant antioxidant activity against the DPPH radical with IC₅₀ values ranging from 7.45–16.12 μ M, with compound **6** having the highest activity. Only compound **4** exhibited potent scavenging activity against ABTS^{•+} with an IC₅₀ value of 0.55 \pm 0.01 μ M, which was better than the positive control ascorbic acid (IC₅₀ 2.35 \pm 0.17 μ M). It is interesting to note that all tested xanthones were found to have antioxidant activities against both of DPPH radical and ABTS^{•+} better than those of isoflavones.

The EtOAc extract of the twig of *M. fruticososa* showed moderate α -glucosidase inhibitory activity with 73.97 % inhibition at 200 μ g/mL. Compounds **3**, **5**, **6**, **9-12**, **14** and **15** were evaluated for their α -glucosidase inhibitory activities. Only compound **12** exhibited significant α -glucosidase inhibitory activity with an IC₅₀ value of 0.02 \pm 0.37 mM which was better than acarbose (IC₅₀ 1.55 \pm 0.39 mM). Compounds **5**, **6**, **12** and **14** were evaluated for their cytotoxicity against a colon cancer cell line and all of them, except compound **14**, were found to have weak cytotoxicity, with IC₅₀ values ranging from 22.35–47.62 μ M (positive control, doxorubicin, IC₅₀ 9.74 μ M). Compound **14** was inactive.

Table 2: Antioxidant Activities of Compounds **3-15**.

Compounds	DPPH assay		ABTS assay	
	%Inhibition (500 μ g/mL)	IC ₅₀ (μ M)	%Inhibition (200 μ g/mL)	IC ₅₀ (μ M)
Crude extract	95.14 \pm 0.52	–	99.36 \pm 1.01	–
3	84.19 \pm 1.26	181.74 \pm 4.78	81.72 \pm 1.48	53.12 \pm 3.00
4	97.47 \pm 0.32	16.12 \pm 0.70	96.37 \pm 0.31	0.55 \pm 0.01
5	80.24 \pm 2.37	208.21 \pm 5.07	28.66 \pm 0.95	–
6	95.89 \pm 0.63	7.45 \pm 0.32	30.06 \pm 0.81	–
7	88.46 \pm 0.79	10.43 \pm 0.44	93.69 \pm 1.22	not tested*
8	32.67 \pm 1.26	–	64.46 \pm 1.80	–
9	24.13 \pm 0.32	–	54.08 \pm 5.00	–
10	25.40 \pm 0.95	–	84.40 \pm 0.50	116.92 \pm 2.73
11	44.52 \pm 3.00	–	94.71 \pm 0.50	64.05 \pm 5.62
12	30.61 \pm 0.16	–	89.30 \pm 3.53	79.61 \pm 4.79
13	38.04 \pm 1.26	–	67.07 \pm 1.89	–
14	50.68 \pm 3.48	–	95.92 \pm 0.95	168.18 \pm 0.30
15	86.72 \pm 1.26	423.06 \pm 8.70	97.32 \pm 0.87	not tested*
Ascorbic acid	–	2.01 \pm 0.22	–	2.35 \pm 0.17

*Inadequate sample available. All determinations were carried out in triplicate, and results were expressed as mean \pm standard deviation (SD).

In summary, we have isolated eight xanthones and seven flavonoids from the twig extract of *M. fruticososa*. These results are in agreement of previous research on the *Maclura* species [2-10]. The difference between *M. fruticososa* and the other species of *Maclura* studied is that both xanthones and flavonoids were major compounds of *M. fruticososa* while in the other species of *Maclura* studied only flavonoids or xanthones were isolated as major compounds. Therefore, the flavonoids and xanthones from this plant might be a useful as a chemotaxonomic marker to differentiate *M. fruticososa* from other species of *Maclura*. In the present work, compounds **4** and **12** exhibited potent antioxidant activity against ABTS^{•+} and α -glucosidase inhibitory activity, respectively, showing better activity than those of the standard controls. This preliminary result indicated that compound **12** might be a good candidate for further study and development as an anti-diabetic agent and compound **4** might be a potential lead compound for antioxidant agent development.

Experimental

General Experimental Procedures: Melting point was determined using a Gallenkamp melting point apparatus. UV-vis spectra were recorded with BMG LABTECH/SPECTROstar Nano spectrometer or Agilent 8453 UV-visible spectrophotometer. The IR spectra were recorded using a PerkinElmer FTS FT-IR spectrometer or SHIMADSU spectrophotometer. The NMR spectra were recorded using a 400 MHz Bruker spectrometer or a 400 MHz Bruker AM400 spectrometer. HRESIMS mass spectra carried out on a Bruker micro TOF mass spectrometer. Unless otherwise indicated, all quick column chromatography (QCC) and column chromatography (CC) were carried out on silica gel 60 H (5-40 μm , SiliCycle[®] Inc.) and silica gel 100 (63-200 μm , SiliCycle[®] Inc.), respectively. Sephadex LH-20, when indicated, was also used for CC. Precoated TLC plates of silica gel 60 F₂₅₄ were used for analytical purposes.

Plant Material: The twigs of *M. fruticosa* were collected from Doi Tung, Chiang Rai Province, Thailand in August 2015. The plant was identified by Mr. Martin Van de Bult, and the voucher specimen (MFU-NPR0111) was deposited at the Natural Products Research Laboratory of Mae Fah Luang University.

Extraction and Isolation: Air-dried twigs of *M. fruticosa* (5.4 kg) were macerated in EtOAc (3 \times 10 L) at room temperature. The extract was filtered and concentrated under reduced pressure to give the crude EtOAc extract (108.3 g). This extract was subjected to QCC over silica gel, eluting with a gradient system of acetone/hexanes (100% hexanes to 100% acetone) to give 16 fractions (1A-1P). Fraction 1J (3.4 g) was further separated by QCC (100% hexanes to 100% acetone) to give six subfractions (2JA-2JF). Compound **1** (1.2 mg) was obtained from subfraction 2JB (126.0 mg) by CC over Sephadex LH-20 (100% MeOH) and then by CC (4:1 CH₂Cl₂/hexanes). Subfraction 3JB (20.1 mg) was purified by CC (1:49 acetone/hexanes) to yield compound **2** (2.2 mg). Compound **9** (25.2 mg) was obtained from subfraction 2JC (820.4 mg) by CC over Sephadex LH-20 (100% MeOH) and then by CC (1:4 EtOAc/hexanes). Fraction 1K (4.6 g) was further separated by QCC (100% hexanes to 100% acetone) to give eight subfractions (2KA-2KH). Compounds **3** (11.1 mg) and **4** (11.3 mg) were obtained from subfraction 2KC (319.1 mg) by repeated CC (1:9 CH₂Cl₂/hexanes). Purification of subfraction 4KC (90.8 mg) by CC over Sephadex LH-20 (100% MeOH) and then by CC (1:49 EtOAc/CH₂Cl₂) afforded compound **10** (16.3 mg). Subfraction 4KD (31.9 mg) was purified by CC over Sephadex LH-20 (100% MeOH) and then by CC (1:49 acetone/hexanes) to yield compound **11** (6.9 mg). Compound **5** (37.3 mg) was obtained from subfraction 2KE (1.2 mg) by CC (100% CH₂Cl₂), followed by repeated CC (1:49 EtOAc/CH₂Cl₂). Separation of subfraction 3KC (944.3 mg) by repeated CC (1:49 acetone/CH₂Cl₂) afforded compound **6** (51.8 mg). Fraction 1L (4.8 g) was further separated by CC over Sephadex LH-20 (100% MeOH) to give six subfractions (2LA-2LF). Compound **7** (4.3 mg) was obtained from subfraction 2LC (360.9 mg) by CC (1:49 MeOH/CH₂Cl₂) and then by repeated CC (1:49 acetone/hexanes). Separation of subfraction 2LD (728.1 mg) by CC (1:49 MeOH/CH₂Cl₂) and further CC (1:24 EtOAc/CH₂Cl₂) yielded compound **8** (4.7 mg). Fraction 1M (6.6 g) was further separated by CC (100% hexanes to 100% acetone) to give nine subfractions (2MA-2MI). Compound **12** (204.5 mg) was obtained from subfraction 2MC (1.2 g) by CC (1:4 EtOAc/hexanes) and further CC (1:49 acetone/CH₂Cl₂). Subfraction 2MD (1.2 g) was subjected to CC over Sephadex LH-20 (100% MeOH) and then by CC (3:7 EtOAc/CH₂Cl₂) to afford compound **13** (8.7 mg). Separation of subfraction 3ME (189.2 mg) by CC over Sephadex LH-20 (100% MeOH) and then by CC (3:7 EtOAc/hexanes) gave compound **14** (21.1 mg). Compound **15** (12.4 mg) was obtained

from subfraction 2MF (332.7 mg) by CC over Sephadex LH-20 (100% MeOH) and further CC (1:49 MeOH/CH₂Cl₂). The isolation flowchart of compounds **1-15** is shown in Figure S8.

Maclurafruticosone (1)

Yellow solid.

MP: 206°C (decomposed).

IR (neat) V_{max} : 3198, 2920, 1652, 1576, 1436, 1286, 798 cm^{-1} .

UV (MeOH) λ_{max} (log ϵ): 286.0 (4.55), 337.0 (4.39) nm.

¹H NMR (CDCl₃, 400 MHz) and ¹³C NMR (CDCl₃, 100 MHz): Table 1.

HRESIMS m/z 407.1480 [M-H]⁻ (calcd for C₂₄H₂₃O₆, 407.1495).

DPPH Radical Scavenging Assay: The DPPH scavenging activity assay was modified from a previous paper [14]. Briefly, 6 \times 10⁻⁵ M DPPH was prepared in absolute EtOH, and then 100 μL of this solution was mixed with 100 μL of sample solution in EtOH in a 96-well microplate. After 30 min of reaction, in the absence of light at room temperature, the absorbance was measured using a microplate reader (SPECTROstar Nano) at 517 nm. The DPPH radical scavenging capacity was calculated using the following equation. % inhibition = [(A_B - A_S) / A_B] \times 100, where A_B and A_S are the absorbance of the blank sample and sample, respectively. All experiments were performed in triplicate, with ascorbic acid used as the positive control ($r^2 = 0.9829$, IC₅₀ 2.01 \pm 0.22 μM). The half maximal inhibitory concentration (IC₅₀) of DPPH scavenging activity was calculated by plotting inhibition percentages against concentrations of the sample. The inhibition values are reported as means \pm SD.

ABTS^{•+} Scavenging Assay: The determination of ABTS^{•+} scavenging activity was carried out using a modified literature procedure [30]. ABTS^{•+} were produced by reacting a 7 mM stock solution of ABTS in DI water with 2.45 mM potassium persulfate (K₂S₂O₈) and allowing the mixture to stand in the dark at room temperature for 16 h before use. The ABTS^{•+} solution was diluted with water to an absorbance of 0.7 \pm 0.02 at 734 nm. Different concentrations of test samples in EtOH (100 μL) and ABTS^{•+} solution (150 μL) were added to each well of the 96-well microplate. The absorbance at 750 nm was determined after 5 min of mixing. The percentage of ABTS free radical scavenging activity was calculated using the same formula as for the DPPH assay. The measurements were performed in triplicate and the positive control was ascorbic acid ($r^2 = 0.9777$, IC₅₀ 2.35 \pm 0.17 μM).

α -Glucosidase Inhibitory Assay: Briefly, the reaction mixture contained 50 μL of sample in DMSO was pre-incubated with α -glucosidase enzyme (0.35 U/mL, 100 μL) at 37°C for 10 min. The reaction was started when 100 μL of *p*-nitrophenyl glucopyranoside (*p*-NPG) substrate was added in phosphate buffer. The reaction was incubated at 37°C for 20 min and stopped by the addition of 1 mL 1 M of Na₂CO₃ solution. The α -glucosidase activity was determined by measuring the amount of *p*-nitrophenol released from *p*-NPG at 405 nm using a microplate reader (SPECTROstar Nano) [16]. Blanks were prepared by adding solvent instead of samples. Acarbose was used as a positive control ($r^2 = 0.9897$, IC₅₀ 1.55 \pm 0.39 mM). All assays were carried out in triplicate.

Cytotoxicity Assay: Colon cancer cells (HCT116) (1 \times 10⁴ cells/well) were cultured in 96 well plate and allowed to adhere for 24 h at 37 °C. The cells were treated with compounds (10 μM or $\mu\text{g/mL}$) in DMEM medium for 24 h. Then the medium were removed and fresh of DMEM containing 0.5 mg/mL of MTT solution were added to each well for 2 h. After that, the medium were discarded by aspirator. The violet formazan crystals in the viable cells were dissolved in 100 μL of DMSO. The absorbance of

each well was then read at a wavelength of 570 nm using a microplate reader. Doxorubicin was used as a positive control with an IC₅₀ value of 9.74 μ M.

% Cell viability is expressed as: $\frac{\text{Absorbance of treated well}}{\text{Absorbance of control well}} \times 100$

% Cytotoxicity = 100 – % cell viability

Supplementary data: ¹H NMR, ¹³C NMR, ¹H-¹H COSY, HSQC, HMBC and HRESIMS of **1**.

Acknowledgments – This work was supported by the Thailand Research Fund through the Direct Basic Research Grant (DBG5980001) and Mae Fah Luang University through the graduate student research grant. Mae Fah Luang University and the University of Wollongong were also acknowledged for laboratory facilities. We would like to thank Mr. Matin Van de Bult, Doi Tung Development Project, Chiang Rai, Thailand for plant collection and identification. S. Laphookhieo thanks the Australian Government via the Endeavour Award 2016 for a research fellowship.

References

- [1] Harry-O'kuru RE, Tisserat B, Gordon SH, Gravett A. (2015) Osage orange (*Maclura pomifera* L.) seed oil poly(α -hydroxydibutylamine) triglycerides: Synthesis and characterization. *Journal of Agricultural and Food Chemistry*, **63**, 6588–6595.
- [2] Tsao R, Yang R, Young JC. (2003) Antioxidant isoflavones in osage orange, *Maclura pomifera* (Raf.) Schneid. *Journal of Agricultural and Food Chemistry*, **51**, 6445–6451.
- [3] Cioffi G, Escobar LM, Braca A, Tommasi ND. (2003) Antioxidant chalcone glycosides and flavanones from *Maclura* (*Chlorophora*). *Journal of Natural Products*, **66**, 1061–1064.
- [4] Kupeli E, Orhan I, Tokar G, Yesilada E. (2006) Anti-inflammatory and antinociceptive potential of *Maclura pomifera* (Rafin.). Schneider fruit extracts and its major constituents, scandenone and auriculasin. *Journal of Ethnopharmacology*, **107**, 169–174.
- [5] Orhan I, Senol FS, Martal M, Dvorska M, Zemlicka M, Smejkal K, Mokry P. (2009) Cholinesterase inhibitory effects of the extracts and compounds of *Maclura pomifera* (Rafin.) Schneider. *Food Chemical Toxicology*, **47**, 1747–1751.
- [6] Lee SJ, Wood AR, Maier CGA, Dixon RA, Mabry TJ. (1994) Prenylated flavonoids from *Maclura pomifera*. *Phytochemistry*, **49**, 2573–2577.
- [7] Groweiss A, Cardellina JH, Boyd MR. (2000) HIV-inhibitory prenylated xanthenes and flavones from *Maclura tinctoria*. *Journal of Natural Products*, **63**, 1537–1539.
- [8] Ribaudo G, Vendrame T, Bova S. (2016) Isoflavones from *Maclura pomifera*: structural elucidation and *in silico* evaluation of their interaction with PDE5. *Natural Products Research*, doi: 10.1080/14786419.2016.1269101.
- [9] Mahmoud Z. (1981) Antimicrobial component from *Maclura pomifera* fruit. *Planta Medica*, **42**, 299–301.
- [10] Son H, Chung M, Lee SI, Yang HD, Moon H. (2007) Pomiferin, histone deacetylase inhibitor isolated from the fruits of *Maclura pomifera*. *Bioorganic Medicinal Chemistry Letters*, **17**, 4753–4755.
- [11] El-Sohly HN, Joshia A, Lia XC, Rossa SA. (1999) Flavonoids from *Maclura tinctoria*. *Phytochemistry*, **52**, 141–145.
- [12] Maneerat W, Ritthiwigrom T, Cheenpracha S, Promgool T, Yossathera K, Deachathai S, Phakhodee W, Laphookhieo S. (2012) Bioactive carbazole alkaloids from *Clausena wallichii* roots. *Journal of Natural Products*, **75**, 741–746.
- [13] Meesakul P, Pudhom K, Pyne SG, Laphookhieo S. (2017) Hybrid flavan–flavanones from *friesodielsia desmoides* and their inhibitory activities against nitric oxide production. *RSC Advances*, **7**, 17545–17550.
- [14] Tantapakul C, Phakhodee W, Ritthiwigrom T, Cheenpracha S, Prawat U, Deachathai S, Laphookhieo S. (2012) Rearranged benzophenones and prenylated xanthenes from *Garcinia propinqua* twigs. *Journal of Natural Products*, **75**, 1660–1664.
- [15] Tantapakul C, Maneerat W, Sripisut T, Ritthiwigrom T, Andersen RJ, Cheng P, Cheenpracha S, Raksat A, Laphookhieo S. (2016) New benzophenones and xanthenes from *Cratoxylum sumatranum* ssp. *Nerifolium* and their antibacterial and antioxidation activities. *Journal of Agricultural and Food Chemistry*, **64**, 8755–8762.
- [16] Sriyatep T, Siridechakorn I, Maneerat W, Pansanit A, Ritthiwigrom T, Andersen RJ, Laphookhieo S. (2015) Bioactive prenylated xanthenes from the young fruits and flowers of *Garcinia cowa*. *Journal of Natural Products*, **78**, 265–271.
- [17] Daud SD, Ee GC, Malek EA, Ahmad Z, Hashim NM, See I.; Teh, SS, Ismail AF. (2016) A new pyranoxanthone from the stem bark of *Calophyllum buxifolium*. *Chemistry of Natural Compounds*, **52**, 807–809.
- [18] Helesbeux JJ, Duval O, Dartiguelongue C, Séraphin D, Oger JM, Richomme P. (2004) Synthesis of 2-hydroxy-3-methylbut-3-enyl substituted coumarins and xanthenes as natural products. Application of the schenck ene reaction of singlet oxygen with ortho-prenylphenol precursors. *Tetrahedron*, **60**, 2293–2300.
- [19] Rukachaisirikul V, Naklue W, Phongpaichit S, Towatana, NH, Maneenoon K. (2006) Phloroglucinols, depsidones and xanthenes from the twigs of *Garcinia parvifolia*. *Tetrahedron*, **62**, 8578–8585.
- [20] Mahabusarakum W, Wiriyachitra P. (1987) Chemical constituents of *Garcinia mangostana*. *Journal of Natural Products*, **50**, 474–478.
- [21] Botta B, Monache GD, Monache FD, Marini GB, Meniching F. (1986) Vismione H and prenylated xanthone from *Vismia guineensis*. *Phytochemistry*, **25**, 1217–1219.
- [22] Wang R, Chen R, Li J, Liu X, Xie K, Chen D, Peng Y, Dai J. (2016) Regiospecific prenylation of hydroxy xanthenes by a plant flavonoid prenyltransferase. *Journal of Natural Products*, **79**, 2143–2147.
- [23] Talla E, Njamen D, Mbafor JT, Fomum ZT, Kamanyi A, Mbanya JC, Giner RM, Recio MC, Máñez S, Ríos JL. (2003) Warangalone, the isoflavonoid anti-inflammatory principle of *Erythrina addisoniae* stem bark. *Journal of Natural Products*, **66**, 891–893.
- [24] Han XH, Hong SS, Hwang JH, Jeong SH, Hwang HJ, Lee MH, Lee MK, Lee D, Ro JH, Hwang BY. (2005) Monoamine oxidase inhibitory constituents from the fruits of *Cudrania tricuspidata*. *Archives of Pharmacal Research*, **28**, 1324–1327.
- [25] Hiep NT, Kwon J, Kim DW, Hwang BY, Lee HJ, Mar W, Lee D. (2015) Isoflavones with neuroprotective activities from fruits of *Cudrania tricuspidata*. *Phytochemistry*, **111**, 141–148.
- [26] Tahara S, Nakahara S, Ingham JL, Mizutani J. (1985) Fungal metabolites of antifungal isoflavone wighteone. *Nippon Nogeik Kaishi*, **59**, 1039–1044.
- [27] Tahara S, Ingham JL, Mizutani J. (1985) New coumaronochromones from white lupin, *Lupinus albus* L. (Leguminosae). *Agricultural Biological Chemistry*, **49**, 1775–1783.
- [28] Yasuda T, Mizunuma S, Kano Y, Saito K, Ohsawa K. (1996) Urinary and biliary metabolites of genistein in rats. *Biological and Pharmaceutical Bulletin*, **19**, 413–417.
- [29] Shinomiya K, Hano Y, Nomura T. (2000) Mechanism on one-sided wessely-moser rearrangement reaction. *Heterocycles*, **53**, 877–886.
- [30] Zheng L, Zhao M, Xiao C, Zhao Q, Su G. (2016) Practical problems when using ABTS assay to assess the radical-scavenging activity of peptides: importance of controlling reaction pH and time. *Food Chemistry*, **192**, 288–294.