



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Surgery

Department of Surgery

November 2015

K Wire fixation of supra-condylar humerus fractures in children. Is ulnar nerve at risk?

Waseem Ahmed

Aga Khan University, waseem.ahmed@aku.edu

Haroon Rashid

Aga Khan University, haroon.rashid@aku.edu

Yasir Mohib

Aga Khan University, yasir.mohib@aku.edu

Rizwan Haroon Rashid

Aga Khan University, rizwan.haroonrashid@aku.edu

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_surg_surg

 Part of the [Surgery Commons](#)

Recommended Citation

Ahmed, W., Rashid, H., Mohib, Y., Rashid, R. (2015). K Wire fixation of supra-condylar humerus fractures in children. Is ulnar nerve at risk?. *JPMA: Journal of Pakistan Medical Association*, 65(11), S-202-S-204.

Available at: http://ecommons.aku.edu/pakistan_fhs_mc_surg_surg/146

K Wire fixation of supra-condylar humerus fractures in children. Is ulnar nerve at risk?

Waseem Ahmed, Haroon Rasheed, Yasir Mohib, Rizwan Haroon Rashid

Abstract

Objective: To compare iatrogenic ulnar nerve injury in lateral entry pin fixation versus medial and lateral entry pin fixation in the treatment of supracondylar fractures of the humerus in children. Measurement of clinical parameters in terms of elbow range of motion and postop radiographic alignment was also targeted.

Methods: The retrospective cohort study was conducted at Aga Khan University Hospital, Karachi, and comprised data of paediatric patients who underwent closed reduction and percutaneous pin fixation for the treatment of displaced extension type supracondylar fractures of the humerus between July 2007 and June 2012. Data regarding socio-demographic status, disease and procedure variables was collected from patient files and was analysed using SPSS 19.

Results: There were 71 patients; 37(52%) in the lateral entry group and 34(48%) in the medial and lateral entry group. The two groups were similar in terms of mean age, gender distribution, and preoperative displacement, comminution, and associated vascular and nerve status ($p>0.05$ each). There were no cases of iatrogenic ulnar nerve injury in either group and no significant differences between groups with respect to the elbow range of motion and radiological parameters ($p>0.05$ each).

Conclusions: With the use of the specific techniques employed, both lateral entry pin fixation and medial and lateral entry pin fixation were effective in the treatment of displaced extension type supracondylar humerus fractures in children.

Keywords: Humerus fracture, Supracondylar fracture, K wire fixation, CRPP. (JPMA 65: S-202 (Suppl. 3); 2015)

Introduction

Supracondylar fractures of the humerus are the most common type of elbow fracture in children. They account for 50% to 70% of all elbow fractures and are seen most frequently in children aged 6-9 years.¹⁻¹⁰ Extension type is more common, accounting for about 95% to 98% of all supracondylar fractures.^{10,11} Due to the close proximity of delicate nerves and vessels around the condyles, these structures can be damaged either by impact of trauma or at risk during fracture reduction and fixation.¹² The incidence of traumatic and iatrogenic nerve injuries with this type of fracture have been recorded to be 12% to 20%.^{10,13} Reduction and percutaneous pin fixation is considered the standard method of management.¹⁴ The configuration of pins remains a source of debate, with crossed medial-lateral pin fixation providing more stability compared to two lateral pins but having the risk of iatrogenic ulnar nerve injury.¹⁵

Supracondylar humerus fractures are a common problem affecting the community and maintenance of anatomical reduction by smooth pins is crucial in the outcome.¹⁶ Iatrogenic ulnar nerve injury still forces the operating

surgeon to use 2 lateral pins, compromising the maximal stability provided by the construct. The safety of using cross-pin fixation in terms of iatrogenic ulnar nerve injury has been mentioned in literature.^{4,12,17,18}

The current study was planned to compare iatrogenic ulnar nerve injury in lateral entry pin fixation versus medial and lateral entry pin fixation in the treatment of supracondylar fractures of the humerus in children. Measurement of clinical parameters in terms of elbow range of motion and postop radiographic alignment was also targeted.

Materials and Methods

The retrospective cohort study was conducted at Aga Khan University Hospital, Karachi, and comprised data of paediatric patients who underwent closed reduction and percutaneous pin fixation for the treatment of displaced extension type supracondylar fractures of the humerus between July 2007 and June 2012. Patients were identified from the Health Information Management System (HIMS), All patients between age 2 and 15 years, extension type supracondylar humerus fracture who underwent closed reduction and K wire fixation were included. All patients were operated by or under direct supervision of the consultant orthopaedic surgeon. Decision regarding the lateral pin fixation or medial and

Department of Surgery, Aga Khan University Hospital, Karachi.

Correspondence: Waseem Ahmed. Email: dr.waseemzubair@gmail.com

lateral pin fixation was governed by the choice of the operating surgeon. Patients with pre-operative ulnar nerve injury and age <2 years or >15 years were excluded.

Data regarding age, gender, fixation method, iatrogenic ulnar nerve injury, range of motion (ROM) at 6 weeks post-operative was collected by an orthopaedic Resident. All pre operative and 6-week post operative radiographs were assessed and data regarding alignment, Bauman angle and carrying angle were calculated. SPSS 19 was used for data analysis and $p < 0.05$ was taken as significant.

Results

A total of 105 files were identified of which 34(32%) did not meet the inclusion criteria and had to be discarded. The final study sample stood at 71. Among them, 37(52%) underwent lateral pin fixation; 17(46%) males and 20(54%) females with overall mean age of 5.67 ± 3.4 years (range: 2-15 years). Likewise, 34(48%) patients had medial and lateral cross-pin fixation; 16(47%) males and 18(53%) females with a mean age of 6.4 ± 2.6 years (range: 2-15 years) (Table-1).

There were no cases of iatrogenic ulnar nerve injury in

Table-1: Demographic Data.

Patients characteristics N=71	Medial-lateral pin entry group N=34	Lateral pin entry group N=37	p-value (significant if $p < 0.05$)
Mean Age (years)	6.4 ± 2.6	5.67 ± 3.4	0.11
Gender			0.83
Male	16	17	
Female	18	20	

Table-2: Iatrogenic ulnar nerve injury.

Complications	Medial-lateral pin entry group N=34	Lateral pin entry group N=37	p-value (significant if $p < 0.05$)
Post-operative Iatrogenic ulnar nerve injury	0	0	1.0

Table-3: Outcome at follow-up.

Assessment measures	Medial-lateral pin entry group N=34	Lateral pin entry group N=37	p-value (significant if $p < 0.05$)
Elbow (Degrees)			
Total Range of motion	$0-131 \pm 4.2$	$0-129 \pm 3.8$	0.50
Carrying angle	6.1 ± 1.5	6.6 ± 1.3	0.31
Baumann angle	80.0 ± 3.2	79.3 ± 3.0	0.73

either group (Table-2).

At 6-week follow-up, there were no significant differences between the groups regarding carrying angle, total elbow motion, Baumann angle and change in Baumann angle and loss of reduction ($p > 0.05$ each) (Table-3).

Discussion

Results of both cross-pin fixation and lateral pin fixation were comparable in terms of iatrogenic ulnar nerve injury, elbow ROM and radiological parameters.⁷ Fortunately, no iatrogenic ulnar nerve injury was encountered in any of our patients, emphasising the safe use of cross-pin construct.

Literature suggests that crossed medial-lateral pin fixation provides increased biomechanical stability, but there is a risk of iatrogenic ulnar nerve injury from placement of the medial pin (14) whereas the two lateral pin fixation avoids the danger of iatrogenic ulnar nerve injury, but it provides less biomechanical stability.^{14,19}

However, a study¹¹ has suggested that it is difficult to compare various studies because pinning technique, pin size and position of elbow during pinning differs and most of the studies have a sample size of less than 50.²⁰

A systematic review in 2010 suggested that the risk of iatrogenic ulnar nerve injury is more with the crossed pinning compared with the lateral pinning.¹⁶ However, paucity of local literature still exists about the issue and many surgeons are found to be apprehensive in placing the medial entry pin to obtain a stable cross-pin construct.¹⁵ Although there is a potential risk of iatrogenic ulnar nerve injury, but with vigilant technique it can be minimised to such an extent that it is to be considered a desirable option in all unstable situations.¹⁶

Conclusion

Both percutaneous fixation techniques are effective in terms of efficacy and safety if a uniform and standardised operative technique is followed in each method.

Reference

1. Mitchelson AJ, Illingworth KD, Robinson BS, Elnimeiry KA, Wilson CJ, Markwell SJ, et al. Patient demographics and risk factors in pediatric distal humeral supracondylar fractures. *Orthopedics* 2013; 36: e700-6. doi: 10.3928/01477447-20130523-12.
2. Abzug JM, Herman MJ. Management of supracondylar humerus fractures in children: current concepts. *J Am Acad Orthop Surg* 2012; 20: 69-77.
3. Balakumar B, Madhuri V. A retrospective analysis of loss of reduction in operated supracondylar humerus fractures. *Indian J Orthop* 2012; 46: 690-7.
4. Howard A, Mulpuri K, Abel MF, Braun S, Bueche M, Epps H, et al. The treatment of pediatric supracondylar humerus fractures. *J Am Acad Orthop Surg* 2012; 20: 320-7.

5. Leaman C, Kotwal R, Williams P. A simple tip to improve the accuracy of crossed K-wire placement in the management of displaced paediatric supracondylar fractures of the humerus. *Ann R Coll Surg Engl* 2012; 94: 602.
 6. Loder RT, Krodel E, D'Amico K. Temporal variation in pediatric supracondylar humerus fractures requiring surgical intervention. *J Child Orthop* 2012; 6: 419-25.
 7. Kish AJ, Hennrikus WL. Fixation of type 2a supracondylar humerus fractures in children with a single pin. *J Pediatr Orthop* 2014; 34:e54-7. doi: 10.1097/BPO.0000000000000247.
 8. Hussain S, Dhar S, Qayoom A. Open reduction and internal fixation of displaced supracondylar fractures of humerus with crossed K-wires via medial approach. *Malays Orthop J* 2014; 8: 29-34.
 9. Pennock AT, Charles M, Moor M, Bastrom TP, Newton PO. Potential causes of loss of reduction in supracondylar humerus fractures. *J Pediatr Orthop* 2014; 34: 691-7.
 10. Alton TB, Werner SE, Gee AO. Classifications in brief: the Gartland classification of supracondylar humerus fractures. *Clin Orthop Relat Res* 2015; 473: 738-41.
 11. Pullagura M, Odak S, Pratt R. Managing supracondylar fractures of the distal humerus in children in a district general hospital. *Ann R Coll Surg Engl* 2013; 95: 582-5.
 12. Khademolhosseini M, Abd Rashid AH, Ibrahim S. Nerve injuries in supracondylar fractures of the humerus in children: is nerve exploration indicated? *J Pediatr Orthop B* 2013; 22: 123-6.
 13. Sahu RL. Percutaneous K-wire fixation in paediatric supracondylar fractures of humerus: a retrospective study. *Niger Med J* 2013; 54: 329-34.
 14. Pedrazzini A, Verdano MA, De Caro F, Pellegrini A, Ceccarelli F. Are percutaneous pinning the best treatment for Gartland type iii supracondylar humeral fractures in children? *Acta Biomed* 2013; 84: 110-6.
 15. Kocher MS, Kasser JR, Waters PM, Bae D, Snyder BD, Hresko MT, et al. Lateral entry compared with medial and lateral entry pin fixation for completely displaced supracondylar humeral fractures in children. A randomized clinical trial. *J Bone Joint Surg Am* 2007; 89: 706-12.
 16. Guner S, Guven N, Karadas S, Ceylan MF, Turktas U, Gokalp MA, et al. Iatrogenic or fracture-related nerve injuries in supracondylar humerus fracture: is treatment necessary for nerve injury? *Eur Rev Med Pharmacol Sci* 2013; 17: 815-9.
 17. Lyons JP, Ashley E, Hoffer MM. Ulnar nerve palsies after percutaneous cross-pinning of supracondylar fractures in children's elbows. *J Pediatr Orthop* 1998; 18: 43-5.
 18. Mayne AI, Perry DC, Stables G, Dhotare S, Bruce CE. Documentation of neurovascular status in supracondylar fractures and the development of an assessment proforma. *Emerg Med J* 2013; 30: 480-2.
 19. Kwak-Lee J, Kim R, Ebramzadeh E, Silva M. Is medial pin use safe for treating pediatric supracondylar humerus fractures? *J Orthop Trauma* 2013; 28: 216-21.
 20. Maity A, Saha D, Roy DS. A prospective randomised, controlled clinical trial comparing medial and lateral entry pinning with lateral entry pinning for percutaneous fixation of displaced extension type supracondylar fractures of the humerus in children. *J Orthop Surg* 2013; 7: 6. doi: 10.1186/1749-799X-7-6.
-