



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Medicine

Department of Medicine

January 2007

Chest radiographs in acute pulmonary embolism.

Ali Bin Sarwar Zubairi

Aga Khan University, ali.zubairi@aku.edu

Shahid Javed Husain

Aga Khan University

Muhammad irfan

Aga Khan University

Kulsoom Fatima

Aga Khan University, kulsoom.fatima@aku.edu

Muhammad Atif Zubairi

Aga Khan University

See next page for additional authors

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med



Part of the [Pulmonology Commons](#)

Recommended Citation

Zubairi, A., Husain, S. J., irfan, M., Fatima, K., Zubairi, M. A., Islam, M. (2007). Chest radiographs in acute pulmonary embolism.. *JAMC: Journal of Ayub Medical College, Abbottabad*., 19(1), 29-31.

Available at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med/314

Authors

Ali Bin Sarwar Zubairi, Shahid Javed Husain, Muhammad irfan, Kulsoom Fatima, Muhammad Atif Zubairi, and Muhammad Islam

CHEST RADIOGRAPHS IN ACUTE PULMONARY EMBOLISM

Ali Bin Sarwar Zubairi, Shahid Javed Husain, Muhammad Irfan, Kulsoom Fatima*,
Muhammad Atif Zubairi, Muhammad Islam**

Department of Pulmonary and Critical Care Medicine, *Department of Radiology, **Department of Medicine,
The Aga Khan University Hospital, Karachi, Pakistan

Background: Pulmonary embolism (PE) is a serious clinical entity carrying significant morbidity and mortality. Clinically, it is a difficult condition to diagnose and remains under treated condition in Pakistan due to non-availability of objective tests and lack of awareness among physicians. This study was conducted to determine the chest radiographic presentation in known cases of acute PE presenting to a tertiary care hospital. **Methods:** Hospital records of patients with a diagnosis of acute PE were reviewed from June 2000 until June 2004. Fifty diagnosed cases of acute PE on Spiral Computed tomography (CT) of the chest demonstrating an intraluminal-filling defect were selected. Two chest physicians reviewed the chest radiographs obtained during that hospitalization. In case of discrepancy, a radiologist made final interpretation. **Results:** The chest radiograph was interpreted as normal in only 18% of patients with acute PE. The most common chest radiographic abnormalities were cardiac enlargement (38%), pulmonary parenchymal infiltrates (34%), atelectasis (26%), pleural effusion (24%), and pulmonary congestion (24%). Other rare findings were elevated hemi diaphragm (14%), pulmonary artery enlargement (14%), and focal oligemia (8%). **Conclusions:** Cardiomegaly is the most common chest radiographic abnormality associated with acute pulmonary embolism. Chest radiography is not useful in making the diagnosis of acute pulmonary embolism. Its major role is in identification of alternative disease processes that can mimic thrombo-embolism

Keywords: Chest radiograph, Echocardiography, Pulmonary embolism

INTRODUCTION

Pulmonary embolism results in significant morbidity and mortality. It remains largely under diagnosed and under treated clinical problem amongst hospitalized patients in Pakistan due to non-availability of objective tests and lack of awareness among physicians. The diagnosis of PE is difficult because clinical features are nonspecific and all diagnostic tests have certain limitations^{1,2}. The ventilation-perfusion (V/Q) scan is non-diagnostic in 60-70% of patients^{1,2}. Helical (spiral) CT has limited sensitivity for sub-segmental embolism. Pulmonary angiogram is the 'gold standard' but is invasive, impractical or unavailable in most clinical settings³.

The chest radiograph is a fundamental test in the initial evaluation of cardiopulmonary disease and has advantage of being non-invasive⁴. It is not a diagnostic test for PE but is extremely helpful in evaluation of other common cardio-respiratory diseases, which mimic PE, like Congestive heart failure (CHF), Pneumonia and Pneumothorax. Data that describe chest radiographic findings associated with the diagnosis of pulmonary embolism are limited⁵.

The purpose of our study was to evaluate the chest radiographic findings in diagnosed cases of acute pulmonary embolism.

MATERIALS AND METHODS

We conducted a retrospective, cross-sectional study on chest radiographs of all patients admitted with a

diagnosis of acute PE from June 2000 until June 2004 at Aga Khan University, a tertiary care hospital in Karachi, Pakistan. We identified 50 consecutive patients who were diagnosed as acute pulmonary embolism on helical CT scan during the study period.

We carefully analyzed the chest radiograph findings along with gender, risk factors, associated diseases, clinical features, laboratory findings, echocardiography, treatment and outcome of these cases.

Two Pulmonologists interpreted chest radiographs. If there was any discrepancy, final interpretation was made by a Radiologist. The chest radiographs were characterized as normal or abnormal. If abnormal, then we looked for presence of one or more abnormal features like; cardiomegaly (cardio thoracic ratio > 0.5), pleural effusion, atelectasis (loss of lung volume), enlargement of a major pulmonary artery (Fleischner's sign), pulmonary parenchymal infiltrate, pulmonary congestion, elevated hemi diaphragm, focal oligemia (Westermark's sign) and a peripheral wedge-shaped opacity above the diaphragm (Hampton's hump).

Statistical package for social science (SPSS) version 11 was used for data analysis. The descriptive analysis was done for demographic, clinical, radiographic features and laboratory results. Results were expressed as mean \pm standard deviation, and number (percentage).

RESULTS

50 patients (21 males, 29 females) were identified during the study period. The mean age was 52 years (12 to 88 years). Almost all of the patients (98%) had symptoms of acute PE and only 2% were asymptomatic. Principal symptoms were breathlessness (88%), chest pain (32%), cough (26%), hemoptysis (8%) and wheezing (4%).

Risk factors included the following: prolonged bed rest or immobilization (34%), major surgery in past 3 months (26%), trauma in past 3 months (18%), malignancy (18%), hypercoagulable states (6%) and traumatic spinal cord injury (2%). Only one patient had positive family history of venous thromboembolism. None had prior history of venous thromboembolism (VTE), pregnancy or postpartum state or use of oral contraceptives.

Chest radiographs of all selected cases were interpreted. Only nine patients (18%) had normal chest radiographs and forty-one (82%) had abnormal chest radiographs. Table 1 describes the findings of chest radiograph.

Echocardiography was done in 29 patients (58%) out of 50. Right ventricular strain was identified in 14 patients (28%); Right ventricle (RV) hypo kinesis in 11 (22%), RV dilatation in 14 (28%), tricuspid regurgitation in 18 (36%), pulmonary hypertension in 18 (36%) and interventricular septal shift in 8 (16%).

Six patients died due to refractory shock and hypoxemia. Thrombolytics were given in 4 patients (8%) out of which 3 survived and discharged home. Embolectomy was done in 4 patients (8%). All of them survived and IVC filter was placed in 5 patients (10%).

Table 1. Chest radiographic abnormalities associated with acute pulmonary embolism

Abnormalities	(%)	95% Confidence Interval
Cardiomegaly	38%	(0.247, 0.513)
Infiltrate	34%	(0.209, 0.471)
Atelactasis	26%	(0.139, 0.381)
Pleural effusion	24%	(0.122, 0.358)
Pulmonary congestion	24%	(0.122, 0.358)
PA enlargement	14%	(0.044, 0.236)
Elevated hemi diaphragm	14%	(0.044, 0.236)
Oligemia	8%	(0.005, 0.155)

DISCUSSION

PE has been described as the most preventable and most under diagnosed cause of hospital deaths. The majority of preventable deaths associated with PE can be ascribed to a missed diagnosis rather than to failure of existing therapies. An estimated 600,000 cases of PE occur per year in United States resulting in 50,000-

100,000 fatalities and 5-10 % of hospital deaths⁶. No data is available from Pakistan. Wharton and Pierson⁷ reported the first radiographic description of pulmonary embolism in 1922.

Historically, many radiographic signs of pulmonary embolism have been described but their specificity has probably been overstated. One of the few prospective studies, which evaluated the reliability of chest radiograph in the diagnosis of PE, has shown that chest radiography in isolation is a notoriously poor indicator of the diagnosis.

In the Prospective Investigation of Pulmonary Embolism Diagnosis (PIOPED) study, the chest radiographs of most patients (88%) were abnormal⁸. Plain chest radiograph cannot be used itself to diagnose or exclude PE, but it may rule out other potentially life-threatening conditions such as pneumothorax⁹.

Although majority of patients with PE have abnormal chest x-ray findings, the most frequent abnormalities are non-specific as in our series. The highly specific findings of focal oligemia (Westermarck’s sign) or a peripheral wedge-shaped opacity (Hampton’s hump) are rare. No abnormalities are seen on plain chest films in approximately 10-15% cases.

The contrast-enhanced helical CT has replaced V/Q scan as the diagnostic study of choice in patients with suspected PE at most centers including our hospital as it allows the direct visualization of emboli as well as the detection of parenchymal abnormalities that may provide an alternate explanation for patient’s symptoms^{10,11}.

Our study extends previous observations of chest X-ray findings associated with acute PE. Patients with acute PE are likely to have an abnormal chest X-ray as reported by others. Cardiomegaly remains the most common radiographic abnormality associated with acute PE as reported by International Cooperative Pulmonary Embolism Registry (ICOPER) database¹². Unlike the ICOPER study, we did not isolate the chest X-ray abnormality unique to PE by eliminating patients with prior cardio-pulmonary disease.

Right ventricular hypo kinesis, detected by echocardiography, is an important predictor of mortality that is associated with acute pulmonary embolism¹³. The ICOPER data suggest that neither cardiomegaly nor PA enlargement is sensitive or specific predictors of RV hypo kinesis.

Our study has some limitations. First, the sample size was small. Second, we did not exclude patients with underlying cardiopulmonary disease, which may influence chest radiograph findings. Interestingly, despite our limitations the data is similar to ICOPER database.

CONCLUSION

In summary, our data suggests that cardiomegaly is the most common chest radiographic abnormality associated with acute PE. Although the chest radiograph cannot be used to diagnose or exclude PE, it contributes to the non-invasive diagnostic assessment of PE through the exclusion of disease processes that may mimic PE.

REFERENCES:

1. Value of the ventilation/perfusion scan in acute pulmonary embolism. Results of the prospective investigation of pulmonary embolism diagnosis (PIOPED). The PIOPED Investigators. *JAMA*. 1990 May 23-30;263(20):2753-9.
2. Johnson MS. Current strategies for the diagnosis of pulmonary embolus. *J Vasc Interv Radiol* 2002;13(1):13-23.
3. Hogg K, Brown G, Dunning J, Wright J, Carley S, Foex B, et al. Diagnosis of pulmonary embolism with CT pulmonary angiography: a systematic review. *Emerg Med J*. 2006; 23(3):172-8.
4. Wells PS, Ginsberg JS, Anderson DR, Kearon C, Gent M, Turpie AG, et al. Use of a clinical model for safe management of patients with suspected pulmonary embolism. *Ann Intern Med*. 1998;129(12): 997-1005.
5. Stein PD, Terrin ML, Hales CA, Palevsky HI, Saltzman HA, Thompson BT et al. Clinical, laboratory, roentgenographic, and electrocardiographic findings in patients with acute pulmonary embolism and no pre-existing cardiac or pulmonary disease. *Chest*. 1991; 100(3): 598-603.
6. Anderson FA Jr, Wheeler HB. Venous thromboembolism. Risk factors and prophylaxis. *Clin Chest Med*. 1995 Jun;16(2):235-51..
7. Wharton LR, Pierson JW. Minor forms of pulmonary embolism after abdominal operations. *JAMA* 1922;79: 1904-10.
8. Worsley DF, Alavi A, Aronchick JM, Chen JT, Greenspan RH, Ravin CE. Chest radiographic findings in patients with acute pulmonary embolism: Observations from the PIOPED Study. *Radiology* 1993;189(1):133-6.
9. Stein PD, Athanasoulis C, Greenspan RH, Henry JW. Relation of plain chest radiographic findings to pulmonary arterial pressure and arterial blood oxygen levels in patients with acute pulmonary embolism. *Am J Cardiol* 1992; 69(4): 394-96.
10. Usman MU, Bari V, Yaqoob J, Murad M. Diagnosis of Pulmonary Embolism with helical CT. *Scan. JPak Med Assoc*. 2003; 53(8): 354-56.
11. Schoepf UJ, Goldhaber SZ, Costello P. Spiral computed tomography for acute pulmonary embolism. *Circulation*. 2004; 109(18): 2160-67.
12. Elliott CG, Goldhaber SZ, Visani L, De Rosa M. Chest radiographs in acute pulmonary embolism: Results from the International Cooperative Pulmonary Embolism Registry. *Chest*. 2000; 118(1): 33-8.
13. A Goldhaber SZ, Visani L, De Rosa M. Acute pulmonary embolism: clinical outcomes in the International Cooperative Pulmonary Embolism Registry (ICOPER). *Lancet*. 1999; 353(9162):1386-9.

Address for Correspondence: Ali Bin Sarwar Zubairi; MD, FCCP, Assistant Professor, Pulmonary and Critical Care Medicine, The Aga Khan University Hospital, Stadium Road, P.O. Box 3500, Karachi 74800, Pakistan.. **Phone:** 92 (21) 4859- 4682, 4859-4685, **Fax:** 92 (21) 493 4294
Email: ali.zubairi@aku.edu