



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Medicine

Department of Medicine

April 2013

My patient had a small vessel stroke, are two antiplatelet agents better than one?

Muhammad Faisal Wadiwala

Aga Khan University

Ayeesha Kamran Kamal

Aga Khan University, ayeesha.kamal@aku.edu

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med



Part of the [Neurology Commons](#)

Recommended Citation

Wadiwala, M. F., Kamal, A. (2013). My patient had a small vessel stroke, are two antiplatelet agents better than one?. *JPMA. The Journal of the Pakistan Medical Association*, 63(4), 535-536.

Available at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med/356

My patient had a small vessel stroke, are two antiplatelet agents better than one?

Muhammad Faisal Wadiwala, Ayeesha Kamran Kamal

Why is this study of clinical importance?

Small subcortical brain infarcts, commonly known as lacunar strokes, constitute about 25% of ischaemic strokes. Aspirin is accepted as a standard antiplatelet therapy in patients with lacunar infarcts. Dual Antiplatelet therapy has been associated with increased bleeding, although it has been shown to reduce the risk of stroke among patients with atrial fibrillation.

The SPS3 trial was a randomized, multicenter clinical trial conducted in 82 clinical centers in North America, Latin America, and Spain. It tested two randomized interventions, in a 2-by-2 factorial design, in patients with MRI-confirmed lacunar stroke: clopidogrel and aspirin versus aspirin alone and two target levels of systolic blood pressure (<130 mm Hg vs. 130 to 149 mm Hg).

This trial was not pharmaceutically sponsored. It was an investigator initiated trial. The final results of the antiplatelet component of the trial have now been published.

Who were the participants?

Between 2003 and 2011, a total of 3020 patients were enrolled in the study: 1503 in the group treated with aspirin plus placebo and 1517 in the group treated with aspirin plus clopidogrel.

Patients were eligible for participation in the study if they were 30 years of age or older, had undergone a symptomatic lacunar stroke within the preceding 180 days, and did not have surgically amenable ipsilateral carotid artery disease or cardioembolism.

What was the intervention?

In accordance with the 2-by-2 factorial design of the study, eligible patients underwent

.....
Stroke Service and Vascular Fellowship Programme, International Cerebrovascular Translational Clinical Research Training Program (Fogarty International Center and National Institute of Neurological Disorders and Stroke), Aga Khan University Hospital, Karachi, Pakistan.

Correspondence: Ayeesha Kamran Kamal. Email: ayeesha.kamal@aku.edu

simultaneous randomization to the antiplatelet intervention (in which both patients and practitioners were unaware of group assignments) and to one of the two groups defined by target levels for systolic blood pressure (<130 mm Hg vs. 130 to 149 mm Hg) (with patients and practitioners aware of the group assignments).

All participants were given 325 mg of enteric-coated aspirin daily and were randomly assigned to receive 75 mg of clopidogrel daily or a matching placebo, with adherence measured by means of pill counts performed at quarterly follow-up visits.

What are the results?

The participants had a mean age of 63 years, and 63% were men. After a mean follow-up of 3.4 years, the risk of recurrent stroke was not significantly reduced with aspirin and clopidogrel (dual antiplatelet therapy) (2.5% per year) as compared with aspirin alone (2.7% per year) (hazard ratio, 0.92; 95% confidence interval [CI], 0.72 to 1.16), nor was the risk of recurrent ischaemic stroke (hazard ratio, 0.82; 95% CI, 0.63 to 1.09) or disabling or fatal stroke (hazard ratio, 1.06; 95% CI, 0.69 to 1.64). The risk of major haemorrhage was almost doubled with dual antiplatelet therapy (105 haemorrhages, 2.1% per year) as compared with aspirin alone (56, 1.1% per year) (hazard ratio, 1.97; 95% CI, 1.41 to 2.71; $P < 0.001$).

What were the conclusions?

Among patients with recent lacunar strokes, the addition of clopidogrel to aspirin did not significantly reduce the risk of recurrent stroke and did not significantly increase the risk of bleeding and death.

How does this impact us?

In Pakistan there is a high prevalence of Diabetes mellitus and Hypertension which frequently causes lacunar stroke. It appears from this trial, that at least in Lacunar strokes, two agents (aspirin and clopidogrel) are not better than one. In addition, given the lifelong nature of therapy after stroke, this data will assist in reducing prescription costs in a resource poor nation.

Acknowledgement and Disclosure Statement

The International Cerebrovascular Translational Clinical Research and Training Program (ICT_CRT) at the Aga Khan University is supported by funds from the Award Number D43TW008660 from the Fogarty International Center and the National Institute of Neurological Disorders and Stroke. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the Fogarty International Center, National Institute of Neurological Disorders and Stroke or the National Institutes of Health.

Recommended Reading

1. SPS3 Investigators. Benavente OR, Hart RG, McClure LA, Szychowski JM, Coffey CS, Pearce LA. Effects of clopidogrel added to aspirin in patients with recent lacunar stroke. *N Engl J Med* 2012; 367: 817-25.