



October 2013

Stroke radiology and distinguishing characteristics of intracranial atherosclerotic disease in native South Asian Pakistanis.

Maria Khan
Aga Khan University

Asif Rasheed
Aga Khan University

Saman Hashmi
Aga Khan University

Moazzam Zaidi
Aga Khan University

Saba Akhtar
Aga Khan University

See next page for additional authors

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med

 Part of the [Neurology Commons](#)

Recommended Citation

Khan, M., Rasheed, A., Hashmi, S., Zaidi, M., Akhtar, S., Bansari, L., Shah, N., Samuel, M., Raza, S., Khan, U. R., Ahmed, B., Ahmed, B., Ahmed, N., Ara, J., Ahsan, T., Munir, S. M., Ali, S., Mehmood, K., Maki, K. U., Ahmed, M. M., Sheikh, N., Memon, A. R., Frossard, P. M., Kamal, A. (2013). Stroke radiology and distinguishing characteristics of intracranial atherosclerotic disease in native South Asian Pakistanis. *International journal of stroke : official journal of the International Stroke Society*, 14-20.
Available at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med/368

Authors

Maria Khan; Asif Rasheed; Saman Hashmi; Moazzam Zaidi; Saba Akhtar; Lajpat Bansari; Nabi Shah; Maria Samuel; Sadaf Raza; Umer Rais Khan; Bilal Ahmed; Bilawal Ahmed; Naveeduddin Ahmed; Jamal Ara; Tasnim Ahsan; S. M. Munir; Shoukat Ali; Khalid Mehmood; Karim Ullah Maki; Muhammad Masroor Ahmed; Niaz Sheikh; Abdul Rauf Memon,; Philippe M. Frossard; and Ayeesha Kamran Kamal

Published in final edited form as:

Int J Stroke. 2013 October ; 8(0 100): . doi:10.1111/j.1747-4949.2012.00878.x.

Stroke radiology and distinguishing characteristics of intracranial atherosclerotic disease in native South Asian Pakistanis

Maria Khan, MBBS, FCPS^{1,*}, Asif Rasheed, MBBS^{2,*}, Saman Hashmi, MBBS², Moazzam Zaidi, MBBS², Muhammad Murtaza, MBBS², Saba Akhtar, MBBS², Lajpat Bansari, MBBS², Nabi Shah, M.Sc², Maria Samuel, M.Sc², Sadaf Raza, M.Sc², Umer Rais Khan, MBBS², Bilal Ahmed, MBBS¹, Bilawal Ahmed, MBBS¹, Naveeduddin Ahmed, FCPS³, Jamal Ara, FCPS, M.Phil⁴, Tasnim Ahsan, MRCP, FRCP⁴, S. M. Munir, FCPS⁴, Shoukat Ali, MD, FCPS⁴, Khalid Mehmood, FCPS MACG FRCP⁵, Karim Ullah Maki, FCPS⁵, Muhammad Masroor Ahmed, MCPS, FCPS⁵, Niaz Sheikh, FRCP⁵, Abdul Rauf Memon, FCPS⁵, Philippe M. Frossard, MSc, PhD², and Ayeesha Kamran Kamal, MD, FAHA, ABVN, ABPN¹

¹Stroke Service and the International Cerebrovascular Translational Clinical Research Training Program, Section of Neurology, Department of Medicine, Aga Khan University Hospital, Karachi, Pakistan

²Center for Non-Communicable Diseases, Karachi, Pakistan

³Liaquat National Hospital, Institute for Postgraduate Medical Studies and Health Sciences, Karachi, Pakistan

⁴Jinnah Postgraduate Medical Centre, Karachi, Pakistan

⁵Dow University of Health Sciences, Civil Hospital, Karachi, Pakistan

Abstract

Background—There are no descriptions of stroke mechanisms from intracranial atherosclerotic disease (ICAD) in native South Asian Pakistanis.

Methods—Men and women aged ≥ 18 years with acute stroke presenting to four tertiary care hospitals in Karachi, Pakistan were screened using Magnetic Resonance Angiography/Transcranial Doppler scans. TOAST criteria were applied to identify strokes from ICAD.

Results—245 patients with acute stroke due to ICAD were studied. 230 scans were reviewed. 206 /230(89.0%) showed acute ischemia.

The most frequent presentation was with cortically based strokes in 42.2% (87/206) followed by border zone infarcts (52/206, 25.2%).Increasing degrees of stenosis correlated with the development of both cortical and border zone strokes ($p=.002$). Important associated findings were frequent atrophy (166/230, 72.2%) silent brain infarcts (66/230, 28%) and a marked lack of severe leukoaraiosis identified in only 68 /230 (29.6 %).

A total of 1870 arteries were studied individually. MCA was the symptomatic stroke vessel in half, presenting with complete occlusion in 66%. Evidence of biological disease, symptomatic or

Corresponding Author: Ayeesha Kamran Kamal, MD, FAHA, ABVN, ABPN Associate Professor Neurology, Section of Neurology, Dept. of Medicine Director Stroke Fellowship Program Co- Director International Cerebrovascular Translational Clinical Research Training Program Aga Khan University Hospital Stadium Road – 74800 Karachi, Pakistan ayeesha.kamal@aku.edu Fax: 9221-493-4294 Phone: 9221-34930051-4559.

*These authors have contributed equally to this work.

asymptomatic was identified in 753 (40.2%) vessels of which 543 (72%) were significantly (>50%) stenosed at presentation.

Conclusion—ICAD is a diffuse process in Pakistani South Asians, with involvement of multiple vessels in addition to the symptomatic vessel. The MCA is the most frequent symptomatic vessel presenting with cortical embolic infarcts. There is a relative lack of leukoariosis. Concomitant atrophy, silent brain infarcts and recent ischemia in the symptomatic territory are all frequently associated findings.

INTRODUCTION

Pakistan is the sixth most populous country in the world with a population of 170 million¹. Over the last four decades, the incidence of stroke has increased by more than a 100% in middle-low income countries like Pakistan². The estimated lifetime prevalence of cerebrovascular disease in Pakistan has been reported to be 20%³ which translates into 34 million individuals with either stroke or TIA. Intracranial atherosclerotic disease (ICAD) may be the most common cause of stroke in the world especially in Pakistani South Asians^{4,5}. To inform better strategies for prevention and treatment of ICAD, a clearer recognition of its presentation and pathophysiologic mechanisms in susceptible populations will be helpful. In Pakistani South Asians however this has not been reported to date.

The Karachi Intracranial Stenosis Study (KISS) is a multicenter case-control study describing the patterns and determinants of intracranial disease in an urban Pakistani population- those most at risk for ICAD⁶. In this paper we describe the detailed radiologic presentations of ICAD stroke and their mechanistic implications.

METHODS

KISS is a multicenter, hospital based, case control study of patients with intracranial atherosclerosis and healthy stroke-free controls. Methods have been described in detail earlier⁶.

Study Site

Subjects were recruited from 4 tertiary care hospitals in Karachi, a southern metropolis of Pakistan with a socioeconomically diverse multiethnic population. Of the contributing hospitals, two are academic referral centers following fee-for-service models and the rest are heavily-subsidized publicly funded hospitals. Data collection was carried out from November 2007 to March 2010.

Sample Size

Over the study inclusion period, 1604 patients with stroke were reviewed. Of these 385 (24%) patients were found to have large artery atherosclerosis (intra or extracranial disease) as the stroke mechanism. Within this group 314 (81.5%) were diagnosed ICAD, rest were extracranial atherosclerosis. We further excluded 69 patients because the certainty of diagnosis was weaker for these patients for definite ICAD as a stroke mechanism. This study describes the radiology and stroke mechanisms of the 245 participants who had intracranial atherosclerotic disease (ICAD), as per the modified TOAST (Trial of ORG 10172 in Acute Stroke Treatment) criteria.⁷

Study Participants

All men and women aged 18 years or older presenting with stroke fulfilling WHO criteria were recruited by trained study physicians who actively reviewed all admissions to the

medicine or neurology wards on a daily basis. Neuro-imaging was mandatory for inclusion. Patients receiving medical attention after 72 hours since they were last seen normal, those diagnosed with non-vascular or iatrogenic stroke, those presenting with concomitant acute coronary syndrome or those who refused to consent were excluded from this study. Surrogate consent was sought if the patient was unable to consent independently.

Study procedures

Appropriate ancillary vascular and cardiac investigations to assign the patient to a diagnosis of ICAD according to TOAST criteria were reviewed in each individual case. No patient was assigned 'concurrent mechanism'. If a patient had atrial fibrillation, they were classified under cardioembolism and excluded for this review. For this analysis, those patients where the diagnosis of ICAD was most definite were selected. All scans were reviewed by a stroke neurologist allowing appropriate classification of each stroke patient; 10% of the scans were read by a second neurologist to assure reliability and consistency of the angiographic data with minimal inter-observer correlation..

A standardized Computerized Tomography (CT)/Magnetic Resonance Imaging (MRI) reading form adapted from the work of J Wardlaw⁸ was used to document imaging findings. Modifications made to the form included using angiographic data to record information on cerebral vasculature. Every artery was examined for the number, degree and distribution of stenosis. Biologically relevant disease ranging from atherosclerotic irregularity to measurable stenosis was noted and measured. Another modification was made to draw out the visualized strokes on "stroke maps". These were derived from published papers^{9, 10}. This was done to better visualize stroke topography in each individual patient. The following radiologic variables were reviewed in each patient: Stroke¹¹, acute stroke lesions¹², silent brain infarcts¹³, border zone infarcts¹⁴, cortical infarctions¹⁵, lacunar infarcts¹⁵, atrophy¹⁶, leukoaraiosis¹⁷, arterial review and assignment of symptomatic vs. asymptomatic status^{9, 10}.

All radiologic data was directly archived into a centralized computer using specially designed software. The software allowed for measurements to be made for calculation of intracranial stenosis using computerized calipers, magnify areas of interest, and quantify stroke volumes. We used the WASID method to calculate stenosis¹⁸.

Ethical Considerations

The study protocol was approved by the relevant Ethical Review Committee. Written informed consent and verbal assent was given by all participants or their surrogate respondents prior to inclusion.

Statistical Analysis

All data was entered and rechecked by two different investigators. Analysis was carried out using Statistical Package for Social Sciences (SPSS) Version 17.0 (SPSS Inc., Chicago, IL). We calculated simple frequencies for stroke locations and subtypes, symptomatic vessels, old vascular lesions and brain atrophy and leukoaraiosis. Means and standard deviations were computed for percentage of stenosis in various symptomatic and asymptomatic vessels and proportions were used for the categorical variables. Chi-square test of independence was applied for correlations between symptomatic vessel stenosis and stroke mechanism, as well as association of cerebral atrophy and leukoaraiosis with the number of diseased vessels. Two sided p-value of 0.05 was taken significant.

RESULTS

Stroke due to ICAD was the predominant mechanism of stroke due to large artery atherosclerosis. Two hundred and forty five patients with stroke due to definite ICAD presented to the study centers between November 2007 and March 2010. Scans (CT/MRI) data were missing for 15 of these patients, who were, therefore, excluded. Out of the final 230 scans reviewed, a majority were judged to be of good quality (72.2%) with only 8.3% being of poor quality. Most of the scans, 161 (70%) were MRIs with Diffusion Weighted Imaging, 49 (21%) were non contrast CT head and 20 (9%) were plain MRIs. MR Angiography (MRA) was available for 170 of these 230 patients.

Acute Stroke Findings

Out of the 230 scans, 24 (10.4%) had no sign of acute ischemic change at presentation. Of these 24, 5 scans were non contrast head CT, these were not repeated. The rest (n=206, 89%) had evidence of acute ischemia with about 11 (5.4%) showing a concomitant recent ischemic lesion. Nine of these patients had acute lesions in two territories. Only 7 (3%) showed hemorrhagic transformation, while 56 (27.2%) had evidence of mass effect with large infarctions. Old vascular lesions were seen in 66 of the 230 scans (28%). Most of these were either old lacunar strokes (43.9%) or old cortical infarcts (33.3%). Only two of these old lesions were hemorrhagic. There were no non-stroke lesions identified in the 230 scans.

Stroke Distributions and Mechanisms

(Figure 1) Of the acute strokes 151/206 (73.3%) were in the anterior circulation and the rest (55/206, 26.7%) were in the posterior circulation. The majority were cortically based strokes (87/206, 42.2%), followed by border-zone infarcts (52/206, 25.2%). Lacunar infarcts were 23 /206 (16%) and were the least common presentation. Cortically based strokes were more frequent in the middle cerebral artery (MCA) territory (66/87; 75.8%). Of the borderzone infarctions, majority were in the internal borderzone (37/52; 71.2%).

Correlation with stroke mechanism and vessel stenosis

(Figure 2) As the degree of arterial stenosis increased from <50% to complete occlusion, there was a statistically significant increase observed in both cortical and borderzone infarcts ($p=0.002$). This difference was more marked for cortically based infarctions than for borderzone infarctions and was negligible for lacunar strokes.

Description of Angiographic findings

(Figure 3) A total of 1,870 intracranial arteries of 170 patients were studied. Of these 170 patients, 151 had acute symptomatic stroke. MCA was the symptomatic vessel in nearly half the patients (69/151, 45.7%) with complete occlusion observed in 66% of these MCAs. ICA and Basilar (BA) were the next common culprit vessels (25/151 each, 16.6%) (Table 1).

Evidence of biological disease (any degree of stenosis) was identified in 753 of the 1870 (40.2%) vessels studied. Significant stenosis (>50%) was found in 543 of the 753 (72%) arteries; 433 of these 543 (79.9%) vessels with significant stenosis were asymptomatic (i.e. not corresponding to the arterial territory of the presenting ischemic change). The greatest number of asymptomatic significant stenosis was observed in PCAs. A mean of 4.4 vessels were diseased per patient with the highest number of affected vessels being 10 of 11.

Associated Radiologic findings and correlations

Important associated radiologic findings were frequent atrophy 166 /230 (72.2%) and the marked lack of leukoaraiosis identified in only 68 of the 230 patients (29.6%). There were

no significant associations between number of diseased vessels and leukoaraiosis ($p = 0.55$), as well as central/ cortical atrophy ($p = 0.87, 0.96$ respectively).

DISCUSSION

We found that ICAD presents most frequently with cortical embolic stroke followed by internal borderzone ischemia in Pakistani South Asians. The stroke is most often due to advanced stenosis of the MCA, and the chances of getting a disabling cortically based stroke correlates with advancing degrees of stenosis. ICAD interestingly, not infrequently presents with lacunar stroke, however this presentation is not influenced by increasing stenosis in the parent artery. Noteworthy concomitants are, associated remote silent brain infarcts, of which one third were cortical strokes, atrophy, recent ischemia, and diffuse atherosclerosis. Unexpectedly infrequent is diffuse white matter disease—given the distribution of risk factors for both being similar – this observation is puzzling and unclear.

The mechanisms of stroke from ICAD in Pakistanis are heterogeneous and it is important for the clinician to recognize these mechanisms for their potential therapeutic implications. Studies from similar populations have shown that cortical embolic stroke in ICAD are often accompanied by platelet microembolism measurable by doppler ultrasound¹⁹. Clinical intervention by the use of double antiplatelet agents in this setting^{20, 21} have shown both reductions in micro-embolic signals and a reduction of recurrent stroke. Internal border-zone stroke, the second commonest mechanism, when measured by Positron Emission Tomography (PET) shows increased Oxygen Extraction Fraction (OEF)²² and by Single Photon Emission CT (SPECT) shows decreased reserve on acetazolamide challenge²³ and is often also prone to progressive infarction^{24, 25}. In actuality there are very few ICAD based studies that tell us what intervention would be reasonable in the face of acute border-zone ischemia. However emerging data suggests that it may not be safe to acutely reduce pressure aggressively²⁶ and perhaps increasing pressure may enhance flow through collaterals and stabilize the progression.

Important also are the lacunar presentations of ICAD, these have been described in other populations due to atheroma in the parent artery²⁷. Getting an expensive intracranial MRA, when the presentation is lacunar is not intuitional for most developing country neurologists, but recognizing this presentation for prognostic reasons is essential. ICAD carries the worst recurrent stroke rate (15% at one year)²⁸ amongst all ischemic stroke presentations- and at this time evidence from Warfarin versus Aspirin in Symptomatic Intracranial Disease (WASID) study clearly informs us that a strategy of aggressive risk factor intervention is the only way forward²⁹.

Important to note is that although the chances of stroke increased with increasing degree of stenosis, strokes of varied mechanisms were seen at all degrees of stenosis³⁰. Autopsy studies on fatal strokes³¹ and high strength magnets³² are informing us that we need to appreciate ICAD as a mechanism of stroke at lesser degrees of stenosis as plaque hemorrhage and instability will ensue in both kinds of strokes regardless of the degree of initial stenosis³³. The associated “preclinical silent brain infarcts” that we report are an additional observation supporting the emergent data on the effect of less than 50% stenosis on brains.

Evidence of atherosclerotic disease was present in nearly half the vessels studied, with more than 70% vessels having significant (>50%) stenosis. Sixty five percent of the culprit vessels had >70% stenosis in our study as compared to WASID which reports 40% of its subjects having severe stenosis (>70% in single or >50% in multiple arteries)³⁴. A study

from the Chinese population showed significant (50%) stenosis in 65% of their patients³⁵. 72.8% of our patients had 50% stenosis in their culprit vessels.

The ratio of asymptomatic stenosis was considerably high in our study (79% vs. 27% in MRAs reviewed for WASID)³⁶. There are currently limited studies on asymptomatic cerebrovascular disease. MRAs in healthy Japanese and Koreans had 3-3.5% ICAD^{37, 38}. Some studies suggest that asymptomatic intracranial disease has a low risk of recurrence^{36, 39}. At least in our cross sectional observation, the involvement of the middle cerebral artery was more likely to be symptomatic and the anterior and posterior cerebral asymptomatic, suggesting that location may determine asymptomatic status due to collaterals⁴⁰. This hypothesis will need further testing. At the very least, if there is diffuse involvement, intervention in high risk populations presents an opportunity to prevent symptomatic stroke. At this stage intervention may not necessarily be technically demanding and expensive like stents or bypass- both difficult interventions in most resource poor countries where ICAD is common.

Leukoaraiosis was surprisingly low. As the burden of hypertension is significantly high in our population, more was expected. In some Asian populations, significant association of leukoaraiosis has been reported with large artery atherosclerosis²⁷ and with multiple vessels³⁵. We are puzzled by this observation and unclear as to why there is small vessel protection.

Lastly the cortical atrophy may be the result of chronic ischemia of multiple vessels, as in a recent observation, revascularization reverses cortical thinning in these patients⁴¹ although our data set failed to find a statistically significant association.

Our study is unique because the disease burden in an individual patient of ICAD has never been quantified before in native South Asians. Our participating locations are national referral centers for stroke, so there is broader generalizability. We followed a systemized, centralized, standardized and supervised protocol. All our variables were predefined and measured exactly using international guidelines. The main limitation of this study is its cross sectional design and lack of longitudinal data. Our study population was urban and hospital based, and our investigation uses non invasive modalities, which might lead to detection bias of the true burden of ICAD. Also, not all scans were MRIs; CT may have missed out on some of the asymptomatic strokes. This may have decreased the detection rate of silent strokes –and the asymptomatic disease burden may be underreported.

Looking at the broader picture, studies aimed at alleviating stenosis in a single vessel may be ignoring the larger problem which perhaps may respond to global interventions aimed at halting the process of progressive pre-symptomatic ICAD⁴².

Acknowledgments

The authors would like to acknowledge the patience, kindness, time and cooperation of all patients and their families who contributed to this study.

Funding Disclosures:

The KISS project was funded by a grant awarded to Ayesha Kamran Kamal (PI) and Philippe M. Frossard (Co PI) by the Higher Education Commission of the Government of Pakistan. Ayesha Kamran Kamal was also the recipient of a seed money grant from the University Research Council. Dr Maria Khan is a neurovascular research fellow whose training is currently funded by Award Number D43TW008660 from the Fogarty International Center and the National Institute of Neurologic Disorders and Stroke. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Fogarty International Center, National Institute of Neurologic Disorders and Stroke or the National Institute of Health.


REFERENCES

1. [03-11-2010] [Www.Census.Gov.Pk](http://www.Census.Gov.Pk)
2. Feigin VL, Lawes CM, Bennett DA, Barker-Collo SL, Parag V. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: A systematic review. *Lancet Neurol.* 2009; 8:355–369. [PubMed: 19233729]
3. Kamal AK, Itrat A, Murtaza M, Khan M, Rasheed A, Ali A, Akber A, Akber Z, Iqbal N, Shoukat S, Majeed F, Saleheen D. The burden of stroke and transient ischemic attack in pakistan: A community-based prevalence study. *BMC Neurol.* 2009; 9:58. [PubMed: 19948076]
4. Caplan LR, Gorelick PB, Hier DB. Race, sex and occlusive cerebrovascular disease: A review. *Stroke.* 1986; 17:648–655. [PubMed: 3526645]
5. Wong LK. Global burden of intracranial atherosclerosis. *Int J Stroke.* 2006; 1:158–159. [PubMed: 18706036]
6. Kamal AK, Taj F, Junaidi B, Rasheed A, Zaidi M, Murtaza M, Iqbal N, Hashmat F, Alam SV, Saleem U, Waheed S, Bansari L, Shah N, Samuel M, Yameen M, Naz S, Khan FS, Ahmed N, Mahmood K, Sheikh N, Makki KU, Ahmed MM, Memon AR, Wasay M, Syed NA, Khealani B, Frossard PM, Saleheen D. The karachi intracranial stenosis study (kiss) protocol: An urban multicenter case-control investigation reporting the clinical, radiologic and biochemical associations of intracranial stenosis in pakistan. *BMC Neurol.* 2009; 9:31. [PubMed: 19604359]
7. Adams HPBB Jr, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE 3rd. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. *Stroke.* 1993; 24:35–41. [PubMed: 7678184]
8. [03-11-2010] Sbirc website, www.Sbirc.Ed.Ac.Uk/imageanalysis.Html, j wardlaw
9. Tatu L, Moulin T, Bogousslavsky J, Duvernoy H. Arterial territories of the human brain: Cerebral hemispheres. *Neurology.* 1998; 50:1699–1708. [PubMed: 9633714]
10. Tatu L, Moulin T, Bogousslavsky J, Duvernoy H. Arterial territories of human brain: Brainstem and cerebellum. *Neurology.* 1996; 47:1125–1135. [PubMed: 8909417]
11. Asplund K, Tuomilehto J, Stegmayr B, Wester PO, Tunstall-Pedoe H. Diagnostic criteria and quality control of the registration of stroke events in the monica project. *Acta Med Scand Suppl.* 1988; 728:26–39. [PubMed: 3202029]
12. Wardlaw JM, Sellar R. A simple practical classification of cerebral infarcts on ct and its interobserver reliability. *AJNR Am J Neuroradiol.* 1994; 15:1933–1939. [PubMed: 7863945]
13. Vermeer SE, Longstreth WT Jr, Koudstaal PJ. Silent brain infarcts: A systematic review. *Lancet Neurol.* 2007; 6:611–619. [PubMed: 17582361]
14. Damasio H. A computed tomographic guide to the identification of cerebral vascular territories. *Arch Neurol.* 1983; 40:138–142. [PubMed: 6830451]
15. Rovira A, Grive E, Rovira A, Alvarez-Sabin J. Distribution territories and causative mechanisms of ischemic stroke. *Eur Radiol.* 2005; 15:416–426. [PubMed: 15657788]
16. Farrell C, Chappell F, Armitage PA, Keston P, MacLulich A, Shenkin S, Wardlaw JM. Development and initial testing of normal reference mr images for the brain at ages 65-70 and 75-80 years. *Eur Radiol.* 2009; 19:177–183. [PubMed: 18690455]
17. van Swieten JC, Hijdra A, Koudstaal PJ, van Gijn J. Grading white matter lesions on ct and mri: A simple scale. *J Neurol Neurosurg Psychiatry.* 1990; 53:1080–1083. [PubMed: 2292703]
18. Samuels OB, Joseph GJ, Lynn MJ, Smith HA, Chimowitz MI. A standardized method for measuring intracranial arterial stenosis. *AJNR Am J Neuroradiol.* 2000; 21:643–646. [PubMed: 10782772]
19. Wong KS, Gao S, Chan YL, Hansberg T, Lam WW, Droste DW, Kay R, Ringelstein EB. Mechanisms of acute cerebral infarctions in patients with middle cerebral artery stenosis: A diffusion-weighted imaging and microemboli monitoring study. *Ann Neurol.* 2002; 52:74–81. [PubMed: 12112050]
20. Wong KS, Chen C, Fu J, Chang HM, Suwanwela NC, Huang YN, Han Z, Tan KS, Ratanakorn D, Chollate P, Zhao Y, Koh A, Hao Q, Markus HS. Clopidogrel plus aspirin versus aspirin alone for reducing embolisation in patients with acute symptomatic cerebral or carotid artery stenosis (clair


- study): A randomised, open-label, blinded-endpoint trial. *Lancet Neurol.* 2010; 9:489–497. [PubMed: 20335070]
21. Nguyen TN, Zaidat OO, Gupta R, Nogueira RG, Tariq N, Kalia JS, Norbash AM, Qureshi AI. Balloon angioplasty for intracranial atherosclerotic disease: Periprocedural risks and short-term outcomes in a multicenter study. *Stroke.* 2011; 42:107–111. [PubMed: 21071722]
 22. Liebeskind DS. Collateral circulation. *Stroke.* 2003; 34:2279–2284. [PubMed: 12881609]
 23. Ozgur HT, Kent Walsh T, Masaryk A, Seeger JF, Williams W, Krupinski E, Melgar M, Labadie E. Correlation of cerebrovascular reserve as measured by acetazolamide-challenged spect with angiographic flow patterns and intra- or extracranial arterial stenosis. *AJNR Am J Neuroradiol.* 2001; 22:928–936. [PubMed: 11337339]
 24. Del Sette M, Eliasziw M, Streifler JY, Hachinski VC, Fox AJ, Barnett HJ. Internal borderzone infarction: A marker for severe stenosis in patients with symptomatic internal carotid artery disease. For the north american symptomatic carotid endarterectomy (nascet) group. *Stroke.* 2000; 31:631–636. [PubMed: 10700496]
 25. Yamauchi H, Nishii R, Higashi T, Kagawa S, Fukuyama H. Hemodynamic compromise as a cause of internal border-zone infarction and cortical neuronal damage in atherosclerotic middle cerebral artery disease. *Stroke.* 2009; 40:3730–3735. [PubMed: 19797182]
 26. Sandset EC, Bath PM, Boysen G, Jatuzis D, Korv J, Luders S, Murray GD, Richter PS, Roine RO, Terent A, Thijs V, Berge E. The angiotensin-receptor blocker candesartan for treatment of acute stroke (scast): A randomised, placebo-controlled, double-blind trial. *Lancet.* 2011; 377:741–750. [PubMed: 21316752]
 27. Lee DK, Kim JS, Kwon SU, Yoo SH, Kang DW. Lesion patterns and stroke mechanism in atherosclerotic middle cerebral artery disease: Early diffusion-weighted imaging study. *Stroke.* 2005; 36:2583–2588. [PubMed: 16269637]
 28. Turan TN, Maida L, Cotsonis G, Lynn MJ, Romano JG, Levine SR, Chimowitz MI. Failure of antithrombotic therapy and risk of stroke in patients with symptomatic intracranial stenosis. *Stroke.* 2009; 40:505–509. [PubMed: 19095991]
 29. Turan TN, Derdeyn CP, Fiorella D, Chimowitz MI. Treatment of atherosclerotic intracranial arterial stenosis. *Stroke.* 2009; 40:2257–2261. [PubMed: 19407238]
 30. Klein IF, Labreuche J, Lavalley PC, Mazighi M, Duyckaerts C, Haww JJ, Amarenco P. Is moderate atherosclerotic stenosis in the middle cerebral artery a cause of or a coincidental finding in ischemic stroke? *Cerebrovasc Dis.* 2010; 29:140–145. [PubMed: 19955738]
 31. Mazighi M, Labreuche J, Gongora-Rivera F, Duyckaerts C, Haww JJ, Amarenco P. Autopsy prevalence of intracranial atherosclerosis in patients with fatal stroke. *Stroke.* 2008; 39:1142–1147. [PubMed: 18309170]
 32. Arenillas JF. Intracranial atherosclerosis: Current concepts. *Stroke.* 2011; 42:S20–23. [PubMed: 21164126]
 33. Ogata J, Masuda J, Yutani C, Yamaguchi T. Mechanisms of cerebral artery thrombosis: A histopathological analysis on eight necropsy cases. *J Neurol Neurosurg Psychiatry.* 1994; 57:17–21. [PubMed: 8301298]
 34. Turan TN, Makki AA, Tsappidi S, Cotsonis G, Lynn MJ, Cloft HJ, Chimowitz MI. Risk factors associated with severity and location of intracranial arterial stenosis. *Stroke.* 2010; 41:1636–1640. [PubMed: 20616323]
 35. Pu Y, Liu L, Zou X, Chen P, Wang Y, Zhou Y, Dong K, Zhao X, Wang C, Wang Y. Relationship between leukoaraiosis and cerebral large artery stenosis. *Neurol Res.* 2009; 31:376–380. [PubMed: 19508822]
 36. Nahab F, Cotsonis G, Lynn M, Feldmann E, Chaturvedi S, Hemphill JC, Zweifler R, Johnston K, Bonovich D, Kasner S, Chimowitz M. Prevalence and prognosis of coexistent asymptomatic intracranial stenosis. *Stroke.* 2008; 39:1039–1041. [PubMed: 18239161]
 37. Park KY, Chung CS, Lee KH, Kim GM, Kim YB, Oh K. Prevalence and risk factors of intracranial atherosclerosis in an asymptomatic korean population. *J Clin Neurol.* 2006; 2:29–33. [PubMed: 20396482]

38. Wong KS, Ng PW, Tang A, Liu R, Yeung V, Tomlinson B. Prevalence of asymptomatic intracranial atherosclerosis in high-risk patients. *Neurology*. 2007; 68:2035–2038. [PubMed: 17548555]
39. Famakin BM, Chimowitz MI, Lynn MJ, Stern BJ, George MG. Causes and severity of ischemic stroke in patients with symptomatic intracranial arterial stenosis. *Stroke*. 2009; 40:1999–2003. [PubMed: 19407228]
40. Liebeskind DS, Cotsonis GA, Saver JL, Lynn MJ, Turan TN, Cloft HJ, Chimowitz MI. Collaterals dramatically alter stroke risk in intracranial atherosclerosis. *Ann Neurol*. 2010
41. Fierstra J, Maclean DB, Fisher JA, Han JS, Mandell DM, Conklin J, Poublanc J, Crawley AP, Regli L, Mikulis DJ, Tymianski M. Surgical revascularization reverses cerebral cortical thinning in patients with severe cerebrovascular steno-occlusive disease. *Stroke*. 2011; 42:1631–1637. [PubMed: 21493908]
42. Spence JD, Hackam DG. Treating arteries instead of risk factors: A paradigm change in management of atherosclerosis. *Stroke*. 2010; 41:1193–1199. [PubMed: 20413738]

A. CORTICAL EMBOLIC INFARCTION a: Left anterior cerebral artery infarct; b & c: Left middle cerebral artery infarcts; d: Bilateral cerebellar infarcts



B. BORDERZONE TERRITORIAL INFARCTIONS: e: left anterior and bilateral posterior borderzone infarcts; f & g: Internal borderzone infarcts



C. LACUNAR STROKES (h and i)




Figure 1. Diffusion weighted images showing the stroke location and mechanisms in order of frequency – most frequent to least

A. CORTICAL EMBOLIC INFARCTION a: Left anterior cerebral artery infarct; b & c: Left middle cerebral artery infarcts; d: Bilateral cerebellar infarcts

B. BORDERZONE TERRITORIAL INFARCTIONS: e: left anterior and bilateral posterior borderzone infarcts; f & g: Internal borderzone infarcts




Figure 2. Correlation of stroke mechanisms with increasing degree of atherosclerotic stenosis ($p=0.002$). Advanced stenosis increases hemodynamic strokes and major cortical embolisms




Figure 3. MRA images showing diffuse severe atherosclerotic disease in South Asian Patients with Intracranial Atherosclerotic Disease (ICAD)

a-d: anterior circulation; frequent bilateral MCA involvement in addition to the symptomatic vessel; **e-h:** posterior circulation; frequent PCA disease in addition to the symptomatic vessel

Table 1

Symptomatic vessel and atherosclerotic stenosis in decreasing order of frequency (n=151)

Vessel	Symptomatic vessel, n (%)	Mean stenosis (SD)	100% stenosis, n (%)
MCA	69 (45.7)	77.4 (36.3)	46 (66)
ICA	25 (16.6)	81.5 (27.6)	16(64)
Basilar	25 (16.6)	44.6 (45.2)	8 (32)
PCA	23 (15.2)	67.6 (39.7)	11 (47.8)
ACA	7 (4.6)	85.7 (37.7)	6 (85.7)
Vertebrals	2(1.3)	72.1 (39.3)	1 (50)

ICA= internal carotid artery; MCA= middle cerebral artery; ACA= anterior cerebral artery; PCA= posterior cerebral artery;