



University of HUDDERSFIELD

University of Huddersfield Repository

Lünse, Christina E., Scott, Fraser J., Suckling, Colin J. and Mayer, Günter

Novel TPP-riboswitch activators bypass metabolic enzyme dependency

Original Citation

Lünse, Christina E., Scott, Fraser J., Suckling, Colin J. and Mayer, Günter (2014) Novel TPP-riboswitch activators bypass metabolic enzyme dependency. *Frontiers in Chemistry*, 2. ISSN 2296-2646

This version is available at <http://eprints.hud.ac.uk/id/eprint/32651/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>



Novel TPP-riboswitch activators bypass metabolic enzyme dependency

Christina E. Lünse¹, Fraser J. Scott², Colin J. Suckling^{2*} and Günter Mayer^{1*}

¹ Life and Medical Sciences Institute, University of Bonn, Bonn, Germany

² Department of Pure and Applied Chemistry, University of Strathclyde, Glasgow, UK

Edited by:

Youla S. Tzantrizos, McGill University, Canada

Reviewed by:

Wenshe R. Liu, Texas A&M University, USA
Krister Wennerberg, University of Helsinki, Finland

*Correspondence:

Colin J. Suckling, Department of Pure and Applied Chemistry, WestCHEM Research School, University of Strathclyde, 295 Cathedral Street, Glasgow, G1 1XL, Scotland, UK
e-mail: c.j.suckling@strath.ac.uk;
Günter Mayer, Life and Medical Sciences Institute, University of Bonn, Gerhard-Domagk-Str. 1, 53121 Bonn, Germany
e-mail: gmayer@uni-bonn.de


Riboswitches are conserved regions within mRNA molecules that bind specific metabolites and regulate gene expression. TPP-riboswitches, which respond to thiamine pyrophosphate (TPP), are involved in the regulation of thiamine metabolism in numerous bacteria. As these regulatory RNAs are often modulating essential biosynthesis pathways they have become increasingly interesting as promising antibacterial targets. Here, we describe thiamine analogs containing a central 1,2,3-triazole group to induce repression of *thiM*-riboswitch dependent gene expression in different *E. coli* strains. Additionally, we show that compound activation is dependent on proteins involved in the metabolic pathways of thiamine uptake and synthesis. The most promising molecule, triazolethiamine (TT), shows concentration dependent reporter gene repression that is dependent on the presence of thiamine kinase ThiK, whereas the effect of pyrithiamine (PT), a known TPP-riboswitch modulator, is ThiK independent. We further show that this dependence can be bypassed by triazolethiamine-derivatives that bear phosphate-mimicking moieties. As triazolethiamine reveals superior activity compared to pyrithiamine, it represents a very promising starting point for developing novel antibacterial compounds that target TPP-riboswitches. Riboswitch-targeting compounds engage diverse endogenous mechanisms to attain *in vivo* activity. These findings are of importance for the understanding of compounds that require metabolic activation to achieve effective riboswitch modulation and they enable the design of novel compound generations that are independent of endogenous activation mechanisms.

Keywords: riboswitches, triazolethiamine, click-chemistry, metal-chelating compounds, metabolic enzymes

INTRODUCTION

Riboswitches are RNA elements mostly found in the 5' UTR of bacterial mRNA that specifically sense the concentration of a small metabolite. Upon metabolite binding, substantial conformational changes occur that ultimately result in an on- or off-switch of gene expression. These regulatory elements represent a fundamental new means of controlling cellular processes in response to environmental conditions (Breaker, 2011). Riboswitches often regulate expression of essential genes and as such they are interesting target structures for the development of novel antibiotic compounds (Blount and Breaker, 2006). Artificial compounds acting successfully on riboswitches have to meet at least two criteria: (i) they must specifically bind to the relevant RNA structure and (ii) induce the conformational changes that finally lead to down- or up-regulation of gene expression. In recent years synthetic molecules that meet these criteria have been discovered for several riboswitch classes (Lünse et al., 2014). One of them is the TPP-riboswitch (Miranda-Rios et al., 2001; Winkler et al., 2002), which selectively interacts with thiamine pyrophosphate (TPP) by binding to its pyrimidine moiety through the so-called pyrimidine sensor helix (P2, J2-3, P3) of the aptamer domain. Conformational changes of the riboswitch resulting in repression of gene expression are only achieved if

the pyrophosphate group of the ligand is recognized by the pyrophosphate sensor helix (P4, J4-5, P5, **Figure 1A**) (Edwards and Ferre-D'Amare, 2006; Serganov et al., 2006; Thore et al., 2006). While this riboswitch class has been shown to allow a greater degree of variation at the position of the thiazole ring, exemplified by the thiamine analog pyrithiamine (PT, **Figure 2A**), the presence of the pyrophosphate moiety is mandatory to interact with and, more importantly, to induce switching of the *E. coli thiM*-riboswitch (Winkler et al., 2002; Rentmeister et al., 2007). This necessity impedes the development of compounds that act on TPP-riboswitches, since either they cannot passively pass the cell wall due to the highly charged pyrophosphate moiety or they cannot efficiently activate the riboswitch if lacking this group. PT's *in vivo* activity most likely relies on hijacking endogenous metabolic enzymes that phosphorylate exogenously added PT, thereby yielding the active derivative pyrithiamine pyrophosphate (PTPP) (Iwashima et al., 1976). In this report we elaborate on a series of triazolethiamines (TT) that have been shown recently to act on TPP-riboswitches *in vitro*, if diphosphorylated yielding triazolethiamine pyrophosphate (TTPP). We investigated TT-activity in relation to the length and modification of the alkyl chain and the presence of endogenous proteins involved in thiamine metabolism. We found that the activity of triazolethiamine




(TT) depends on the metabolic enzyme thiamine kinase (ThiK) and thiamine transporters. Most importantly, we demonstrated that ThiK dependency is bypassed by triazolethiamine-derivatives that bear phosphate mimicking and metal-ion chelating moieties. This strategy opens new avenues toward riboswitch activating compounds, in particular those that rely on phosphate groups to recognize and switch cognate RNA structures.

MATERIALS AND METHODS

β -GALACTOSIDASE REPORTER GENE ASSAY

For β -galactosidase assays 5 ml pre-cultures of DH5 α Z1, BW25113 or Keio deletion strains containing the appropriate plasmid constructs were prepared in LB Lennox standard medium (lysogeny broth Lennox, 10 g/l tryptone, 5 g/l yeast extract, 5 g/l NaCl in water) and incubated over night at 37°C and 155 rpm. Incubation was followed by photometric measurement of optical density at $\lambda = 600$ nm (OD₆₀₀) and dilution of cells to an optical density of 0.5. This dilution was used to inoculate β -galactosidase expression cultures at a ratio of 1:500 in a final volume of 2 or 4 ml in M9 medium (5x M9 medium containing 15 g/l KH₂PO₄, 5 g/l NH₄Cl, 2.5 g/l NaCl, 30 g/l Na₂HPO₄) containing final concentrations of 5 mM MgSO₄, 0.2 wt% glucose, 0.2 μ g/ μ l casamino acids (Difco), and 100 μ g/ml ampicillin. Even though vitamin-deprived casamino acids were used, growth of thiamine auxotrophic strains was observed without addition of thiamine indicating that a small, but constant amount of thiamine must be present in the casamino acid stock. Minimal inhibitory concentration determinations in casamino acid free minimal medium with separately added amino acids revealed that at least 1 nM of thiamine is necessary to enable bacterial growth



of the thiamine auxotrophs investigated. The expression cultures contained either 20 or 500 μ M of thiamine (Sigma), 500 μ M pyrithiamine (Sigma) or 500 μ M of the compounds, respectively. Controls received neither thiamine, or pyrithiamine nor compounds. If compounds had to be dissolved in DMSO, controls were supplemented with equal amounts (final DMSO concentration: 1%). The cultures were incubated for 24 h at 37°C and 150 rpm. After 24 h incubation the optical density was measured. The cells were centrifuged at 4500 g for 5 min and the pellets were washed twice in 400 μ l 1x PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄·2H₂O, 2 mM KH₂PO₄, pH 7.4). The pellets were resuspended in 200 μ l 1x lysis buffer (Reporter Lysis Buffer, Promega), incubated for 15 min at room temperature and cells were pelleted. Supernatants were used for all following steps. 75 μ l of the respective lysates were mixed with 75 μ l of 2x assay buffer (200 mM sodium phosphate buffer pH 7.3, 2 mM MgCl₂, 100 mM β -mercaptoethanol, 1.33 mg/ml ONPG) and incubated for 5–15 min at 37°C. The reaction was stopped by the addition of 250 μ l 1 M Na₂CO₃ and absorbance at 420 nm was

measured immediately using a Nanoquant (Infinite 200, Tecan). Reporter gene assays were performed in clear, flat bottom 96-well plates. Miller units were calculated according to the following formula: Miller units = $(1000 \times \text{OD}_{420}) / (\text{OD}_{600} \times \text{incubation time [min]} \times \text{culture volume [L]})$ and Miller units of controls without thiamine or compound added were set to 1. Samples were run in duplicate and experiments were repeated at least three times.

RESULTS

Thiamine analogs that bear a 1,2,3-triazole group instead of the thiazole heterocyclic moiety as found in thiamine were synthesized and investigated for TPP-riboswitch activation using a β -galactosidase reporter gene assay (Figure 1B). In this assay the TPP-riboswitch *thiM* from *E. coli* was cloned into the 5'-UTR of β -galactosidase. The DH5 α Z1 *E. coli* strain was transformed by this plasmid (Simons et al., 1987; Diederich et al., 1992; Lutz and Bujard, 1997). Due to the thi-1 mutation, DH5 α Z1 cells are thiamine auxotroph, which means they are dependent on thiamine uptake from the growth medium (Supplementary Table 1). This enables direct external control of the availability of this vitamin. Synthesis of TT-compounds was achieved using a “click” chemistry approach utilizing a common azide intermediate, which was then further reacted with the appropriate substituted alkyne yielding corresponding thiamine analogs (Figure 2A, Supplementary Figure 1). Subsequent monitoring of β -galactosidase activity was performed in the presence and absence of thiamine or triazolethiamine (TT) derivatives. Addition of thiamine to the *E. coli* cultures resulted in a complete loss of β -galactosidase expression; an IC₅₀-value of 0.015 μ M was observed (Figure 2 and Table 1). In line with previous findings, pyrithiamine (PT) was shown incapable of β -galactosidase repression in DH5 α Z1 (Figure 2B) (Sudarsan et al., 2005). On the other hand, the addition of 500 μ M TT (1) represses the expression of β -galactosidase expression to a level similar to that of thiamine. TT has been shown to depend on diphosphorylation to gain activity *in vitro* (Chen et al., 2012). Thus, we anticipate that TT is phosphorylated by the endogenous set of thiamine converting enzymes. Shortening (2) or elongating the alkyl side chain (3 and 4, Figure 1A) results in a loss of repression with increasing deviation from the natural alkyl chain length (Figure 1B). This is most likely due to reduced uptake and/or phosphorylation of these compounds, but could also be because of a poor fit of phosphorylated triazolethiamines to the riboswitch. Previously, it has been shown that the *thi*-box aptamer domain is somewhat adjustable to ligand size through compaction of pyrimidine and phosphate sensor helices (Edwards and Ferre-D'Amare, 2006). However, this low affinity ligand binding [thiamine $K_D = 50 \mu\text{M}$ vs. TPP $K_D = 50 \text{nM}$, (Sudarsan et al., 2005)] is unlikely to efficiently induce subsequent changes in the expression platform (Serganov et al., 2006) and thus reporter gene expression. Interestingly, compound 5, which bears an amino group instead of a hydroxyl residue on the ethyl-alkyl side chain of TT, also represses reporter gene activity. How this compound would be activated through *in vivo* phosphorylation is unclear. However, it is possible that 5 is also phosphorylated in the same way as TT. Experiments using a mutant version of the *thiM* riboswitch (*thiM*-Mu, Figure 1A), which is incapable of binding

to TPP showed that neither of the investigated TT-compounds represses gene expression in these assays. These data underline that repression of gene expression truly relies on riboswitch binding and activation. We next measured IC₅₀-values of these compounds, to evaluate concentration dependent repression of gene expression. TT (1) revealed the strongest effect with an IC₅₀-value of 8.4 μ M (Table 1 and Supplementary Figure 2). Compounds with increasing alkyl-chain length (3 and 4) showed no concentration dependent inhibition curves (Table 1) and, as expected for this bacterial strain, PT did not show an effect on β -galactosidase expression (Table 1, Figure 2A). In order to investigate the dependence of TT activity on endogenous proteins and enzymes involved in thiamine biosynthesis, several strains of the Keio collection were investigated, that all originate from the strain BW25113, and contain deletions of non-essential genes known to be involved in thiamine metabolism of *E. coli* (Scheme 1) (Baba et al., 2006).

Thiamine is synthesized from two precursors, hydroxy-/methyl-/pyrimidine diphosphate (HMP-PP) and hydroxyethylthiazole phosphate (HET-P), which are produced independently and finally joined to form thiamine phosphate (Begley et al., 1999). Further phosphorylation yields the cofactor TPP (Scheme 1). The Keio deletion strains were transformed with the β -galactosidase reporter plasmid containing either the wild type or the mutant TPP-riboswitch and effects of thiamine, TT or PT on reporter gene expression were assayed (Table 2, Figure 3, Supplementary Figures 9, 10). In most strains the addition of thiamine led to a strong decrease in β -galactosidase expression that is comparable to that observed for the wild type strain. As expected, the *thiK* deletion strain (Δ *thiK*) revealed decreased thiamine sensitivity, since this enzyme is required for the phosphorylation of thiamine to TPP. When adding thiamine, *thiI* and *iscS* deletion strains showed slightly increased levels of reporter gene expression indicating a loss of TPP-riboswitch activation (Figure 3A). Investigating the impact of TT (Figure 3B) on TPP-riboswitch dependent reporter gene expression in the deletion strains revealed a prominent increase in reporter gene expression for those strains containing deletions of thiamine transport genes (Scheme 1; *thiP*, *thiQ*, *tbpA*) in comparison to wild type cells (Figure 3B, magenta bars). This may indicate that TT uptake is, at least in part, mediated


Table 1 | IC₅₀-values of thiamine, pyrithiamine and compound 1–5 for inhibition of reporter gene expression in DH5 α Z1 cells are listed.

Compound	IC ₅₀ (μ M) (DH5 α Z1)	95% Confidence interval
Thiamine	0.015	0.012–0.019
1 (TT)	8.4	4.6–15.6
2	43.3	13.4–140
3	n.a.	n.a.
4	n.a.	n.a.
5	22.9	12.1–43.4
PT	n.a.	n.a.

Means and 95% confidence intervals of at least three independent experiments, measured in duplicates, are shown. n.a., not applicable; where data points could not be fitted to a dose-response curve. TT, Triazolethiamine; PT, Pyrithiamine.

by these ABC-transporters. However in the presence of **PT**, reporter gene expression in these strains was scarcely increased in comparison to the wild type strain (**Figure 3C**). This suggests that **PT**, just like thiamine (**Figure 3A**), must use other routes of cell entry apart from the thiamine-specific active transporter. Possibly, its positive charge supports passive diffusion processes, as believed to occur for thiamine as well (Webb et al., 1998). Besides the dependence on transport proteins, differential effects on several other deletion strains involved in phosphate-transfer or thiazole formation were observed (**Figure 3**, green and gray bars, respectively). Assaying $\Delta thiK$, $\Delta thiD$, and $\Delta thiM$ strains, whose deleted genes are involved in phosphate transfer

reactions, revealed increased β -galactosidase expression in the presence of **TT** compared to wild type, whereas $\Delta thiE$ showed decreased β -galactosidase expression (**Figure 3B**). As the deletion of *thiE* renders the bacteria incapable of synthesizing any TPP from its HMP and HET precursors (**Scheme 1**), the reporter gene expression is mainly influenced by external thiamine or compound supply. Interestingly, the prominent decrease of β -galactosidase expression in the presence of **TT** is comparable to that found for thiamine at the same concentration in the wild type strain BW25113 (**Figure 3A**, black bars). In $\Delta thiE$ cells, lower intracellular concentrations of thiamine compete for any possible enzymatic **TT**-activation mechanism (such as



phosphorylation or take-up). Therefore, the same amounts of TT lead to a stronger decrease of reporter gene expression in these cells in comparison to other deletion strains (Figure 3B). In the presence of TT, *thiC* and *iscS* deletion strains which are involved in formation of the thiazole moiety of thiamine revealed increased β -galactosidase expression (Figure 3B). At the same time β -galactosidase expression in *thiL*, *thiG*, *thiH* and *sufS* deletion strains remained unaffected or was slightly decreased in the presence of PT (Figure 3C). Finally, PT-treated *E. coli* strains containing deletions of the genes *thiD* and *thiC*, which are involved in the biosynthesis of the pyrimidine-containing precursor HMP-PP (Scheme 1), show an increase in reporter gene expression (Figure 3C). The same is observed for the Δ *thiE* strain. The deletion of *iscS* renders bacteria less susceptible to thiamine, TT or PT (Supplementary Figure 3). However, it should be noted that the Δ *iscS* strain showed slower growth than any of the other deletion strains investigated. This reduced fitness may be due to the fact that ThiI and IscS are not only involved in thiamine metabolism but also needed in the 4-thiouridine biosynthetic pathway (Scheme 1). Interestingly, deleting the thiamine kinase ThiK abolished thiamine and TT dependent inhibition of reporter gene expression (Figures 3A,B). This finding underlines the hypothesis that TT is only active *in vivo* in its phosphorylated form. Due to its close structural similarity to the natural substrate thiamine, TT is likely to be recognized and phosphorylated by endogenous bacterial enzymes such as ThiK. Using ITC measurements Chen and colleagues revealed that the chemically synthesized, diphosphorylated form of TT,


named triazolethiamine pyrophosphate (TTPP), binds to the *thiM* (Figure 4A) riboswitch with an affinity of 370 nM, whereas for the natural ligand TPP a K_D of 8 nM was measured (Chen et al., 2012). Additionally, *in vitro* translation assays proved that binding of TTPP induces a change in riboswitch secondary structure to sequester the SD-sequence and inhibits efficient protein translation (Chen et al., 2012). Therefore, endogenous diphosphorylation of TT would generate a very potent TPP-riboswitch activator. Hence, we were intrigued to find that reporter gene expression for the *thiK* deletion strain is dramatically decreased in the presence of PT (Figure 3C). This suggests that PT either does not need to be activated by pyrophosphorylation through

Table 2 | IC₅₀-values of thiamine for inhibition of reporter gene expression in *E. coli* containing the indicated deletion of a non-essential thiamine biosynthesis gene are shown as mean and 95% confidence interval of at least three independent experiments, measured in duplicates.

Strain	IC ₅₀ (μ M)	95% Confidence Interval
BW25113 (wt)	0.032	0.022–0.047
Δ <i>thiP</i>	0.93	0.74–1.2
Δ <i>thiQ</i>	1.5	1.0–2.1
Δ <i>tbpA</i>	1.3	0.97–1.7
Δ <i>thiL</i>	0.027	0.018–0.039
Δ <i>thiK</i>	n.a.	n.a.
Δ <i>sufS</i>	0.025	0.018–0.036
Δ <i>thiD</i>	0.019	0.012–0.030
Δ <i>thiM</i>	0.016	0.010–0.026
Δ <i>iscS</i>	n.a.	n.a.
Δ <i>thiH</i>	0.028	0.021–0.037
Δ <i>thiS</i>	0.033	0.018–0.061
Δ <i>thiF</i>	0.009	0.0062–0.013
Δ <i>thiE</i>	0.071	0.045–0.11
Δ <i>thiC</i>	0.023	0.015–0.035
Δ <i>thiG</i>	0.0091	0.0054–0.015

n.a., not applicable, where data points could not be fitted to a dose-response curve.

Respective curves are shown in Supplementary Figures 9, 10.



ThiK, that it can also be phosphorylated by other endogenous enzymes like ThiE or that it is converted into TPP through salvage pathways.

In order to bypass ThiK-dependency of TT activity in *E. coli*, triazolethiamine derivatives that were intended to be independent of endogenous phosphorylation were synthesized and their impact on riboswitch regulation of β -galactosidase expression was studied. These compounds contain a number of phosphate mimics (sulfate, sulfonamide, and sulfone) and metal-chelating groups (dicarbonyl compounds) that might be expected to interact with phosphate-binding metal ions in proteins (Supplementary Figures 4–8). A “click” chemistry approach was used to generate sulfone **7** and the methotrexate- or folate- like compounds **15** and **16** (Figure 4A). Amide couplings were performed to obtain the sulfonamides **8** and **9** and also to obtain the pyruvate, malonate and salicylic acid containing compounds **10**, **11**, and **12**, respectively (Figure 4A) (Erixon et al., 2008). Additionally, a Michael addition was used to generate the sulfones **13** and **14**. With respect to the phosphate mimics, the sulfonamide **13** and the sulfone **14** that are connected to a benzyl group scarcely activate the *thiM* riboswitch in DH5 α Z1 cells. The sulfonamide **8** had a similar effect on β -galactosidase expression in the thiamine auxotroph strain. Interestingly, compounds containing the planar benzene ring are not as active in *thiM* riboswitch activation, even though it is imaginable that this aromatic ring could tighten compound-riboswitch interaction by π - π -stacking with RNA nucleobases (Edwards and Ferre-D’Amare, 2006;

Thore et al., 2006). Compounds **15** and **16** were found to be the most potent repressors of reporter gene expression (Figure 4B, ~ 0.5 fold β -galactosidase expression). Compounds of this group have enough flexibility in principle to interact with two metal ions via their ester or carboxyl groups. Interestingly, these molecules activate the *thiM* riboswitch extensively, despite their relatively large size. The carboxylate ethyl esters like **15** are likely to enter the cell passively and are probably hydrolyzed by esterases resulting in **16**. To investigate whether the most potent phosphate mimics exhibit a ThiK independent reduction in reporter gene expression, compounds **7** and **16** were chosen and assayed in the thiamine kinase deficient *E. coli* strain (Figure 5A). In contrast to addition of thiamine or TT, which do not reduce reporter gene expression, the addition of the same amount of **7** and **16** induces a decrease of β -galactosidase expression to approximately




FIGURE 4 | Chemical formulas of triazole thiamine analogs with phosphate mimicking groups (A). Relative β -galactosidase expression in the presence of thiamine or compound (500 μ M) in DH5 α Z1 cells is shown (B).




FIGURE 5 | Effect of thiamine, TT, PT and phosphate mimics **7** and **16** on β -galactosidase expression in *E. coli* strains Δ thiK (A), BW25113 (B), and Δ thiP (C). Minimal medium (refer to experimental procedures for details) was supplemented with 500 μ M of analyte. Compound **16** was dissolved to a stock solution of 100mM in 100% DMSO, assay controls also contained a final DMSO concentration of 1%.

0.25 fold (**Figure 5A**). However, $\Delta thiK$ strains transformed by the mutated *thiM* riboswitch construct do not or only slightly respond to thiamine or compound addition (Supplementary Figure 11A). In wild type *E. coli* strain BW25113 thiamine has the strongest influence on β -galactosidase repression. **TT** and compound **7** are almost as efficient in inhibiting gene expression whereas **PT** and compound **16** show somewhat weaker effects (**Figure 5B**). Again no significant influence on reporter gene expression in cells containing the mutant *thiM* riboswitch was observed (Supplementary Figure 11C). Investigating the deletion of one subunit of the active thiamine transporter (ThiP) revealed increased levels of reporter protein for **TT**, **7**, and **16**, indicating that these molecules are at least in part taken-up via this mechanism (**Figure 5C**). The investigation of the mutant riboswitch construct *thiM*-Mu proves these compound effects to be specific for *thiM* riboswitch activation, as the mutated RNA does not alter protein expression (Supplementary Figure 11).

DISCUSSION

TPP riboswitch are interesting and attractive target structures for developing antibacterial compounds. They form the most extensive riboswitch class with representatives found in bacteria, archaea and plants (Kubodera et al., 2003; Sudarsan et al., 2003; Bocobza et al., 2007). Here we show that replacing the thiazole heterocycle with 1,2,3-triazole is a valuable strategy to generate thiamine analogs that interact with TPP riboswitches and, thus, induce repression of gene expression. These findings are in line with the observations of Chen et al. recently reporting that the diphosphorylated triazothiamine indeed interacts with the *thiM* riboswitch *in vitro*, albeit with decreased affinity (Chen et al., 2012). We went a significant step further and demonstrated that these compounds are effective in *E. coli* and, more importantly, that their activity depends on proteins involved in the metabolic pathways of thiamine uptake and synthesis.

The data shown above indicate that activity of **TT** depends on active uptake and endogenous phosphorylation by ThiK. Surprisingly, **PT** was found to be independent of ThiK and transporters. Therefore, **PT** is either phosphorylated by other kinases apart from ThiK or its phosphorylation, against all previous hypotheses, is not necessary for *in vivo* activity (Edwards and Ferre-D'Amare, 2006; Warner et al., 2014). Unphosphorylated **PT** is known to have a more than 1000 fold lower affinity for the TPP-riboswitch, whereas diphosphorylation results in a K_D that is only 3fold higher than that of TPP (Sudarsan et al., 2005). Although **PT** was shown to bind to the pyrimidine sensor helix of the aptamer domain by one hydrogen bond and the stacking of the aminopyrimidine ring, crystal structures revealed that two hydrogen bonds have been lost in comparison to TMP or TPP interactions. The pyrophosphate sensor helix is largely disordered and it is questionable whether this riboswitch conformation allows gene regulation upon **PT** binding (Edwards and Ferre-D'Amare, 2006). Furthermore, the increase in *thiM*-riboswitch dependent reporter gene expression in the presence of **PT** in $\Delta thiD$ or $\Delta thiC$ strains suggests the existence of a **PT**-detoxification mechanism in *E. coli* as proposed for *B. subtilis* by Sudarsan et al. (2005). If indeed there was a thiaminase-II-like enzyme present in *E. coli*, HMP could be recovered from **PT**

and used for the generation of thiamine through combination with *de novo* synthesized thiazole (**Scheme 1**). This **PT** recycling would lead to a decrease of its intracellular concentration, hence a diminished influence on the TPP-riboswitch-dependent gene expression. At the same time, the newly generated TPP is likely to be used immediately for its biological purpose without accumulation and hence a need for riboswitch-mediated down regulation of gene expression. Even though a TenA-like protein has not yet been identified in *E. coli*, *in silico* predictions indicate that ThiC may represent a functional equivalent of TenA (Morett et al., 2003).

We further show that equipping **TT** with metal-chelating groups results in ThiK- and transport-independent activity. This indirectly proves that **TT** requires phosphorylation and indicates a possible route in riboswitch activator design which circumvents the necessity of endogenous proteins and renders riboswitch inhibitors independent of bacterial pathways. The most promising phosphate-mimics were tested for their dependence on ThiK or active thiamine transport mechanisms. Whereas activity of compound **16** appears to be ThiK-independent, its transport is strongly influenced by ABC transporter integrity. At the same time, effects of compound **16** in wild type cells are not as prominent as those observed for **TT**. Compound **7** however, does not only reveal strong reporter gene repression, but this effect is even independent of ThiK or ThiP presence. Hence, future compound designs should take characteristics of compound **7** into account in order to provide novel chemical entities whose *in vivo* activity does not depend on the individual enzyme repertoire of the target cell. Also, investigating potential routes of metabolism of phosphate-mimicking compounds may unveil further information and support the generation of even more precise riboswitch-regulating compounds.

The distinct efficiencies of **TT**-derivatives and **PT** in repression of gene expression, further point at possible secondary effects of the employed metabolite analogs. These side effects may rely on interactions with enzymes involved in thiamine pyrophosphate metabolic pathways. Indeed it has been shown that triazole-derivatives of thiamine pyrophosphate inhibit enzymes involved in the biosynthesis of TPP *in vitro* (Erixon and Dabalos, 2007).

Others have shown that analogs with a positively charged middle ring bind much better to the *thiM* riboswitch than ones with a neutral ring at this position (Chen et al., 2012). It was suggested that this is due to the electron drawing effect of the middle ring, which enhances the binding of the pyrimidine moiety to the cognate part of the aptamer domain. The reporter gene assay presented herein however suggests that *in vitro* binding does not always necessarily correlate with *in vivo* riboswitch regulation. Even though **TT** does not contain a charged middle ring, it induces reporter gene repression more efficiently than **PT**. Moreover, our study shows that riboswitch regulation by artificial activators is dependent on the individual set of metabolic enzymes present in the organism.

Since **TT** (**1**) is active in *E. coli* and reveals superior activity when compared to pyrithiamine it represents a very promising starting point for developing novel antibacterial compounds that target TPP-riboswitches. Furthermore, our study shows that even

being very similar from a structural point of view, TT and PT engage diverse endogenous mechanisms to attain activity. These findings are of importance for the design and understanding of compounds that require intracellular activation to achieve an effective repression of gene expression.

ACKNOWLEDGMENTS

We thank the groups of Hans-Georg Sahl and Gabriele Bierbaum (University of Bonn) for helpful comments. This study was supported by funds from the German Research Council (DFG Ma 3442/2-1 and 3442/2-2).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://www.frontiersin.org/journal/10.3389/fchem.2014.00053/abstract>

REFERENCES

- Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., et al. (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* 2:2006.0008. doi: 10.1038/msb4100050
- Begley, T. P., Downs, D. M., Ealick, S. E., McLafferty, F. W., Van Loon, A. P., Taylor, S., et al. (1999). Thiamin biosynthesis in prokaryotes. *Arch. Microbiol.* 171, 293–300. doi: 10.1007/s002030050713
- Blount, K. F., and Breaker, R. R. (2006). Riboswitches as antibacterial drug targets. *Nat. Biotechnol.* 24, 1558–1564. doi: 10.1038/nbt1268
- Bocobza, S., Adato, A., Mandel, T., Shapira, M., Nudler, E., and Aharoni, A. (2007). Riboswitch-dependent gene regulation and its evolution in the plant kingdom. *Genes Dev.* 21, 2874–2879. doi: 10.1101/gad.443907
- Breaker, R. R. (2011). Prospects for riboswitch discovery and analysis. *Mol. Cell* 43, 867–879. doi: 10.1016/j.molcel.2011.08.024
- Chen, L., Cressina, E., Dixon, N., Erixon, K., Agyei-Owusu, K., Micklefield, J., et al. (2012). Probing riboswitch-ligand interactions using thiamine pyrophosphate analogues. *Org. Biomol. Chem.* 10, 5924–5931. doi: 10.1039/c2ob07116a
- Diederich, L., Rasmussen, L. J., and Messer, W. (1992). New cloning vectors for integration in the lambda attachment site attB of the *Escherichia coli* chromosome. *Plasmid* 28, 14–24. doi: 10.1016/0147-619X(92)90032-6
- Edwards, T. E., and Ferre-D'Amare, A. R. (2006). Crystal structures of the thi-box riboswitch bound to thiamine pyrophosphate analogs reveal adaptive RNA-small molecule recognition. *Structure* 14, 1459–1468. doi: 10.1016/j.str.2006.07.008
- Erixon, K. M., and Dabalos, C. L. (2007). Inhibition of pyruvate decarboxylase from *Z. mobilis* by novel analogues of thiamine pyrophosphate: investigating pyrophosphate mimics. *Chem. Commun. (Camb)*, 960–962. doi: 10.1039/b615861g
- Erixon, K. M., Dabalos, C. L., and Leeper, F. J. (2008). Synthesis and biological evaluation of pyrophosphate mimics of thiamine pyrophosphate based on a triazole scaffold. *Org. Biomol. Chem.* 6, 3561–3572. doi: 10.1039/b806580b
- Iwashima, A., Wakabayashi, Y., and Nose, Y. (1976). Formation of pyrithiamine pyrophosphate in brain-tissue. *J. Biochem.* 79, 845–847.
- Kubodera, T., Watanabe, M., Yoshiuchi, K., Yamashita, N., Nishimura, A., Nakai, S., et al. (2003). Thiamine-regulated gene expression of *Aspergillus oryzae* thiA requires splicing of the intron containing a riboswitch-like domain in the 5'-UTR. *FEBS Lett.* 555, 516–520. doi: 10.1016/S0014-5793(03)01335-8
- Lünse, C. E., Schüller, A., and Mayer, G. (2014). The promise of riboswitches as potential antibacterial drug targets. *Int. J. Med. Microbiol.* 304, 79–92. doi: 10.1016/j.ijmm.2013.09.002
- Lutz, R., and Bujard, H. (1997). Independent and tight regulation of transcriptional units in *Escherichia coli* via the LacR/O, the TetR/O and AraC/I1-12 regulatory elements. *Nucleic Acids Res.* 25, 1203–1210. doi: 10.1093/nar/25.6.1203
- Mayer, G., Raddatz, M. S., Grunwald, J. D., and Famulok, M. (2007). RNA ligands that distinguish metabolite-induced conformations in the TPP riboswitch. *Angew. Chem. Int. Ed. Engl.* 46, 557–560. doi: 10.1002/anie.200603166
- Miranda-Rios, J., Navarro, M., and Soberon, M. (2001). A conserved RNA structure (thi box) is involved in regulation of thiamin biosynthetic gene expression in bacteria. *Proc. Nat. Acad. Sci. U.S.A.* 98, 9736–9741. doi: 10.1073/pnas.161168098
- Morett, E., Korb, J. O., Rajan, E., Saab-Rincon, G., Olvera, L., Olvera, M., et al. (2003). Systematic discovery of analogous enzymes in thiamin biosynthesis. *Nat. Biotechnol.* 21, 790–795. doi: 10.1038/nbt834
- Rentmeister, A., Mayer, G., Kuhn, N., and Famulok, M. (2007). Conformational changes in the expression domain of the *Escherichia coli* thiM riboswitch. *Nucleic Acids Res.* 35, 3713–3722. doi: 10.1093/nar/gkm300
- Rodionov, D. A., Vitreschak, A. G., Mironov, A. A., and Gelfand, M. S. (2002). Comparative genomics of thiamin biosynthesis in prokaryotes. New genes and regulatory mechanisms. *J. Biol. Chem.* 277, 48949–48959. doi: 10.1074/jbc.M208965200
- Serganov, A., Polonskaia, A., Phan, A. T., Breaker, R. R., and Patel, D. J. (2006). Structural basis for gene regulation by a thiamine pyrophosphate-sensing riboswitch. *Nature* 441, 1167–1171. doi: 10.1038/nature04740
- Simons, R. W., Houman, F., and Kleckner, N. (1987). Improved single and multi-copy lac-based cloning vectors for protein and operon fusions. *Gene* 53, 85–96. doi: 10.1016/0378-1119(87)90095-3
- Sudarsan, N., Barrick, J. E., and Breaker, R. R. (2003). Metabolite-binding RNA domains are present in the genes of eukaryotes. *RNA* 9, 644–647. doi: 10.1261/rna.5090103
- Sudarsan, N., Cohen-Chalamish, S., Nakamura, S., Emilsson, G. M., and Breaker, R. R. (2005). Thiamine pyrophosphate riboswitches are targets for the antimicrobial compound pyrithiamine. *Chem. Biol.* 12, 1325–1335. doi: 10.1016/j.chembiol.2005.10.007
- Thore, S., Leibundgut, M., and Ban, N. (2006). Structure of the eukaryotic thiamine pyrophosphate riboswitch with its regulatory ligand. *Science* 312, 1208–1211. doi: 10.1126/science.1128451
- Warner, K. D., Homan, P., Weeks, K. M., Smith, A. G., Abell, C., and Ferré-D'Amare, A. R. (2014). Validating fragment-based drug discovery for biological RNAs: Lead fragments bind and remodel the TPP riboswitch specifically. *Chem. Biol.* 21, 591–595. doi: 10.1016/j.chembiol.2014.03.007
- Webb, E., Claas, K., and Downs, D. (1998). thiBPQ encodes an ABC transporter required for transport of thiamine and thiamine pyrophosphate in *Salmonella typhimurium*. *J. Biol. Chem.* 273, 8946–8950. doi: 10.1074/jbc.273.15.8946
- Winkler, W., Nahvi, A., and Breaker, R. R. (2002). Thiamine derivatives bind messenger RNAs directly to regulate bacterial gene expression. *Nature* 419, 952–956. doi: 10.1038/nature01145

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 09 May 2014; accepted: 01 July 2014; published online: 28 July 2014.

Citation: Lünse CE, Scott FJ, Suckling CJ and Mayer G (2014) Novel TPP-riboswitch activators bypass metabolic enzyme dependency. *Front. Chem.* 2:53. doi: 10.3389/fchem.2014.00053

This article was submitted to *Chemical Biology*, a section of the journal *Frontiers in Chemistry*.

Copyright © 2014 Lünse, Scott, Suckling and Mayer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.