

Original citation:

Schlichting, Marco (2017) Euler class groups and the homology of elementary and special linear groups. *Advances in Mathematics*, 320 . pp. 1-81.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/91210>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

EULER CLASS GROUPS AND THE HOMOLOGY OF ELEMENTARY AND SPECIAL LINEAR GROUPS

MARCO SCHLICHTING

ABSTRACT. We improve homology stability ranges for elementary and special linear groups over rings with many units. Our result implies stability for unstable Quillen K -groups and proves a conjecture of Bass. For commutative local rings with infinite residue fields, we show that the obstruction to further stability is given by Milnor-Witt K -theory. As an application we construct Euler classes of projective modules with values in the cohomology of the Milnor-Witt K -theory sheaf. For d -dimensional commutative noetherian rings with infinite residue fields we show that the vanishing of the Euler class is necessary and sufficient for an oriented projective module P of rank d to split off a rank 1 free direct summand. Along the way we obtain a new presentation of Milnor-Witt K -theory and of symplectic K_2 simplifying the classical Matsumoto-Moore presentation.

CONTENTS

1. Introduction	1
2. The homology of affine groups	5
3. Stability in homology and K -theory	10
4. Milnor-Witt K -theory	17
5. The obstruction to further stability	23
5.1. Multiplicative properties of the spectral sequence	24
5.2. Presentation and decomposability	28
5.3. The Steinberg relation and $H_2(SL_2A)$	34
5.4. Centrality of $[-1, 1]$ and $H_n(SL_nA, SL_{n-1}A)$	41
5.5. Prestability	45
6. Euler class groups	47
Appendix A. The affine B.G.-property for the Zariski topology	55
References	64

1. INTRODUCTION

The purpose of this paper is to improve stability ranges in homology and algebraic K -theory of elementary and special linear groups, and to apply these results to construct obstruction classes for projective modules to split off a free direct summand.

Our first result concerns a conjecture of Bass [Bas73, Conjecture XVI on p. 43]. In *loc. cit.* he conjectured that for a commutative noetherian ring A whose maximal

Date: August 10, 2017.

ideal spectrum has dimension d the canonical maps

$$\pi_i BGL_{n-1}^+(A) \rightarrow \pi_i BGL_n^+(A)$$

are surjective for $n \geq d + i + 1$ and bijective for $n \geq d + i + 2$. Here, for a connected space X , we denote by X^+ Quillen's plus-construction with respect to the maximal perfect subgroup of $\pi_1 X$, and we write $BGL_n^+(A)$ for $BGL_n(A)^+$. In this generality, there are counterexamples to Bass' conjecture; see [vdK76, §8]. The best general positive results to date concerning the conjecture are due to van der Kallen [vdK80] and Suslin [Sus82]. They prove that the maps are surjective for $n - 1 \geq \max(2i, \text{sr}(A) + i - 1)$ and bijective for $n - 1 \geq \max(2i, \text{sr}(A) + i)$ where $\text{sr}(A)$ denotes the stable rank of A [Vas71]. Here A need not be commutative nor noetherian.

In this paper we prove Bass' conjecture for rings with many units. Recall [NS89] that a ring A (always associative with unit) has many units if for every integer $n \geq 1$ there is a family of n *central elements* of A such that the sum of each non-empty subfamily is a unit. Examples of rings with many units are infinite fields, commutative local rings with infinite residue field and algebras over a ring with many units. Here is our first main result.

Theorem 1.1 (Theorem 3.10). *Let A be a ring with many units. Then the natural homomorphism*

$$\pi_i BGL_{n-1}^+(A) \rightarrow \pi_i BGL_n^+(A)$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

The ring A in Theorem 1.1 is not assumed to be commutative. If A is commutative noetherian with maximal ideal spectrum of dimension d then $\text{sr}(A) \leq d + 1$ [Bas64, Theorem 11.1]. So, our theorem proves Bass' conjecture in case A has many units. If A is commutative local with infinite residue field then $\text{sr}(A) = 1$ and the theorem admits the following refinement which shows that the stability range in Theorem 1.1 is sharp in many cases. Denote by $K_n^{MW}(A)$ the n -th Milnor-Witt K -theory of A [Mor12, Definition 3.1] which makes sense for any commutative ring A ; see Definition 4.10.

Theorem 1.2 (Theorem 5.38). *Let A be a commutative local ring with infinite residue field. Then the natural homomorphism*

$$\pi_i BGL_{n-1}^+(A) \rightarrow \pi_i BGL_n^+(A)$$

is an isomorphism for $n \geq i + 2$ and surjective for $n \geq i + 1$. Moreover, there is an exact sequence for $n \geq 2$

$$\pi_n BGL_{n-1}^+(A) \rightarrow \pi_n BGL_n^+(A) \rightarrow K_n^{MW}(A) \rightarrow \pi_{n-1} BGL_{n-1}^+(A) \rightarrow \pi_{n-1} BGL_n^+(A).$$

Theorem 1.1 follows from the following homology stability result for elementary linear groups. Recall [Bas64, §1] that the group of elementary $r \times r$ -matrices of a ring A is the subgroup $E_r(A)$ of $GL_r(A)$ generated by the elementary matrices $e_{i,j}(a) = 1 + a \cdot e_i e_j^T$, $a \in A$ where $e_i \in A^r$ is the i -th standard column basis vector.

Theorem 1.3 (Theorem 3.9). *Let A be a ring with many units. Then the natural homomorphism*

$$H_i(E_{n-1}(A), \mathbb{Z}) \rightarrow H_i(E_n(A), \mathbb{Z})$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

For a division ring A with infinite center, Theorem 1.3 proves a conjecture of Sah [Sah89, 2.6 Conjecture]. From Theorem 1.3 one easily deduces the following homology stability result for the special linear groups of commutative rings.

Theorem 1.4 (Theorem 3.12). *Let A be a commutative ring with many units. Then the natural homomorphism*

$$H_i(SL_{n-1}(A), \mathbb{Z}) \rightarrow H_i(SL_n(A), \mathbb{Z})$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

When A is a commutative local ring with infinite residue field Theorem 1.4 says that $H_i(SL_n(A), SL_{n-1}(A)) = 0$ for $i < n$. The following theorem gives an explicit presentation of these groups for $i = n$.

Theorem 1.5 (Theorem 5.37). *Let A be a commutative local ring with infinite residue field. Then for all $n \geq 2$ we have*

$$H_n(SL_n(A), SL_{n-1}(A)) \cong K_n^{MW}(A).$$

Moreover, for n even, the map $H_n(SL_n(A)) \rightarrow H_n(SL_n(A), SL_{n-1}(A))$ is surjective. In particular, the map $H_i(SL_{n-1}(A)) \rightarrow H_i(SL_n(A))$ is an isomorphism for $i \leq n - 2$ and surjective (bijective) for $i = n - 1$ and n odd (n even).

Theorems 1.4 and 1.5 generalize a result of Hutchinson and Tao [HT10] who proved them for fields of characteristic zero, though for n odd, the identification of the relative homology with Milnor-Witt K -theory is only implicit in their work. Contrary to [HT10], our proof is independent of the characteristic of the residue field, works for local rings other than fields and does not use the solution of the Milnor conjecture on quadratic forms. In Theorem 5.39 we also give explicit computations of the kernel and cokernel of the stabilization map in homology at the edge of stabilization recovering and generalizing the remaining results of [HT10]. This, however, requires the solution of the Milnor conjecture. Theorem 1.5 answers questions raised in [BM99].

Our proof of Theorem 1.5 uses a new presentation of the Milnor-Witt K -groups $K_n^{MW}(A)$ for $n \geq 2$. Denote by $\mathbb{Z}[A^*]$ the group ring of the group of units A^* in A , and $I[A^*]$ the augmentation ideal. For $a \in A^*$ denote by $\langle a \rangle \in \mathbb{Z}[A^*]$ the corresponding element in the group ring, and by $[a] \in I[A^*]$ the element $\langle a \rangle - 1$. We define the graded ring $\hat{K}^{MW}(A)$ as the graded $\mathbb{Z}[A^*]$ -algebra generated in degree 1 by $I[A^*]$ modulo the two sided ideal generated by the Steinberg relations $[a][1 - a]$ for all $a, 1 - a \in A^*$; see Definition 4.2.

Theorem 1.6 (Theorem 4.18). *Let A be a commutative local ring. If A is not a field assume that the cardinality of its residue field is at least 4. Then the natural map of graded rings $\hat{K}^{MW}(A) \rightarrow K^{MW}(A)$ induces an isomorphism $\hat{K}_n^{MW}(A) \cong K_n^{MW}(A)$ for $n \geq 2$.*

In particular, for a local ring A with infinite residue field, the Schur multiplier $H_2(SL_2(A))$ has a presentation as the quotient of $I[A^*] \otimes_{A^*} I[A^*]$ by the Steinberg relations (Theorem 5.27); compare [Moo68, Theorem 9.2], [Mat69, Corollaire 5.11], [vdK77, Theorem 3.4].

Theorems 1.4 and 1.5 are the SL_n -analogs of a result of Nesterenko and Suslin [NS89]. They proved that Theorems 1.4 and 1.5 hold when $SL_n(A)$ and $K^{MW}(A)$

are replaced with $GL_n(A)$ and Milnor K-theory $K^M(A)$. Suslin and Nesterenko's proof rests on the computation of the homology of affine groups [NS89, Theorem 1.11] which is false if one simply replaces $GL_n(A)$ with $SL_n(A)$. Our innovation is the correct replacement of [NS89, Theorem 1.11] in the context of $SL_n(A)$ and of groups related to $E_n(A)$. This is done in Section 2 whose main result is Theorem 2.5 and its Corollary 2.6. With our new presentation of Milnor-Witt K-theory in Section 4, Sections 3 and 5 more or less follow the treatment in [NS89].

The importance of homology stability and the computation of the obstruction to further stability in Theorem 1.5 lies in the following application. Let R be a commutative noetherian ring of dimension n all of whose residue fields are infinite. Let P be an oriented rank n projective R -module. In §6, we define an "Euler class"

$$e(P) \in H_{Zar}^n(R, \mathcal{K}_n^{MW})$$

such that $e(P) = 0$ if P splits off a free direct summand of rank 1. Here, \mathcal{K}_n^{MW} denotes the Zariski sheaf associated with the presheaf $A \mapsto K_n^{MW}(A)$, and H_{Zar}^n denotes Zariski cohomology. We prove the following.

Theorem 1.7 (Theorem 6.18). *Let R be a commutative noetherian ring of dimension $n \geq 2$. Assume that all residue fields of R are infinite. Let P be an oriented rank n projective R -module. Then*

$$P \cong Q \oplus R \Leftrightarrow e(P) = 0 \in H_{Zar}^n(R, \mathcal{K}_n^{MW}).$$

The theorem also holds if we replace Zariski cohomology with Nisnevich cohomology (Theorem 6.24).

If R has dimension n and is of finite type over an algebraically closed field k , then the canonical map $\mathcal{K}_n^{MW} \rightarrow \mathcal{K}_n^M$ of sheaves on X is an isomorphism. In particular, if R has dimension n and is smooth over an algebraically closed field, then $H_{Zar}^n(R, \mathcal{K}_n^{MW}) = H_{Zar}^n(R, \mathcal{K}_n^M)$ is isomorphic to the Chow group of codimension n cycles on $X = \text{Spec } R$, by [Ker09, Theorem 7.5], and we recover a result of Murthy [Mur94]. If R is smooth over a field of characteristic not 2 (which is not assumed algebraically closed) then $H_{Zar}^n(R, \mathcal{K}_n^{MW})$ is isomorphic to the Chow-Witt groups introduced by Barge and Morel [BM00] and studied by Fasel [Fas08].

Theorem 1.7 is a generalization to singular affine varieties of a theorem of Morel [Mor12, Theorem 8.14] who proved the result for R smooth of finite type over a perfect field. Our arguments don't use \mathbb{A}^1 -homotopy theory but they can be used to simplify some proofs in [Mor12]; see proof of Theorem 6.22 and [AHW17].

The proof of Theorem 1.7 relies on Theorem 1.5 and a representability result of vector bundles on noetherian affine schemes (Theorem 6.15) which is of independent interest. There is also a version (Theorem 6.21) of Theorem 1.7 for projective modules with orientation in a line bundle other than R .

Advice. The reader only interested in the results for local rings A and the application to Euler classes in Theorem 1.7 should replace in Sections 2 and 3 the stable rank $\text{sr}(A)$ with 1, the groups $G_n A$ with $GL_n A$, $SG_n(A)$ with $SL_n(A)$, \bar{A}^* with A^* and Lemma 3.1 with Lemma 5.6. This should make for a smoother reading.

Conventions. All rings are associative with unit. The group of units of a ring A is denoted A^* . The stable rank [Vas71] of a ring A is denoted $\text{sr}(A)$. By "space" we mean "simplicial set". Unless otherwise stated, tensor products are

over \mathbb{Z} and homology has coefficients in \mathbb{Z} . For a commutative ring A and integer $n \geq 1$, the group $SL_n(A)$ is the group of $n \times n$ matrices with entries in A and determinant 1. The symbol $SL_0(A)$ will stand for the discrete set (or discrete groupoid) A^* . This has the effect that for all $n \geq 0$ and any $GL_n(A)$ -module M , we have $H_i(SL_n(A), M) = \text{Tor}_i^{GL_n(A)}(\mathbb{Z}[A^*], M)$ where $\mathbb{Z}[A^*]$ is a $GL_n(A)$ -module via the determinant map $GL_n(A) \rightarrow A^*$. Moreover, for all $n \geq 0$, we have a homotopy fibration of classifying spaces $BSL_n(A) \rightarrow BGL_n(A) \rightarrow BA^*$.

We denote by sSets the category of simplicial sets endowed with its standard Kan model structure.

Acknowledgements. Most of the results of this article were found in Spring 2014 while the author was visiting Max-Planck-Institute for Mathematics in Bonn. I would like to thank MPIM Bonn for its hospitality. I also would like to thank Marc Hoyois whose comments lead to the appendix, and Wataru Kai for pointing out an error in a previous version.

2. THE HOMOLOGY OF AFFINE GROUPS

For a group G , we denote by $\mathbb{Z}[G]$ its integral group ring and we write $\langle g \rangle$ for the element in $\mathbb{Z}[G]$ corresponding to $g \in G$. Furthermore, we denote by $\varepsilon : \mathbb{Z}[G] \rightarrow \mathbb{Z} : \langle g \rangle \mapsto 1$ the augmentation ring homomorphism, and by $I[G] = \ker(\varepsilon)$ its kernel, the augmentation ideal. Let G be an abelian group and $s \in \mathbb{Z}[G]$ an element in its group ring. A G -module M is called s -torsion if for every $x \in M$ there is $n \in \mathbb{N}$ such that $s^n x = 0$, or equivalently if $[s^{-1}]M = 0$. The category of s -torsion G -modules is closed under taking subobjects, quotient objects and extensions in the category of all G -modules.

Many of our computations concern the homology $H_i(G, M)$ of a group G with coefficients in a left G -module M . We recall the basic functoriality of this construction [Bro82, §III.8]. Let G, G' be groups, and M, M' be G, G' -modules, respectively. A pair of maps $(\varphi, f) : (G, M) \rightarrow (G', M')$ where $\varphi : G \rightarrow G'$ is a group homomorphism and $f : M \rightarrow M'$ is a homomorphism of abelian groups with $f(gx) = \varphi(g)f(x)$ for all $g \in G$ and $x \in M$ induces a map of homology groups $(\varphi, f)_* : H_*(G, M) \rightarrow H_*(G', M')$. Given two such pairs of maps $(\varphi_0, f_0), (\varphi_1, f_1) : (G, M) \rightarrow (G', M')$. If there is an element $h \in G'$ such that $\varphi_1(g) = h\varphi_0(g)h^{-1}$ and $f_1(x) = hf_0(x)$ for all $g \in G$ and $x \in M$, then the induced maps on homology agree: $(\varphi_0, f_0)_* = (\varphi_1, f_1)_* : H_*(G, M) \rightarrow H_*(G', M')$.

Definition 2.1. Let A be a ring with group of units A^* , and let $m \geq 0$ be an integer. An $S(m)$ -sequence in A is a sequence (a_1, \dots, a_m) of m central¹ elements a_1, \dots, a_m in A all of whose non-empty partial sums are units, that is, for every index set $\emptyset \neq J \subset \{1, \dots, m\}$, we require

$$a_J = \sum_{j \in J} a_j \in A^*.$$

Following [NS89], an $S(m)$ -algebra is a ring which has an $S(m)$ -sequence. A ring with many units is a ring that has an $S(m)$ -sequence for every integer $m \geq 1$. For instance, a commutative local ring with infinite residue field has many units, and any algebra over a ring with many units has many units.

¹In [NS89] centrality of the a_i 's is not required. However, it is used in the proof of [NS89, Theorem 1.11] where A^* is considered a normal subgroup of $GL_n(A)$ via the diagonal embedding.

$S(m)$ -algebras are algebras over the commutative ring

$$S(m) = \mathbb{Z}[X_1, \dots, X_m][\Sigma^{-1}]$$

obtained by localizing the polynomial ring $\mathbb{Z}[X_1, \dots, X_m]$ in the m variables X_1, \dots, X_m at the set of all non-empty partial sums of the variables

$$\Sigma = \{X_J \mid \emptyset \neq J \subset \{1, \dots, m\}\}, \quad \text{where } X_J = \sum_{j \in J} X_j.$$

By construction, $S(m)$ has an $S(m)$ -sequence $X = (X_1, \dots, X_m)$. Sending X_i to a_i makes a ring A with $S(m)$ -sequence (a_1, \dots, a_m) into an $S(m)$ -algebra.

Let A be a ring with $S(m)$ -sequence $a = (a_1, \dots, a_m)$. For an integer $m \geq 1$, write $[1, m]$ for the set $\{1, \dots, m\}$ of integers between 1 and m . We will denote by $s(a) \in \mathbb{Z}[A^*]$ the following element in the integral group ring of the group of units of A :

$$s(a) = - \sum_{\emptyset \neq J \subset [1, m]} (-1)^{|J|} \langle a_J \rangle \in \mathbb{Z}[A^*].$$

Note that the augmentation homomorphism $\varepsilon : \mathbb{Z}[A^*] \rightarrow \mathbb{Z}$ sends $s(a)$ to 1:

$$s(a) \xrightarrow{\varepsilon} - \sum_{\emptyset \neq J \subset [1, m]} (-1)^{|J|} = (-1)^{|\emptyset|} - \sum_{J \subset [1, m]} (-1)^{|J|} = 1 - (1 - 1)^m = 1.$$

More generally, for an integer $t \in \mathbb{Z}$ we will write $s_t(a)$ for the image of $s(a)$ under the ring homomorphism $t : \mathbb{Z}[A^*] \rightarrow \mathbb{Z}[A^*] : \langle x \rangle \mapsto \langle x^t \rangle$, that is,

$$s_t(a) = - \sum_{\emptyset \neq J \subset [1, m]} (-1)^{|J|} \langle (a_J)^t \rangle \in \mathbb{Z}[A^*].$$

When $A = S(m)$ with $S(m)$ -sequence $X = (X_1, \dots, X_m)$, we may write s_m and $s_{m,t}$ for $s(X)$ and $s_t(X)$.

Lemma 2.2. *Let A be a commutative ring with $S(m)$ -sequence $a = (a_1, \dots, a_m)$ and let $k, t \geq 1$ be integers such that $k \cdot t < m$. Then the ring homomorphism $\varphi_k : \mathbb{Z}[A^*] \rightarrow A^{\otimes k}$, induced by $\langle x \rangle \mapsto x \otimes \dots \otimes x$ for $x \in A^*$, sends $s_t(a) \in \mathbb{Z}[A^*]$ to $0 \in A^{\otimes k}$.*

Proof. We first show the case $t = 1$. For a function $\sigma : [1, k] \rightarrow [1, m]$ we write $a^\sigma = a_{\sigma(1)} \otimes \dots \otimes a_{\sigma(k)}$. Note that $a_\emptyset = 0$ and $(a_\emptyset)^{\otimes k} = 0$. In $A^{\otimes k}$ we have

$$\begin{aligned} -s(a) &= \sum_{J \subset [1, m]} (-1)^{|J|} (a_J)^{\otimes k} = \sum_{\substack{J \subset [1, m] \\ \sigma : [1, k] \rightarrow J}} (-1)^{|J|} a^\sigma \\ &= \sum_{\sigma : [1, k] \rightarrow [1, m]} a^\sigma \sum_{\text{Im } \sigma \subset J \subset [1, m]} (-1)^{|J|} = 0 \end{aligned}$$

since for $I \subset [1, m]$ with $I \neq [1, m]$ we have

$$\sum_{I \subset J \subset [1, m]} (-1)^{|J|} = (-1)^{|I|} \sum_{J \subset [1, m] - I} (-1)^{|J|} = (-1)^{|I|} (1 - 1)^{|[1, m] - I|} = 0.$$

This shows that $\varphi_k(s(a)) = 0 \in A^{\otimes k}$ for all $m > k \geq 1$, that is, the case $t = 1$. For general $t \geq 1$, the lemma follows from the case $t = 1$ and the commutative diagram of rings

$$\begin{array}{ccc} \mathbb{Z}[A^*] & \xrightarrow{t} & \mathbb{Z}[A^*] \\ \varphi_{tk} \downarrow & & \downarrow \varphi_k \\ (A^{\otimes t})^{\otimes k} & \xrightarrow{\mu^{\otimes k}} & A^{\otimes k} \end{array}$$

where the top horizontal map is induced by $\langle x \rangle \mapsto \langle x^t \rangle$ and the lower one by the multiplication $\mu : A^{\otimes t} \rightarrow A : x_1 \otimes \cdots \otimes x_t \mapsto x_1 \cdots x_t$. \square

For an integer $k \geq 1$ and an abelian group M , consider the k -th exterior power $\Lambda_{\mathbb{Z}}^k M$ of M over \mathbb{Z} ; see [Bro82, §V.6]. This is the quotient of the abelian group $M^{\otimes k}$ modulo the abelian subgroup generated by elements of the form $x_1 \otimes \cdots \otimes x_k$ where $x_i = x_{i+1}$ for some $i = 1, \dots, k-1$. If M is an A -module, then $\Lambda_{\mathbb{Z}}^k M$ becomes an A^* -module under the diagonal action $a \cdot (x_1 \wedge \cdots \wedge x_k) = ax_1 \wedge \cdots \wedge ax_k$ where $a \in A^*$ and $x_i \in M$.

Corollary 2.3. *Let A be a commutative ring with $S(m)$ -sequence $a = (a_1, \dots, a_m)$. Let M be an A -module. Then for all integers $k, t \geq 1$ with $k \cdot t < m$ the $\mathbb{Z}[A^*]$ -module $\Lambda_{\mathbb{Z}}^k M$ satisfies*

$$s_t(a) \cdot \Lambda_{\mathbb{Z}}^k M = 0.$$

Proof. Denote by $V_k(A)$ the image of the ring homomorphism $\varphi_k : \mathbb{Z}[A^*] \rightarrow A^{\otimes k}$ from Lemma 2.2. This is a subring of $A^{\otimes k}$. The abelian group $M^{\otimes k}$ has a natural $A^{\otimes k}$ -module structure

$$A^{\otimes k} \times M^{\otimes k} \rightarrow M^{\otimes k} : (b_1 \otimes \cdots \otimes b_k, x_1 \otimes \cdots \otimes x_k) \mapsto b_1 x_1 \otimes \cdots \otimes b_k x_k,$$

hence, a natural $V_k(A)$ -module structure, by restriction of scalars. The abelian subgroup of $M^{\otimes k}$ generated by $x_1 \otimes \cdots \otimes x_k$ with $x_i = x_{i+1}$ for some $i = 1, \dots, k-1$ is a $V_k(A)$ -submodule since $V_k(A)$ is generated as an abelian group by elements of the form $b \otimes \cdots \otimes b$ with $b \in A^*$. It follows that the quotient module $\Lambda_{\mathbb{Z}}^k M$ has a natural structure of a $V_k(A)$ -module. Restriction of scalars along the ring homomorphism $\varphi_k : \mathbb{Z}[A^*] \rightarrow V_k(A)$ induces the diagonal A^* -action on $\Lambda^k M$. By Lemma 2.2, the map φ_k sends $s_t(a)$ to zero in $V_k(A)$. The result follows. \square

Let R be a ring and M an R -module. Then the group of units R^* acts on M via abelian group automorphisms. In particular, R^* acts linearly on the integral homology groups $H_q(M, \mathbb{Z})$ of M , by functoriality of group homology. This action makes $H_q(M, \mathbb{Z})$ into a $\mathbb{Z}[R^*]$ -module.

Proposition 2.4. *Let R be a commutative ring with $S(m)$ -sequence $a = (a_1, \dots, a_m)$. Let M be an R -module. Then for all integers $t, q \geq 1$ with $tq < m$ the integral homology groups $H_q(M, \mathbb{Z})$ of M are $s_t(a)$ -torsion, that is, localization at $s_t(a)$ yields*

$$[s_t(a)^{-1}] H_q(M, \mathbb{Z}) = 0.$$

Proof. Recall that the $S(m)$ -sequence $a = (a_1, \dots, a_m)$ makes R into an $S(m)$ -algebra. Restriction of scalars makes M into an $S(m)$ -module and $H_q(M, \mathbb{Z})$ into an $S(m)^*$ -module such that

$$[s_t(X)^{-1}] H_q(M, \mathbb{Z}) = [s_t(a)^{-1}] H_q(M, \mathbb{Z}).$$

So, we can assume $R = S(m)$ and $X = a$.

Consider M as a constant simplicial $S(m)$ -module. Choose a weak equivalence $P_* \rightarrow M$ in the category of simplicial $S(m)$ -modules such that P_* is projective in each degree. For instance, the image under the Dold-Kan correspondence of an $S(m)$ -projective resolution of M will do. The classifying space-functor applied to the map of simplicial abelian groups $P_* \rightarrow M$ yields an $S(m)^*$ -equivariant weak equivalence of simplicial sets $BP_* \rightarrow BM$ and hence $S(m)^*$ -equivariant isomorphisms $H_q(BP_*) \cong H_q(BM)$ of integral homology groups. To the $S(m)^*$ -equivariant simplicial space $s \mapsto BP_s$ is associated a strongly convergent first quadrant spectral sequence of $S(m)^*$ -modules

$$E_{r,s}^1 = H_r(BP_s, \mathbb{Z}) \Rightarrow H_{r+s}(BP_*, \mathbb{Z}) = H_{r+s}(BM, \mathbb{Z})$$

where $d^1 : H_r(P_s) \rightarrow H_r(P_{s-1})$ is the alternating sum of the face maps of the simplicial abelian group $s \mapsto H_r(BP_s)$. Since the ring $S(m)$ is flat over \mathbb{Z} , each P_s is a torsion-free abelian group, and thus, the Pontryagin map $\Lambda_{\mathbb{Z}}^r P_s \rightarrow H_r(BP_s)$ is an isomorphism of $S(m)^*$ -modules [Bro82, Theorem V.6.4.(ii)]. By Corollary 2.3, the $S(m)^*$ -module $\Lambda_{\mathbb{Z}}^r P_s$ is $s_t(X)$ -torsion for all $tr < m$. Since $E_{0,s}^2 = 0$ for $s \geq 1$, it follows from the spectral sequence that $H_q(M)$ is $s_t(X)$ -torsion whenever $1 \leq tq < m$. \square

The rest of the section is devoted to proving Theorem 2.5 below which is an analog of [NS89, Theorem 1.11] for groups related to SL_n and E_n . Let A be a ring and $Z(A)$ its center. Let $q \geq 1$ be an integer. The inclusions

$$GL_q(A) \subset GL_{q+1}(A) \subset GL(A) : M \mapsto \begin{pmatrix} M & 0 \\ 0 & 1 \end{pmatrix}$$

define group homomorphisms $\det : GL_q(A) \rightarrow GL(A)^{ab} = K_1(A)$ whose kernel we denote by $SG_q(A)$. If A is an $S(m)$ -algebra, we will need an action of $S(m)^*$ on the integral homology groups of $SG_q(A)$. For that end, let \bar{A}^* be the image in $K_1(A)$ of the map $Z(A)^* \subset GL_1(A) \rightarrow K_1(A)$. We denote by $G_q(A)$ the subgroup of $GL_q(A)$ consisting of those matrices $T \in GL_q(A)$ whose class $\det(T) \in K_1(A)$ lies in the subgroup $\bar{A}^* \subset K_1(A)$. Note that $G_q(A)$ contains all invertible diagonal matrices with entries in $Z(A)$. In particular, the map $\det : GL_q(A) \rightarrow K_1(A)$ restricts to a surjective group homomorphism $\det : G_q(A) \rightarrow \bar{A}^*$, and we have an exact sequence of groups

$$1 \rightarrow SG_q(A) \longrightarrow G_q(A) \xrightarrow{\det} \bar{A}^* \rightarrow 1.$$

We will write $\text{Aff}_{p,q}^G(A)$ and $\text{Aff}_{p,q}^{SG}(A)$ for the following subgroups of $GL_{p+q}(A)$

$$\text{Aff}_{p,q}^G = \begin{pmatrix} G_q(A) & 0 \\ M_{p,q}(A) & 1_p \end{pmatrix} \quad \text{and} \quad \text{Aff}_{p,q}^{SG} = \begin{pmatrix} SG_q(A) & 0 \\ M_{p,q}(A) & 1_p \end{pmatrix}.$$

For any $M \in M_{p,q}(A)$ and $T \in GL_q(A)$, the matrices T and

$$\begin{pmatrix} T & 0 \\ M & 1_p \end{pmatrix} = \begin{pmatrix} 1_q & 0 \\ MT^{-1} & 1_p \end{pmatrix} \begin{pmatrix} T & 0 \\ 0 & 1_p \end{pmatrix}$$

have the same class in $K_1(A)$. It follows that the map $\det : GL_{p+q}(A) \rightarrow K_1(A)$ restricts to a surjective group homomorphism $\text{Aff}_{p,q}^G(A) \rightarrow \bar{A}^*$ with kernel the group $\text{Aff}_{p,q}^{SG}(A)$. Hence, for integers $q \geq 1$, $p \geq 0$ the exact sequence of groups

$$1 \rightarrow \text{Aff}_{p,q}^{SG}(A) \longrightarrow \text{Aff}_{p,q}^G(A) \xrightarrow{\det} \bar{A}^* \rightarrow 1$$

makes the homology groups $H_r(\text{Aff}_{p,q}^{SG}(A))$ into left \bar{A}^* -modules [Bro82, Corollary III.8.2].

For an $S(m)$ -algebra A , we denote by $s_{m,t} \in \mathbb{Z}[\bar{A}^*]$ the image of $s_{m,t} \in \mathbb{Z}[S(m)^*]$ under the ring homomorphism $\mathbb{Z}[S(m)^*] \rightarrow \mathbb{Z}[\bar{A}^*]$ induced by the group homomorphism $S(m)^* \rightarrow Z(A)^* \rightarrow \bar{A}^*$. The following is our analog of [NS89, Theorem 1.11].

Theorem 2.5. *Let A be an $S(m)$ -algebra. Let $t, q \geq 1$ be integers such that q divides t . Then for all integers $p, r \geq 0$ such that $rt < mq$ the inclusion*

$$SG_q(A) \rightarrow \text{Aff}_{p,q}^{SG}(A) : M \mapsto \begin{pmatrix} M & 0 \\ 0 & 1_p \end{pmatrix}$$

induces an isomorphism of \bar{A}^* -modules

$$H_r(SG_q(A)) \cong s_{m,-t}^{-1} H_r(\text{Aff}_{p,q}^{SG}(A)).$$

Proof. A matrix $T \in G_q(A)$ defines an automorphism of the exact sequence of groups

$$(2.1) \quad 0 \rightarrow M_{p,q}(A) \rightarrow \begin{pmatrix} SG_q(A) & 0 \\ M_{p,q}(A) & 1_p \end{pmatrix} \rightarrow SG_q(A) \rightarrow 1$$

through right multiplication by T^{-1} on $M_{p,q}(A)$, through conjugation by $\begin{pmatrix} T & 0 \\ 0 & 1_p \end{pmatrix}$ on the middle term and through conjugation by T on $SG_q(A)$. This defines a left action of the group $G_q(A)$ on the exact sequence and hence an action on the associated Hochschild-Serre spectral sequence

$$(2.2) \quad E_{i,j}^2 = H_i(SG_q A, H_j(M_{p,q} A)) \Rightarrow H_{i+j} \begin{pmatrix} SG_q(A) & 0 \\ M_{p,q}(A) & 1_p \end{pmatrix}$$

which descends to an $G_q/SG_q = \bar{A}^*$ -action via the determinant map $G_q(A) \rightarrow \bar{A}^*$, in view of the basic functoriality of group homology recalled at the beginning of this section. Since the surjection in the exact sequence (2.1) splits, we have $H_i(SG_q, \mathbb{Z}) = E_{i,0}^2 = E_{i,0}^\infty$.

On the homology groups $H_i(SG_q A, H_j(M_{p,q} A))$, the element $T \in G_q(A)$ acts through conjugation on $SG_q(A)$ and right multiplication by T^{-1} on $M_{p,q}(A)$. Since q divides t , we can write $t = q \cdot k$ for some integer $k \geq 1$. For $\bar{a} \in \bar{A}^*$, the element $\langle \bar{a}^{-t} \rangle$ acts on the spectral sequence as the diagonal matrix $T = a^{-k} \cdot 1_q \in G_q$ where $a \in Z(A)^*$ is a lift of $\bar{a} \in \bar{A}^*$. This element acts on the pair $(SG_q A, H_j(M_{p,q} A))$ through conjugation by $a^{-k} \cdot 1_q$ on $SG_q A$ which is the identity map, and through right translation by $T^{-1} = (a^{-k} \cdot 1_q)^{-1} = a^k \cdot 1_q$ on $M_{p,q} A$ which is the action by $\langle a^k \rangle \in \mathbb{Z}[\bar{A}^*]$ induced by the usual left $Z(A)$ -module structure on $M_{p,q} A$. In view of Proposition 2.4 with $R = S(m)$ it follows that for $j \geq 1$ and $kj < m$ we have

$$s_{m,-t}^{-1} H_i(SG_q A, H_j(M_{p,q} A)) = H_i(SG_q A, s_{m,k}^{-1} H_j(M_{p,q} A)) = 0.$$

Moreover,

$$s_{m,-t}^{-1} H_i(SG_q A, H_0(M_{p,q} A)) = H_i(SG_q A, s_{m,k}^{-1} H_0(M_{p,q} A)) = H_i(SG_q A, \mathbb{Z})$$

since $s_{m,k}$ acts through $\varepsilon(s_{m,k}) = 1$ on $H_0(M_{p,q} A) = \mathbb{Z}$. Localizing the spectral sequence (2.2) at $s_{m,-t} \in \mathbb{Z}[\bar{A}^*]$ yields a spectral sequence which satisfies $s_{m,-t}^{-1} E_{i,j}^2 = s_{m,-t}^{-1} E_{i,j}^\infty = 0$ for $tj < mq$ and $j \geq 1$, and $s_{m,-t}^{-1} E_{i,0}^2 = s_{m,-t}^{-1} E_{i,0}^\infty = H_i(SG_q, \mathbb{Z})$. The claim follows. \square

It will be convenient to reinterpret this result in somewhat different notation. To that end, let A be an $S(m)$ -algebra. We introduce the rings Λ and $\Lambda_{m,t}$ as

$$\Lambda = \mathbb{Z}[\bar{A}^*], \quad \Lambda_{m,t} = (s_{m,-t})^{-1} \Lambda.$$

Note that the natural maps of groups $GL_q(A) \rightarrow K_1(A)$ induce ring homomorphisms $\mathbb{Z}[G_q(A)] \rightarrow \Lambda \rightarrow \Lambda_{m,t}$ compatible with the inclusions $G_q(A) \subset G_{q+1}(A)$.

Corollary 2.6. *Let A be an $S(m)$ -algebra. Let $t, q \geq 1$ be integers such that q divides t . Then for all integers $p \geq 0$ the canonical inclusions of groups and rings $SG_q(A) \subset G_q(A) \subset \text{Aff}_{p,q}^G(A)$ and $\mathbb{Z} \subset \Lambda_{m,t}$ induce maps of complexes*

$$\mathbb{Z} \otimes_{SG_q(A)}^L \mathbb{Z} \longrightarrow \Lambda_{m,t} \otimes_{G_q(A)}^L \mathbb{Z} \longrightarrow \Lambda_{m,t} \otimes_{\text{Aff}_{p,q}^G(A)}^L \mathbb{Z}$$

which are isomorphisms on homology groups in degrees $r < mq/t$.

Proof. Recall that for a subgroup $N \subset G$ of a group, we have Shapiro's Lemma

$$\mathbb{Z}[N \backslash G] \otimes_G^L \mathbb{Z} = \mathbb{Z} \otimes_N^L \mathbb{Z}[G] \otimes_G^L \mathbb{Z} = \mathbb{Z} \otimes_N^L \mathbb{Z}$$

since $\mathbb{Z}[N \backslash G] = \mathbb{Z} \otimes_N \mathbb{Z}[G] = \mathbb{Z} \otimes_N^L \mathbb{Z}[G]$ as $\mathbb{Z}[G]$ is a free N -module. If N is normal in G then $G/N = N \backslash G$ is a group and $\mathbb{Z}[G/N] \otimes_G^L \mathbb{Z}$ is a complex of left G/N -modules where G/N acts as left multiplication on $\mathbb{Z}[G/N]$. On homology, the isomorphism

$$H_i(\mathbb{Z}[G/N] \otimes_G^L \mathbb{Z}) \cong H_i(\mathbb{Z} \otimes_N^L \mathbb{Z}) = H_i(N)$$

is an isomorphism of G/N -modules where the action on $H_i(N)$ is the usual conjugation action. Applied to $N = \text{Aff}_{p,q}^{SG}$ and $G = \text{Aff}_{p,q}^G$, we have an isomorphism

$$H_i(\Lambda \otimes_{\text{Aff}_{p,q}^G}^L \mathbb{Z}) \cong H_i(\text{Aff}_{p,q}^{SG})$$

of Λ -modules. Localizing at $s_{m,-t}$ using Theorem 2.5, the result follows. \square

3. STABILITY IN HOMOLOGY AND K -THEORY

Let A be a ring and recall from Section 2 the definition of the groups $G_q(A)$ for $q \geq 1$. We set $G_0(A) = \{1\}$, the one-element group. Let $n \geq r \geq 0$ be integers. We denote by $U_r(A^n) \subset M_{n,r}(A)$ the set of left invertible $n \times r$ matrices with entries in A , and by $GU_r(A^n) \subset U_r(A^n)$ the subset of those left invertible matrices which can be completed to a matrix in $G_n(A)$. For instance $U_0(A^n) = GU_0(A^n) = 0$ is the one element set and $GU_n(A^n) = G_n(A)$. By convention, $U_r(A^n) = GU_r(A^n) = \emptyset$ whenever $r < 0$ or $r > n$. By [NS89, Lemma 2.1] we have $U_r(A^n) = GU_r(A^n)$ for $r \leq n - \text{sr}(A)$.

We define a complex $C(A^n)$ of abelian groups whose degree r component is the free abelian group $C_r(A^n) = \mathbb{Z}[GU_r(A^n)]$ generated by the set $GU_r(A^n)$. For $i = 1, \dots, r$ one has maps of abelian groups $\delta_r^i : C_r(A^n) \rightarrow C_{r-1}(A^n)$ defined on basis elements by $\delta_r^i(v_1, \dots, v_r) = (v_1, \dots, \hat{v}_i, \dots, v_r)$ omitting the i -th entry where (v_i, \dots, v_r) is a left invertible matrix with i -th column the vector v_i . We set

$$(3.1) \quad d_r = \sum_{i=1}^r (-1)^{i-1} \delta_r^i : C_r(A^n) \rightarrow C_{r-1}(A^n),$$

and it is standard that $d_r d_{r+1} = 0$. This defines the chain complex $C(A^n)$.

Lemma 3.1. *Let A be a ring and $n \geq 0$ an integer. Then for all $i \leq n - \text{sr}(A)$ we have*

$$H_i(C(A^n)) = 0.$$

Proof. We check that the proof of [NS89, Lemma 2.2] goes through with $G_n(A)$ in place of $GL_n(A)$. If we denote by $\tilde{C}(A^n)$ the complex with $\tilde{C}_q(A^n) = \mathbb{Z}[U_q(A^n)]$ in degree q and differential given by the same formula as for $C(A^n)$, then we have an inclusion of complexes $C(A^n) \subset \tilde{C}(A^n)$ with $C_q(A^n) = \tilde{C}_q(A^n)$ for $q \leq n - \text{sr } A$, by [NS89, Lemma 2.1]. By [vdK80, 2.6. Theorem (i)] we have $H_i(\tilde{C}(A^n)) = 0$ whenever $i \leq n - \text{sr } A$. Thus, it suffices to show that the boundary dx of every $x \in U_{n-r+1}(A)$ is a boundary in $C(A^n)$ where $r = \text{sr}(A)$. By [NS89, Lemma 2.1], there is a matrix $\alpha \in E_n(A) \subset G_n(A)$ such that $\alpha x = \begin{pmatrix} 1_{n-r} & u \\ 0 & v \end{pmatrix}$ with $u \in M_{n-r,1}(A)$ and $v \in M_{r,1}(A)$. Left invertibility of αx implies that there are $T \in M_{n-r,r}(A)$ and $b \in M_{1,r}$ such that $u + Tv = 0$ and $bv = 1$. The matrix $\beta = \begin{pmatrix} 1_{n-r} & B \\ 0 & 1_{r,r} \end{pmatrix} \begin{pmatrix} 1_{n-r} & -T \\ 0 & 1_{r,r} \end{pmatrix}$ is a product of elementary matrices where $B \in M_{n-r,r}(A)$ is the matrix all of whose rows equal b . Then $\beta\alpha x = \begin{pmatrix} 1_{n-r} & e_1 + \dots + e_{n-r} \\ 0 & v \end{pmatrix}$ and the matrix $w = (\beta\alpha x, e_{n-r+1})$ satisfies $\delta_{n-r+2}^{n-r+2}(w) = \beta\alpha x$. For $i = 1, \dots, n - r + 1$, the matrix $\delta_{n-r+2}^i(w)$ can be completed to the matrix $(\delta_{n-r+2}^i(w), e_{n-r+2}, \dots, e_n) \in SG_n(A)$. It follows that

$$y = x + (-1)^{n-r+2} d\alpha^{-1} \beta^{-1} w \in C_{n-r+1}(A^n)$$

and $dy = dx$. \square

The group $GL_n(A)$ acts on $U_r(A^n)$ by left matrix multiplication, and so does its subgroup $G_n(A)$. This makes the complex $C(A^n)$ into a complex of left $G_n(A)$ -modules. We may sometimes drop the letter A in the notation $G_n(A)$, $SG_n(A)$ etc when the ring A is understood.

Lemma 3.2. *For any ring A , right $G_n(A)$ -module M and integers $i, n \geq 0$ with $i \leq n - \text{sr}(A)$ we have*

$$H_i(M \otimes_{G_n(A)}^L C(A^n)) = 0.$$

Proof. This follows from Lemma 3.1 in view of the spectral sequence

$$E_{p,q}^2 = \text{Tor}_p^{G_n}(M, H_q(C(A^n))) \Rightarrow H_{p+q}(M \otimes_{G_n}^L C(A^n)).$$

\square

For our arguments below we frequently need the following assumptions using the notation from Section 2.

- (*) Let $m, n_0, t \geq 1$, $n \geq 0$ be integers such that A has an $S(m)$ -algebra structure, $n_0 \cdot t < m$ and $0 \leq n \leq n_0$, and t is a multiple of every positive integer $\leq n_0$. Set $\sigma = s_{m,-t} \in \mathbb{Z}[\bar{A}^*]$.

For an integer r , we denote by $C_{\leq r}(A^n)$ the subcomplex of $C(A^n)$ which is $C_{\leq r}(A^n)_i = C_i(A^n)$ for $i \leq r$ and $C_{\leq r}(A^n)_i = 0$ otherwise. So, $C_{\leq r}(A^n)/C_{\leq r-1}(A^n)$ is $C_r(A^n)$ placed in homological degree r . This defines a filtration on $C(A^n)$ by complexes of $G_n(A)$ -modules and thus a spectral sequence of Λ -modules

$$(3.2) \quad E_{p,q}^1(A^n) = \text{Tor}_p^{G_n}(\Lambda_{m,t}, C_q(A^n)) \Rightarrow H_{p+q}(\Lambda_{m,t} \otimes_{G_n}^L C(A^n))$$

with differential d^r of bidegree $(r-1, -r)$. The spectral sequence (3.2) comes with a filtration by Λ -modules

$$0 \subset F_{p+q,0} \subset F_{p+q-1,1} \subset F_{p+q-2,2} \subset \dots \subset F_{0,p+q} = H_{p+q}(\Lambda_{m,t} \otimes_{G_n}^L C(A^n))$$

where $F_{p+q-s,s}$ is the image of

$$H_{p+q}(\Lambda_{m,t} \overset{L}{\otimes}_{G_n} C_{\leq s}(A^n)) \rightarrow H_{p+q}(\Lambda_{m,t} \overset{L}{\otimes}_{G_n} C(A^n)),$$

and $F_{p+q-s,s}/F_{p+q-s+1,s-1} \cong E_{p+q-s,s}^\infty(A^n)$.

Lemma 3.3. *Assume (*). Then the spectral sequence (3.2) has*

$$E_{p,q}^1(A^n) = \begin{cases} H_p(SG_{n-q}(A), \mathbb{Z}), & 0 \leq q < n, \ p \leq n_0 \\ \Lambda_{m,t}, & p = 0, \ q = n \\ 0, & q = n \text{ and } p \neq 0, \text{ or } q < 0, \text{ or } q > n. \end{cases}$$

Proof. By definition, the group $G_n(A)$ acts transitively on the set $GU_q(A^n)$ with stabilizer at $(e_{n-q+1}, \dots, e_n) \in GU_q(A^n)$ the subgroup $\text{Aff}_{q,n-q}^G(A)$. Recall (Shapiro's Lemma) that for any right G -module M we have the quasi-isomorphism

$$M \overset{L}{\otimes}_N \mathbb{Z} \xrightarrow{\sim} M \overset{L}{\otimes}_G \mathbb{Z}[G/N]$$

induced by the inclusions $N \subset G$ and $\mathbb{Z} \subset \mathbb{Z}[G/N]$. For $M = \Lambda_{m,t}$, $G = G_n(A)$ and $N = \text{Aff}_{q,n-q}^G(A)$ we therefore have quasi-isomorphisms

$$\Lambda_{m,t} \overset{L}{\otimes}_{\text{Aff}_{q,n-q}^G} \mathbb{Z} \xrightarrow{\sim} \Lambda_{m,t} \overset{L}{\otimes}_{G_n} \mathbb{Z}[G_n/\text{Aff}_{q,n-q}^G] = \Lambda_{m,t} \overset{L}{\otimes}_{G_n} C_q(A^n).$$

In view of Corollary 2.6, for $q < n$ we have the map of complexes

$$\mathbb{Z} \overset{L}{\otimes}_{SG_{n-q}} \mathbb{Z} \longrightarrow \Lambda_{m,t} \overset{L}{\otimes}_{\text{Aff}_{q,n-q}^G} \mathbb{Z}.$$

which induces an isomorphism on homology in degrees $\leq n_0 < m/t \leq m(n-q)/t$. For $q = n$ we have $C_n(A^n) = \mathbb{Z}[G_n]$ and thus,

$$\Lambda_{m,t} \overset{L}{\otimes}_{G_n} C_n(A^n) = \Lambda_{m,t} \overset{L}{\otimes}_{G_n} \mathbb{Z}[G_n] = \Lambda_{m,t}.$$

□

Lemma 3.4. *Assume (*). Then for $0 < q < n$ and $p \leq n_0$, the differential $d_{p,q}^1 : E_{p,q}^1(A^n) \rightarrow E_{p,q-1}^1(A^n)$ in the spectral sequence (3.2) is zero if q is even and for q odd it is the map*

$$H_p(SG_{n-q}(A), \mathbb{Z}) \rightarrow H_p(SG_{n-q+1}(A), \mathbb{Z})$$

induced by the standard inclusion $SG_{n-q}(A) \rightarrow SG_{n-q+1}(A)$.

For $q = n$ we have $d_{p,n}^1 = 0$ for $p \neq 0$, and $d_{0,n}^1 = 0$ for n even and for n odd $d_{0,n}^1$ is the map $\Lambda_{m,t} \rightarrow \mathbb{Z}$ induced by the augmentation $\varepsilon : \Lambda = \mathbb{Z}[\bar{A}^*] \rightarrow \mathbb{Z}$.

Proof. The differential $d_{p,q}^1$ is the map

$$d_{p,q}^1 = \sum_{j=1}^q (-1)^{j-1} (1 \otimes \delta^j)_* : H_p(\Lambda_{m,t} \overset{L}{\otimes}_{G_n} \mathbb{Z}[GU_q]) \rightarrow H_p(\Lambda_{m,t} \overset{L}{\otimes}_{G_n} \mathbb{Z}[GU_{q-1}])$$

where $\delta^j : \mathbb{Z}[GU_q] \rightarrow \mathbb{Z}[GU_{q-1}]$ is induced by the map $\delta^j : GU_q \rightarrow GU_{q-1}$ defined by $(v_1, \dots, v_q) \mapsto (v_1, \dots, \hat{v}_j, \dots, v_n)$.

Assume first that $q < n$. Consider the diagram

$$\begin{array}{ccccc} \mathbb{Z} \otimes_{SG_{n-q}}^L \mathbb{Z} & \xrightarrow{1 \otimes u_q^n} & \mathbb{Z} \otimes_{SG_n}^L \mathbb{Z}[GU_q(A^n)] & \longrightarrow & \Lambda_{m,t} \otimes_{G_n}^L \mathbb{Z}[GU_q(A^n)] \\ \downarrow & & \downarrow 1 \otimes \delta^j & & \downarrow 1 \otimes \delta^j \\ \mathbb{Z} \otimes_{SG_{n-q+1}}^L \mathbb{Z} & \xrightarrow{1 \otimes u_{q-1}^n} & \mathbb{Z} \otimes_{SG_n}^L \mathbb{Z}[GU_{q-1}(A^n)] & \longrightarrow & \Lambda_{m,t} \otimes_{G_n}^L \mathbb{Z}[GU_{q-1}(A^n)] \end{array}$$

induced by the natural inclusions of groups and rings $SG_{n-q} \subset \text{Aff}_{q,n-q}^{SG} \subset G_n$, $SG_{n-q} \subset SG_{n-q+1}$, $\mathbb{Z} \subset \Lambda_{m,t}$ and where $u_q^n : \mathbb{Z} \rightarrow \mathbb{Z}[GU_q(A^n)]$ sends 1 to the left invertible matrix (e_{n-q+1}, \dots, e_n) . The horizontal compositions induce the isomorphisms in Lemma 3.3 upon taking homology in degrees $\leq n_0$. We will show that the outer diagram commutes upon taking homology groups. This implies the claim for $q < n$. Since the right square commutes, it suffices to show that the left square commutes upon taking homology.

We will use the basic functoriality of group homology as recalled at the beginning of section 2. Upon taking homology, the left hand square of the diagram is the diagram

$$\begin{array}{ccc} H_*(SG_{n-q}, \mathbb{Z}) & \xrightarrow{(i_{n-q}^n, u_q^n)_*} & H_*(SG_n, \mathbb{Z}[GU_q]) \\ \downarrow (i_{n-q}^{n-q+1}, 1)_* & & \downarrow (1, \delta^j)_* \\ H_*(SG_{n-q+1}, \mathbb{Z}) & \xrightarrow{(i_{n-q+1}^n, u_{q-1}^n)_*} & H_*(SG_n, \mathbb{Z}[GU_{q-1}]) \end{array}$$

where u_r^n is the left invertible matrix (e_{n-r+1}, \dots, e_n) and $i_r^s : SG_r \rightarrow SG_s$ denotes the standard embedding for $r \leq s$. Consider the matrix

$$h = (e_1, \dots, e_{n-q}, \tau e_{n-q+j}, \delta^j u_q^n) \in SG_n$$

where $\tau \in \{1, -1\}$ is chosen so that $\det(h) = 1 \in K_1(A)$. Then $h \cdot i_{n-q}^n \cdot h^{-1} = i_{n-q}^n$ and $\delta^j(u_q^n) = h \cdot u_{q-1}^n$. In view of the basic functoriality of group homology, the two compositions $(i_{n-q}^n, \delta^j u_q^n)_*$ and $(i_{n-q}^n, u_{q-1}^n)_*$ are equal, and the square commutes. Hence the lemma for $q < n$.

For $q = n$ and $p \neq 0$ we have $d_{p,n}^1 = 0$ since $E_{p,n}^1(A^n) = 0$. To prove the claim for $q = n$ and $p = 0$, we need to show the commutativity of the diagram

$$\begin{array}{ccc} \Lambda_{m,t} & \xrightarrow[\cong]{1 \otimes u_n^n} & \Lambda_{m,t} \otimes_{G_n} \mathbb{Z}[G_n(A)] \\ \varepsilon \downarrow & & \downarrow 1 \otimes \delta^j \\ \mathbb{Z} & \xrightarrow{1 \otimes u_{n-1}^n} & \Lambda_{m,t} \otimes_{G_n} \mathbb{Z}[GU_{n-1}(A^n)]. \end{array}$$

Since (3.2) is a spectral sequence of $\Lambda_{m,t}$ -modules, this diagram is a diagram of $\Lambda_{m,t}$ -modules. So, it suffices to show that the two compositions send $1 \in \Lambda_{m,t}$ to the same element, that is, we need to see that

$$1 \otimes \delta^j(u_n^n) = 1 \otimes u_{n-1}^n \in \Lambda_{m,t} \otimes_{G_n} \mathbb{Z}[GU_{n-1}(A^n)]$$

for all $j = 1, \dots, n$. Consider the matrix $h_j = (\tau e_j, e_1, e_2, \dots, \hat{e}_j, \dots, e_n) \in SG_n(A)$ where $\tau \in \{1, -1\}$ is chosen so that $\det(h_j) = 1 \in K_1(A)$. Then $\delta^j(u_n^n) = h_j \cdot u_{n-1}^n$ and thus

$$1 \otimes \delta^j(u_n^n) = 1 \otimes h_j \cdot u_{n-1}^n = h_j \otimes u_{n-1}^n = 1 \otimes u_{n-1}^n \in \Lambda_{m,t} \otimes_{G_n} \mathbb{Z}[GU_{n-1}(A^n)]$$

since $h_j \in SG_n(A)$ goes to $1 \in \bar{A}^*$ under the map $G_n(A) \rightarrow \bar{A}^*$. \square

Proposition 3.5. *Assume (*). The differentials $d_{p,q}^r$ in the spectral sequence (3.2) are zero for $r \geq 2$ and $p \leq n_0$.*

Proof. We argue by induction on n . For $n = 0, 1$, the differentials d^r , $r \geq 2$, are zero since $E_{p,q}^r(A^n) = 0$ for $q \neq 0, 1$. Assume $n \geq 2$. Similar to [NS89], consider the homomorphism of complexes $\psi : C(A^{n-2})[-2] \rightarrow C(A^n)$ defined in degree q by $\psi = \psi_0 - \psi_1 + \psi_2$ where for $(v_1, \dots, v_{q-2}) \in GU_{q-2}(A^{n-2})$ the map ψ_i is given by

$$\begin{aligned}\psi_0(v_1, \dots, v_{q-2}) &= (v_1, \dots, v_{q-2}, e_{n-1}, e_n) \\ \psi_1(v_1, \dots, v_{q-2}) &= (v_1, \dots, v_{q-2}, e_{n-1}, e_n - e_{n-1}) \\ \psi_2(v_1, \dots, v_{q-2}) &= (v_1, \dots, v_{q-2}, e_n, e_n - e_{n-1}).\end{aligned}$$

The map ψ commutes with differentials and is compatible with the actions by $G_{n-2}(A)$ and $G_n(A)$ via the standard inclusion $G_{n-2}(A) \subset G_n(A)$. Hence, ψ induces a map of spectral sequences $E(A^{n-2})[0, -2] \rightarrow E(A^n)$. Denote by E and \tilde{E} the spectral sequences $E(A^n)$ and $E(A^{n-2})[0, -2]$. From Lemma 3.3, we have

$$\tilde{E}_{p,q}^1 = E_{p,q-2}^1(A^{n-2}) = \begin{cases} H_p(SG_{n-q}(A), \mathbb{Z}) & 2 \leq q < n, p \leq n_0 \\ \Lambda_{m,t} & q = n, p = 0 \\ 0 & q = n \text{ and } p \neq 0, \text{ or } q < 2, \text{ or } q > n. \end{cases}$$

The claim follows by induction on r using the following lemma. \square

Lemma 3.6. *Assume (*). Under the identifications of Lemma 3.3, the homomorphism $\psi : \tilde{E}_{p,q}^1 \rightarrow E_{p,q}^1$ is the identity for $2 \leq q \leq n$ and $p \leq n_0$.*

Proof. Keeping the notations of the proof of Lemma 3.4, we will check the commutativity of the following diagrams for $j = 0, 1, 2$ upon taking homology groups

$$(3.3) \quad \begin{array}{ccc} \mathbb{Z} \otimes_{SG_{n-q}}^L \mathbb{Z} & \xrightarrow{1 \otimes u_{q-2}^{n-2}} & \mathbb{Z} \otimes_{SG_{n-2}}^L \mathbb{Z}[GU_{q-2}(A^{n-2})] & \longrightarrow & \Lambda_{m,t} \otimes_{G_{n-2}}^L \mathbb{Z}[GU_{q-2}(A^{n-2})] \\ & \searrow^{1 \otimes u_q^n} & \downarrow^{1 \otimes \psi_j} & & \downarrow^{1 \otimes \psi_j} \\ & & \mathbb{Z} \otimes_{SG_n}^L \mathbb{Z}[GU_q(A^n)] & \longrightarrow & \Lambda_{m,t} \otimes_{G_n}^L \mathbb{Z}[GU_q(A^n)] \end{array}$$

$$(3.3) \quad \begin{array}{ccc} \Lambda_{m,t} & \xrightarrow{1 \otimes u_{n-2}^{n-2}} & \Lambda_{m,t} \otimes_{G_{n-2}} \mathbb{Z}[G_{n-2}] \\ & \searrow^{1 \otimes u_n^n} & \downarrow^{1 \otimes \psi_j} \\ & & \Lambda_{m,t} \otimes_{G_n} \mathbb{Z}[G_n] \end{array}$$

where in the first diagram $2 \leq q < n$, and the second diagram takes care of $q = n$. Since $\psi = \psi_0 - \psi_1 + \psi_2$, commutativity of the diagrams on homology groups will imply the claim.

In the first diagram the right hand square commutes, and so we are left with showing the commutativity of the left hand square on homology, that is, the commutativity of the diagram

$$(3.4) \quad \begin{array}{ccc} H_p(SG_{n-q}, \mathbb{Z}) & \xrightarrow{(i_{n-q}^{n-2}, u_{q-2}^{n-2})^*} & H_p(SG_{n-2}, \mathbb{Z}[GU_{q-2}(A^{n-2})]) \\ & \searrow (i_{n-q}^n, u_q^n)^* & \downarrow (i_{n-2}^n, \psi_j)^* \\ & & H_p(SG_n, \mathbb{Z}[GU_q(A^n)]). \end{array}$$

Commutativity for $j = 0$ is clear. For $j = 1$ we consider the following matrix $h = (e_1, \dots, e_{n-1}, e_n - e_{n-1}) \in SG_n(A)$. For $2 \leq q < n$ we have $h \cdot i_{n-q}^n \cdot h^{-1} = i_{n-q}^n$ and $h \circ u_q^n = \psi_1(u_{q-2}^{n-2})$, this shows commutativity of diagram (3.4) in this case. For $j = 2$, we replace the matrix h with the matrix $(e_1, \dots, e_{n-2}, e_n, e_n - e_{n-1})$ in $SG_n(A)$. This finishes the proof of commutativity of (3.4) for $j = 0, 1, 2$.

To show commutativity of the second diagram (3.3), note that it is a diagram of $\Lambda_{m,t}$ -modules and that the horizontal and diagonal arrows are isomorphisms with inverses the multiplication maps $\Lambda_{m,t} \otimes_G \mathbb{Z}[G] \xrightarrow{1 \otimes \det} \Lambda_{m,t} \otimes \Lambda_{m,t} \rightarrow \Lambda_{m,t}$. Since $\psi_j(u_{n-2}^n) = 1 = u_n^n \in \bar{A}^* \subset K_1(A)$, the claim follows. \square

Theorem 3.7. *Let A be a ring with many units. Then the natural homomorphism*

$$H_i(SG_{n-1}(A), \mathbb{Z}) \rightarrow H_i(SG_n(A), \mathbb{Z})$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

Proof. Choose n_0, m, t as in (*). In particular $n \leq n_0$. Then we have the spectral sequence (3.2) with E^1 -term given by Lemma 3.3 and $d_{p,q}^1$ was computed in Lemma 3.4 for $p \leq n_0$. By Proposition 3.5 and Lemma 3.2 we have $E_{p,q}^2 = E_{p,q}^\infty = 0$ for $n \geq p + q + \text{sr}(A)$ and $p \leq n_0$. The claim follows. \square

We can reformulate Theorem 3.7 in terms of elementary linear groups. Recall [Bas64, §1] that the group of elementary $r \times r$ -matrices of a ring A is the subgroup $E_r(A)$ of $GL_r(A)$ generated by the elementary matrices $e_{i,j}(a) = 1 + a \cdot e_i e_j^T$, $a \in A$.

Lemma 3.8. *Let A be a ring with many units. Then for $n > \text{sr}(A)$, we have $E_n(A) = SG_n(A)$, and this group is the commutator and the maximal perfect subgroup of $GL_n(A)$. Moreover, the natural map $GL_n(A) \rightarrow K_1(A)$ is surjective for $n \geq \text{sr}(A)$.*

Proof. We clearly have $E_n(A) \subset SG_n(A)$. It follows from the GL_n -version of Theorem 3.7 proved in [NS89] that the natural map $GL_n(A) \rightarrow K_1(A)$ is surjective for $n \geq \text{sr}(A)$ and $GL_n(A)^{ab} = K_1(A)$ for $n > \text{sr}(A)$. Therefore, $GL_n(A)/SG_n(A) \cong K_1(A)$ for $n \geq \text{sr}(A)$ and $SG_n(A) = [GL_n(A), GL_n(A)]$ for $n > \text{sr}(A)$.

For the rest of the proof assume $n > \text{sr}(A)$. From [Vas69, Theorem 3.2], we have $GL_n(A)/E_n(A) = K_1(A)$, hence $E_n(A) = SG_n(A)$. Classically, the group $E_n(A)$ is perfect for $n \geq 3$ [Bas64, Corollary 1.5]. Alternatively, from Theorem 3.7, we have an isomorphism

$$H_1(SG_n(A)) \cong H_1(E(A)) = 0$$

since the infinite elementary linear group $E(A)$ is perfect and $\text{colim}_r SG_r(A) = E(A)$. Hence, the group $E_n(A)$ is perfect (also when $n = 2$). Since the quotient $GL_n(A)/E_n(A) = K_1(A)$ is abelian, the group $E_n(A)$ is the maximal perfect subgroup of $GL_n(A)$. \square

Theorem 3.9. *Let A be a ring with many units. Then the natural homomorphism*

$$H_i(E_{n-1}(A), \mathbb{Z}) \rightarrow H_i(E_n(A), \mathbb{Z})$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

Proof. This follows from Theorem 3.7 in view of Lemma 3.8. \square

Denote by $BGL_n^+(A)$ the space obtained by applying Quillen's plus construction to the classifying space $BGL_n(A)$ of $GL_n(A)$ with respect to the maximal perfect subgroup of $GL_n(A)$. The following proves a conjecture of Bass [Bas73, Conjecture XVI on p. 43] in the case of rings with many units.

Theorem 3.10. *Let A be a ring with many units, and let $n \geq 1$ be an integer. Then the natural homomorphism*

$$\pi_i BGL_{n-1}^+(A) \rightarrow \pi_i BGL_n^+(A)$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

Proof. The case $i = 0$ is trivial and the case $i = 1$ follows at once from Lemma 3.8.

Now, assume $i \geq 2$ and $n \geq i + \text{sr}(A) \geq 2 + \text{sr}(A)$. Denote by $\mathcal{F}_n(A)$ the homotopy fibre of the map $BGL_{n-1}^+(A) \rightarrow BGL_n^+(A)$. It follows from Lemma 3.8, that \mathcal{F}_n is also the homotopy fibre of $BE_{n-1}^+(A) \rightarrow BE_n^+(A)$. Since $E_{n-1}(A)$ and $E_n(A)$ are perfect, $\mathcal{F}_n(A)$ is connected and $\pi_1 \mathcal{F}_n(A)$ is abelian as a quotient of $\pi_2 BE_n^+(A)$. From Theorem 3.9 and the (relative) Serre spectral sequence

$$E_{r,s}^2 = H_r(BE_n^+(A), H_s(\text{pt}, \mathcal{F}_n)) \Rightarrow H_{r+s}(BE_n^+(A), BE_{n-1}^+(A))$$

associated to the fibration $\mathcal{F}_n \rightarrow BE_{n-1}^+(A) \rightarrow BE_n^+(A)$ with simply connected base we find

$$H_{i-1}(\mathcal{F}_n, \text{pt}) = H_i(\text{pt}, \mathcal{F}_n) = H_i(E_n(A), E_{n-1}(A))$$

for $i \leq n - \text{sr}(A) + 1$ where $\text{pt} \in \mathcal{F}_n(A)$ denotes the base point of $\mathcal{F}_n(A)$. By Hurewicz's Theorem, it follows from Theorem 3.9 that the natural map

$$\pi_{i-1}(\mathcal{F}_n(A), \text{pt}) \rightarrow H_{i-1}(\mathcal{F}_n(A), \text{pt})$$

is an isomorphism for $i \leq n - \text{sr}(A) + 1$. In particular, we have

$$\pi_{i-1}(\mathcal{F}_n(A), \text{pt}) = H_i(E_n(A), E_{n-1}(A)) = 0, \quad i \leq n - \text{sr}(A).$$

Hence the result. \square

Recall that for a commutative ring A and integer $n \geq 1$, the special linear group $SL_n(A)$ of A is the kernel of the determinant map $GL_n(A) \rightarrow A^*$.

Theorem 3.11. *Let A be a local commutative ring with infinite residue field, and let $n \geq 2$ be an integer. Denote by $\mathcal{F}_n(A)$ the homotopy fibre of the map $BGL_{n-1}^+(A) \rightarrow BGL_n^+(A)$. Then for $i \geq 1$ we have*

$$\pi_{i-1} \mathcal{F}_n(A) = \begin{cases} 0, & i < n \\ H_i(SL_n(A), SL_{n-1}(A)), & i = n. \end{cases}$$

Proof. Note that for A as in the theorem $E_r(A) = SL_r(A)$ is the maximal perfect subgroup of $GL_r(A)$ for all $r \geq 1$. Now the proof is the same as for Theorem 3.10, the only improvement being that \mathcal{F}_n is already the homotopy fibre of $BE_{n-1}(A)^+ \rightarrow BE_n(A)^+$ for $n \geq 2$. \square

Theorem 3.12. *Let A be a commutative ring with many units and $n \geq 2$ an integer. Then the natural homomorphism*

$$H_i(SL_{n-1}(A), \mathbb{Z}) \rightarrow H_i(SL_n(A), \mathbb{Z})$$

is an isomorphism for $n \geq i + \text{sr}(A) + 1$ and surjective for $n \geq i + \text{sr}(A)$.

Proof. For $r \geq \text{sr}(A)$, the natural map $GL_r(A) \rightarrow K_1(A)$ is surjective, by Lemma 3.8. When A is commutative we have an exact sequence of groups for $r \geq \text{sr}(A)$

$$1 \rightarrow SG_r(A) \rightarrow SL_r(A) \rightarrow SK_1(A) \rightarrow 0.$$

Hence, for $n \geq \text{sr}(A) + 1$ we have the associated Serre spectral sequence

$$H_i(SK_1(A), H_j(SG_n A, SG_{n-1} A)) \Rightarrow H_{i+j}(SL_n(A), SL_{n-1}(A)).$$

The result now follows from Theorem 3.7. \square

4. MILNOR-WITT K -THEORY

Let A be a commutative ring. Recall that we denote by A^* the group of units in A . Elements in the integral group ring $\mathbb{Z}[A^*]$ of A^* corresponding to $a \in A^*$ are denoted by $\langle a \rangle$. Note that $\langle 1 \rangle = 1 \in \mathbb{Z}[A^*]$. We denote by $\langle\langle a \rangle\rangle$ the element $\langle\langle a \rangle\rangle = \langle a \rangle - 1 \in \mathbb{Z}[A^*]$. Let $I[A^*]$ be the augmentation ideal in $\mathbb{Z}[A^*]$, that is, $I[A^*]$ is the kernel of the ring homomorphism $\mathbb{Z}[A^*] \rightarrow \mathbb{Z} : \langle a \rangle \mapsto 1$. We denote by $[a]$ the element $[a] = \langle a \rangle - 1 \in I[A^*]$. Under the canonical embedding $I[A^*] \subset \mathbb{Z}[A^*]$, the element $[a]$ maps to $\langle\langle a \rangle\rangle$.

Lemma 4.1. *The augmentation ideal $I[A^*]$ is the $\mathbb{Z}[A^*]$ -module generated by symbols $[a]$ for $a \in A^*$ subject to the relation*

$$[ab] = [a] + \langle a \rangle [b].$$

Proof. Clearly, the equation $[ab] = [a] + \langle a \rangle [b]$ holds in $I[A^*]$. Let \tilde{I}_{A^*} be the $\mathbb{Z}[A^*]$ -module generated by symbols $[a]$ for $a \in A^*$ subject to the relation $[ab] = [a] + \langle a \rangle [b]$. Note that $[1] = 0$ in \tilde{I}_{A^*} because $[1 \cdot 1] = [1] + \langle 1 \rangle [1]$ and $\langle 1 \rangle = 1 \in \mathbb{Z}[A^*]$.

Consider the $\mathbb{Z}[A^*]$ -module map $\tilde{I}_{A^*} \rightarrow I[A^*] : [a] \mapsto \langle a \rangle - 1$. The set consisting of $\langle a \rangle - 1 \in I[A^*]$, $a \in A^*$, $a \neq 1$, defines a \mathbb{Z} -basis of $I[A^*]$. This allows us to define a \mathbb{Z} -linear homomorphism $I[A^*] \rightarrow \tilde{I}_{A^*} : \langle a \rangle - 1 \mapsto [a]$. Clearly, the composition $I[A^*] \rightarrow \tilde{I}_{A^*} \rightarrow I[A^*]$ is the identity. Finally, the map $I[A^*] \rightarrow \tilde{I}_{A^*}$ is surjective because $\langle b \rangle [a] = [ab] - [b]$. \square

Definition 4.2. Let A be a commutative ring. We define the graded ring $\hat{K}_*^{MW}(A)$ as the tensor algebra of the augmentation ideal $I[A^*]$ (placed in degree 1) over the group ring $\mathbb{Z}[A^*]$ modulo the Steinberg relation $[a][1-a] = 0$ for $a, 1-a \in A^*$:

$$\hat{K}_*^{MW}(A) = \bigoplus_{n \geq 0} \hat{K}_n^{MW}(A) = \text{Tens}_{\mathbb{Z}[A^*]}(I[A^*]) / [a][1-a].$$

In view of Lemma 4.1, the graded ring $\hat{K}_*^{MW}(A)$ is the $\mathbb{Z}[A^*]$ -algebra generated by symbols $[a]$, $a \in A^*$, in degree 1 subject to the relations

- (1) For $a, b \in A^*$ we have $[ab] = [a] + \langle a \rangle [b]$.
- (2) For $a, 1-a \in A^*$ we have the Steinberg relation $[a][1-a] = 0$.

It is convenient to write $[a_1, \dots, a_n]$, h and ε for the following elements in $\hat{K}_*^{MW}(A)$

$$[a_1, \dots, a_n] = [a_1] \cdots [a_n] \in \hat{K}_n^{MW}(A),$$

$$h = 1 + \langle -1 \rangle, \quad \varepsilon = -\langle -1 \rangle \in \hat{K}_0^{MW}(A)$$

where $a_1, \dots, a_n \in A^*$.

Lemma 4.3. *Let A be a commutative ring. Then in the ring $\hat{K}_*^{MW}(A)$ we have for all $a, b, c, d \in A^*$ the following relations.*

- (1) $[ab] = [a] + \langle a \rangle [b]$
- (2) $\langle 1 \rangle = 1$ and $[1] = 0$,
- (3) $\langle ab \rangle = \langle a \rangle \cdot \langle b \rangle$ and $\langle a \rangle$ is central in $\hat{K}_*^{MW}(A)$.
- (4) $\left[\frac{a}{b}\right] = [a] - \langle \frac{a}{b} \rangle [b]$, in particular, $[b^{-1}] = -\langle b^{-1} \rangle [b]$.
- (5) $\langle \langle a \rangle \rangle [b] = \langle \langle b \rangle \rangle [a]$.
- (6) If $a + b = 1$ then $\langle \langle a \rangle \rangle [b, c] = 0$.
- (7) If $a + b = 1$ then $\langle \langle a \rangle \rangle \langle \langle b \rangle \rangle [c, d] = 0$.
- (8) $\langle \langle a \rangle \rangle [b, c] = \langle \langle a \rangle \rangle [c, b]$.

Proof. Items (1)- (3) follow from the definition of $\hat{K}_*^{MW}(A)$.

(4) We have $[a] = \left[\frac{a}{b} \cdot b\right] = \left[\frac{a}{b}\right] + \langle \frac{a}{b} \rangle [b]$.

(5) We have $[a] + \langle a \rangle [b] = [ab] = [b] + \langle b \rangle [a]$ which is $\langle \langle a \rangle \rangle [b] = \langle \langle b \rangle \rangle [a]$.

(6) By (5) and the Steinberg relation we have $\langle \langle a \rangle \rangle [b, c] = \langle \langle c \rangle \rangle [b, a] = 0$.

(7) Similarly, we have $\langle \langle a \rangle \rangle \langle \langle b \rangle \rangle [c, d] = \langle \langle c \rangle \rangle \langle \langle d \rangle \rangle [a, b] = 0$, by (5) and the Steinberg relation.

(8) From (5) and (3) we have $\langle \langle a \rangle \rangle [b, c] = \langle \langle b \rangle \rangle [a, c] = \langle \langle c \rangle \rangle [a, b] = \langle \langle a \rangle \rangle [c, b]$. \square

Lemma 4.4. *Let A be either a field or a commutative local ring whose residue field has at least 4 elements. Then for all $a, b, c \in A^*$ the following relations hold in $\hat{K}_*^{MW}(A)$.*

- (1) $[a][-a] = 0$
- (2) $[a][a] = [a][-1] = [-1][a]$
- (3) $[a][b] = \varepsilon [b][a]$ where $\varepsilon = -\langle -1 \rangle$
- (4) $\langle \langle a \rangle \rangle \cdot h \cdot [b, c] = 0$ where $h = 1 + \langle -1 \rangle$.
- (5) $[a^2, b] = h \cdot [a, b]$
- (6) $\langle \langle a^2 \rangle \rangle [b, c] = 0$, in particular, $\langle a^2 \rangle [b, c] = [b, c]$.

Proof. (1) First assume $\bar{a} \neq 1$ where \bar{a} means reduction modulo the maximal ideal in A . Then $1 - a, 1 - a^{-1} \in A^*$ and $-a = \frac{1-a}{1-a^{-1}}$. Therefore, $[a][-a] = [a]\left[\frac{1-a}{1-a^{-1}}\right] = [a]([1-a] - \langle -a \rangle [1-a^{-1}]) = -\langle -a \rangle [a][1-a^{-1}] = -\langle -a \rangle \langle a \rangle [a^{-1}][1-a^{-1}] = 0$ where the second to last equation is from Lemma 4.3 (4). This already implies the case when A is a field.

Now assume that A is local whose residue field has at least 4 elements. Assume $\bar{a} = 1$ and choose $b \in A^*$ with $\bar{b} \neq 1$. Then $\bar{a}\bar{b} \neq 1$. Therefore, $0 = [ab][-ab] = ([a] + \langle a \rangle [b])[-ab] = [a][-ab] + \langle a \rangle [b][-ab] = [a]([-a] + \langle -a \rangle [b]) + \langle a \rangle [b]([a] + \langle a \rangle [-b]) = [a][-a] + \langle -a \rangle [a][b] + \langle a \rangle [b][a]$. Hence, for all $\bar{b} \neq 1$ we have $[a][-a] = -\langle -a \rangle [a][b] - \langle a \rangle [b][a]$. Now, choose $b_1, b_2 \in A^*$ such that $\bar{b}_1, \bar{b}_2, \bar{b}_1\bar{b}_2 \neq 1$. This is possible if the residue field of A has at least 4 elements. Then $[a][-a] = -\langle -a \rangle [a][b_1b_2] - \langle a \rangle [b_1b_2][a] = -\langle -a \rangle [a](\langle b_1 \rangle + \langle b_2 \rangle [b_1]) - \langle a \rangle (\langle b_1 \rangle + \langle b_2 \rangle [b_1])[a] = -\langle -a \rangle [a][b_1] - \langle a \rangle [b_1][a] + \langle b_1 \rangle (-\langle -a \rangle [a][b_2] - \langle a \rangle [b_2][a]) = [a][-a] + \langle b_1 \rangle [a][-a]$. Hence, $\langle b_1 \rangle [a][-a] = 0$. Multiplying with $\langle b_1^{-1} \rangle$ yields the result.

(2) We have $[a][a] = [(-1)(-a)][a] = ([-1] + \langle -1 \rangle)[-a][a] = [-1][a]$. Similarly, $[a][a] = [a][(-1)(-a)] = [a][(-1) + \langle -1 \rangle][-a] = [a][-1]$.

(3) We have $0 = [ab][-ba] = ([a] + \langle a \rangle)[b][-ab] = [a][-ab] + \langle a \rangle[b][-ab] = [a][(-a) + \langle -a \rangle][b] + \langle a \rangle[b][(-a) + \langle -a \rangle][b] = \langle -a \rangle[a][b] + \langle a \rangle[b][a]$. Multiplying with $\langle a^{-1} \rangle$ yields $[a][b] = -\langle -1 \rangle[b][a]$.

(4) From Lemma 4.3 (8) and Lemma 4.4 (3) we have $\langle \langle a \rangle \rangle[b, c] = -\langle -1 \rangle \langle \langle a \rangle \rangle[b, c]$ and thus $\langle \langle a \rangle \rangle(1 + \langle -1 \rangle)[b, c] = 0$.

(5) We have $[a^2, b] = [a, b] + \langle a \rangle[a, b] = 2[a, b] + \langle \langle a \rangle \rangle[a, b] = 2[a, b] + \langle \langle b \rangle \rangle[a, a] = 2[a, b] + \langle \langle b \rangle \rangle[-1, a] = 2[a, b] + \langle \langle -1 \rangle \rangle[a, b] = h \cdot [a, b]$.

(6) We have $\langle \langle a^2 \rangle \rangle[b, c] = \langle \langle b \rangle \rangle[a^2, c] = \langle \langle b \rangle \rangle \cdot h \cdot [a, c] = 0$. \square

Corollary 4.5. *Let A be either a field or a commutative local ring whose residue field has at least 4 elements. Then in $\hat{K}_*^{MW}(A)$ we have $[a_1, \dots, a_n] = 0$ if $a_i + a_j = 1$ or $a_i + a_j = 0$ for some $i \neq j$.*

Proof. This follows from the Steinberg relation and Lemma 4.4 (1) and (3). \square

Definition 4.6. Let A be a commutative ring. We define the ring

$$GW(A)$$

as the quotient of the group ring $\mathbb{Z}[A^*]$ modulo the following relations.

- (1) For all $a \in A^*$ we have $\langle \langle a \rangle \rangle h = 0$.
- (2) (Steinberg relation) For all $a, 1 - a \in A^*$ we have $\langle \langle a \rangle \rangle \langle \langle 1 - a \rangle \rangle = 0$.

Definition 4.7. Let A be a commutative ring. We define the $\mathbb{Z}[A^*]$ -module

$$V(A)$$

as the quotient of the augmentation ideal $I[A^*]$ modulo the relations

- (1) For all $a, b \in A^*$ we have $\langle \langle a \rangle \rangle \cdot h \cdot [b] = 0$.
- (2) (Steinberg relation) For all $a, 1 - a \in A^*$ we have $\langle \langle a \rangle \rangle [1 - a] = 0$.

Since $\langle \langle a \rangle \rangle [b] = \langle \langle b \rangle \rangle [a]$ in $I[A^*]$ and hence in $V(A)$, we see that the $\mathbb{Z}[A^*]$ -module $V(A)$ is naturally a $GW(A)$ -module.

Proposition 4.8. *Let A be commutative ring. Then the natural surjections $\mathbb{Z}[A^*] \rightarrow GW(A)$ and $I[A^*] \rightarrow V(A)$ induce a surjective map of graded rings*

$$\hat{K}_*^{MW}(A) \rightarrow \text{Tens}_{GW(A)} V(A) / [a][1 - a].$$

If A is either a field or a local ring whose residue field has at least 4 elements then this map is an isomorphism in degrees ≥ 2 .

Proof. It is clear that the map in the proposition is a surjective ring homomorphism. Its kernel is the (homogeneous) ideal in $\hat{K}_*^{MW}(A)$ generated by the elements of the form $\langle \langle a \rangle \rangle h$ ($a \in A^*$), $\langle \langle a \rangle \rangle \langle \langle 1 - a \rangle \rangle$ and $\langle \langle a \rangle \rangle [1 - a]$ ($a, 1 - a \in A^*$). If A is a field or a local ring with residue field cardinality ≥ 4 , then this ideal is zero in degrees ≥ 2 , in view of Lemmas 4.3 (6), (7) and 4.4 (3), (4). \square

Remark 4.9. Since the homomorphism of graded rings in Proposition 4.8 is surjective for any commutative ring A , all formulas in Lemma 4.3 also hold in the target of that map.

In case A is a field the following definition is due to Hopkins and Morel [Mor12, Definition 3.1]. For commutative local rings, the definition was also considered in [GSZ16].

Definition 4.10. Let A be a commutative ring. The Milnor-Witt K -theory of A is the graded associative ring $K_*^{MW}(A)$ generated by symbols $[a]$, $a \in A^*$, of degree 1 and one symbol η of degree -1 subject to the following relations.

- (1) For $a, 1 - a \in A^*$ we have $[a][1 - a] = 0$.
- (2) For $a, b \in A^*$, we have $[ab] = [a] + [b] + \eta[a][b]$.
- (3) For each $a \in A^*$, we have $\eta[a] = [a]\eta$, and
- (4) $\eta^2[-1] + 2\eta = 0$.

Definition 4.11. Let A be a commutative ring and n an integer. Let $\tilde{K}_n^{MW}(A)$ be the abelian group generated by symbols of the form $[\eta^m, u_1, \dots, u_{n+m}]$ with $m \geq 0$, $n + m \geq 0$, $u_i \in A^*$, subject to the following three relations.

- (1) $[\eta^m, u_1, \dots, u_{n+m}] = 0$ if $u_i + u_{i+1} = 1$ for some $i = 1, \dots, n + m - 1$.
- (2) For all $a, b \in A^*$, $m \geq 0$ and $i = 1, \dots, n + m$ we have

$$[\eta^m, u_1, \dots, u_{i-1}, ab, u_{i+1}, \dots] =$$

$$[\eta^m, \dots, u_{i-1}, a, u_{i+1}, \dots] + [\eta^m, \dots, u_{i-1}, b, u_{i+1}, \dots] + [\eta^{m+1}, \dots, u_{i-1}, a, b, u_{i+1}, \dots]$$

- (3) For each $m \geq 0$ and $i = 1, \dots, n + m + 2$ we have

$$[\eta^{m+2}, u_1, \dots, u_{i-1}, -1, u_{i+1}, \dots, u_{n+m+2}] + 2[\eta^{m+1}, \dots, u_{i-1}, u_{i+1}, \dots, u_{n+m+2}] = 0.$$

We make $\tilde{K}_*^{MW}(A) = \bigoplus_n \tilde{K}_n^{MW}(A)$ into a graded ring with multiplication

$$\tilde{K}_r^{MW}(A) \otimes \tilde{K}_s^{MW}(A) \rightarrow \tilde{K}_{r+s}^{MW}(A)$$

defined by

$$[\eta^m, u_1, \dots, u_{r+m}] \otimes [\eta^n, v_1, \dots, v_{s+n}] \mapsto [\eta^{m+n}, u_1, \dots, u_{r+m}, v_1, \dots, v_{s+n}].$$

By going through the 3 relations in Definition 4.11 this map is well-defined as map of abelian groups. The multiplication is obviously associative and unital with unit $1 = [\eta^0]$. We define a map of graded rings

$$K_*^{MW}(A) \rightarrow \tilde{K}_*^{MW}(A)$$

by sending η to $[\eta]$ and $[u]$ to $[\eta^0, u]$. It is easy to check that the defining relations for $K_*^{MW}(A)$ hold in $\tilde{K}_*^{MW}(A)$.

Lemma 4.12. [Mor12, Lemma 3.4] *For any commutative ring A , the maps*

$$\tilde{K}_n^{MW}(A) \rightarrow K_n^{MW}(A) : [\eta^m, u_1, \dots, u_{n+m}] \rightarrow \eta^m [u_1] \cdots [u_{n+m}]$$

define an isomorphism of graded rings

$$\tilde{K}_*^{MW}(A) \xrightarrow{\cong} K_*^{MW}(A).$$

Proof. The composition $\tilde{K}_*^{MW}(A) \rightarrow K_*^{MW}(A) \rightarrow \tilde{K}_*^{MW}(A)$ is the identity, and the first map is surjective. \square

For $a \in A^*$ set $\langle a \rangle = 1 + \eta[a] \in K_0^{MW}(A)$ and $\langle \langle a \rangle \rangle = \langle a \rangle - 1 = \eta[a] \in K_0^{MW}(A)$.

Lemma 4.13. *Let A be a commutative ring. Then for all $a, b \in A^*$ we have in $K_*^{MW}(A)$ the following.*

- (1) $[ab] = [a] + \langle a \rangle [b]$
- (2) $\langle 1 \rangle = 1$ and $[1] = 0$,
- (3) $[\frac{a}{b}] = [a] - \langle \frac{a}{b} \rangle [b]$, in particular, $[b^{-1}] = -\langle b^{-1} \rangle [b]$.
- (4) $\langle ab \rangle = \langle a \rangle \cdot \langle b \rangle$ and $\langle a \rangle$ is central in $K_*^{MW}(A)$.

- (5) The map $\mathbb{Z}[A^*] \rightarrow K_0^{MW}(A) : \langle a \rangle \mapsto \langle a \rangle$ is a surjective ring homomorphism making $K_*^{MW}(A)$ into a $\mathbb{Z}[A^*]$ -algebra.
(6) $\langle \langle a \rangle \rangle \cdot h = 0$ where $h = 1 + \langle -1 \rangle$.
(7) If $a + b = 1$ then $\langle \langle a \rangle \rangle [b] = 0$.

Proof. (1) We have $[ab] = [a] + [b] + \eta[a][b] = [a] + (1 + \eta[a])[b] = [a] + \langle a \rangle [b]$.

(2) We have $[-1] = [-1] + [1] + \eta[-1][1]$, hence $0 = \eta[1] + \eta^2[-1][1] = \eta[1] - 2\eta[1] = -\eta[1]$. This shows $\langle 1 \rangle = 1$. Now, $[1] = [1 \cdot 1] = [1] + \langle 1 \rangle [1] = [1] + [1]$ from which we obtain $[1] = 0$.

(3) We have $[a] = [\frac{a}{b}b] = [\frac{a}{b}] + \langle \frac{a}{b} \rangle [b]$.

(4) We have $\langle a \rangle \cdot \langle b \rangle = (1 + \eta[a])(1 + \eta[b]) = 1 + \eta([a] + [b] + \eta[a][b]) = 1 + \eta[ab] = \langle ab \rangle$. Moreover, $\langle a \rangle [b] = (1 + \eta[a])[b] = [b] + \eta[a][b] = [ab] - [a] = [ba] - [a]$ whereas $[b]\langle a \rangle = [b](1 + \eta[a]) = [b] + \eta[b][a] = [ba] - [a]$. Hence $\langle a \rangle [b] = [b]\langle a \rangle$.

(5) It is clear that $\mathbb{Z}[A^*] \rightarrow K_0^{MW}(A)$ is a ring homomorphism. By Lemma 4.12, the group $K_0^{MW}(A)$ is additively generated by $[\eta^0]$ and $[\eta, a]$, equivalently, by $[\eta^0] = 1 = \langle 1 \rangle$ and $[\eta^0] + [\eta, a] = \langle a \rangle$. Hence, the map of rings is surjective. The ring $K_*^{MW}(A)$ is a $\mathbb{Z}[A^*]$ -algebra since $K_0^{MW}(A)$ is central in $K_*^{MW}(A)$.

(6) We have $\langle a \rangle \cdot h = (1 + \eta[a]) \cdot h = h$ because $\eta h = 0$. Hence $\langle \langle a \rangle \rangle \cdot h = 0$

(7) If $a + b = 1$ then $\langle \langle a \rangle \rangle [b] = \eta[a][b] = 0$. \square

Lemma 4.14. *Let A be a commutative ring. Then the following map defines an isomorphism of rings*

$$GW(A) \xrightarrow{\cong} K_0^{MW}(A) : \langle a \rangle \mapsto \langle a \rangle$$

Proof. By Lemma 4.13 the map in the lemma is a surjective ring homomorphism. Using Lemma 4.12, we define the inverse by

$$\tilde{K}_0^{MW}(A) \rightarrow GW(A) : [\eta^m, a_1, \dots, a_m] \mapsto \prod_{i=1}^m \langle \langle a_i \rangle \rangle$$

It is easy to check that this also defines a surjective ring homomorphism. Since the composition $GW(A) \rightarrow \tilde{K}_0^{MW}(A) \rightarrow GW(A)$ is the identity, we are done. \square

Lemma 4.15. *Let A be a commutative ring. Then for $a, b \in A^*$ we have in $V(A)$*

- (1) $\langle \langle a \rangle \rangle [b] = \langle \langle b \rangle \rangle [a]$
(2) $h[b] = 2[b] + \langle \langle b \rangle \rangle [-1]$

Proof. (1) The equation holds in $I[A^*]$ and hence in its quotient $V(A)$.

(2) We have $h[b] = [b] + \langle -1 \rangle [b] = 2[b] + \langle \langle -1 \rangle \rangle [b] = 2[b] + \langle \langle b \rangle \rangle [-1]$. \square

Lemma 4.16. *Let A be a commutative ring. Then we have an isomorphism of $GW(A) = K_0^{MW}(A)$ -modules*

$$V(A) \xrightarrow{\cong} K_1^{MW}(A) : [a] \mapsto [a].$$

Proof. By Lemma 4.13 the map in the lemma is well-defined. Using Lemma 4.12 we define the inverse by

$$\tilde{K}_1^{MW}(A) \rightarrow V(A) : [\eta^m, u_1, \dots, u_{m+1}] \mapsto \left(\prod_{i=1}^m \langle \langle u_i \rangle \rangle \right) [u_{m+1}].$$

This map preserves the relations of Definition 4.11, the last one follows from Lemma 4.15 (2) which implies $0 = \langle \langle a \rangle \rangle h[b] = \langle \langle a \rangle \rangle \langle \langle b \rangle \rangle [-1] + 2\langle \langle a \rangle \rangle [b] \in V(A)$. \square

For A a field, the following is a remark (without proof) in [Mor12].

Proposition 4.17. *Let A be a commutative ring. The isomorphisms $GW(A) \cong K_0^{MW}(A)$ and $V(A) \cong K_1^{MW}(A)$ in Lemmas 4.14 and 4.16 extend to an isomorphism of graded rings*

$$\mathrm{Tens}_{GW(A)} V(A) / \{[a][1-a] \mid a, 1-a \in A^*\} \xrightarrow{\cong} K_{\geq 0}^{MW}(A).$$

Proof. Using Lemma 4.12, the inverse is given by the ring map

$$\begin{aligned} \tilde{K}_{\geq 0}^{MW}(A) &\rightarrow \mathrm{Tens}_{GW(A)} V(A) / \{[a][1-a] \mid a, 1-a \in A^*\} \\ [\eta^m, u_1, \dots, u_{m+n}] &\mapsto \left(\prod_{i=1}^m \langle\langle u_i \rangle\rangle \right) [u_{m+1}] \cdots [u_{m+n}] \end{aligned}$$

This map is well-defined and indeed the required inverse in view of Lemma 4.15 and Remark 4.9. \square

Now we come to the main result of this section. For fields, a related but different presentation is given in [HT13].

Theorem 4.18. *Let A be either a field or a local ring whose residue field has at least 4 elements. Then the homomorphism of graded $\mathbb{Z}[A^*]$ -algebras*

$$\hat{K}_*^{MW}(A) \rightarrow K_*^{MW}(A) : [a] \mapsto [a]$$

induces an isomorphism for all $n \geq 2$

$$\hat{K}_n^{MW}(A) \xrightarrow{\cong} K_n^{MW}(A).$$

Proof. The theorem follows from Propositions 4.8 and 4.17. \square

Proposition 4.19. *Let A be either a field or a local ring with residue field cardinality at least 4. Let $n \geq 1$ an integer. Then for $a_i \in A^*$ ($1 \leq i \leq n$) and $\lambda_i \in A^*$ ($1 \leq i \leq n$) with $\bar{\lambda}_i \neq \bar{\lambda}_j$ for $i \neq j$ ($\bar{\lambda}$ denotes reduction of λ modulo the maximal ideal), the following relation holds in $\hat{K}_*^{MW}(A)$:*

$$\begin{aligned} &[\lambda_1 a_1, \dots, \lambda_n a_n] - [a_1, \dots, a_n] \\ &= \sum_{i=1}^n \varepsilon^{i+n} \cdot \langle a_i \rangle \cdot [(\lambda_1 - \lambda_i) a_1, \dots, (\widehat{\lambda_i - \lambda_i}) a_i, \dots, (\lambda_n - \lambda_i) a_n, \lambda_i]. \end{aligned}$$

Proof. We will prove the statement by induction on n . For $n = 1$ this is Lemma 4.3 (1). For $n \geq 2$ we have

$$\begin{aligned}
& \sum_{i=1}^{n-1} \varepsilon^{i+n} \cdot \langle a_i \rangle \cdot [(\lambda_1 - \lambda_i)a_1, \dots, (\widehat{\lambda_i - \lambda_i})a_i, \dots, (\lambda_n - \lambda_i)a_n, \lambda_i] \\
\stackrel{(1)}{=} & \sum_{i=1}^{n-1} \varepsilon^{i+n} \cdot \langle a_i \rangle \cdot [(\lambda_1 - \lambda_i)a_1, \dots, (\widehat{\lambda_i - \lambda_i})a_i, \dots, \lambda_n a_n, \lambda_i] \\
& + \sum_{i=1}^{n-1} \varepsilon^{i+n} \langle a_i \rangle \langle \lambda_n a_n \rangle [(\lambda_1 - \lambda_i)a_1, \dots, \widehat{}, \dots, (\lambda_{n-1} - \lambda_i)a_{n-1}, 1 - \lambda_i/\lambda_n, \lambda_i] \\
\stackrel{(2)}{=} & \left(\sum_{i=1}^{n-1} \varepsilon^{i+n-1} \langle a_i \rangle [(\lambda_1 - \lambda_i)a_1, \dots, (\widehat{\lambda_i - \lambda_i})a_i, \dots, (\lambda_{n-1} - \lambda_i)a_{n-1}, \lambda_i] \right) [\lambda_n a_n] \\
& + \langle \lambda_n a_n \rangle \sum_{i=1}^{n-1} \varepsilon^{i+n} \langle a_i \rangle [(\lambda_1 - \lambda_i)a_1, \dots, \widehat{}, \dots, (\lambda_{n-1} - \lambda_i)a_{n-1}, 1 - \lambda_i/\lambda_n, \lambda_n] \\
\stackrel{(3)}{=} & [\lambda_1 a_1, \dots, \lambda_n a_n] - [a_1, \dots, a_{n-1}, \lambda_n a_n] \\
& + \varepsilon \langle \lambda_n a_n \rangle \sum_{i=1}^{n-1} \varepsilon^{i+n-1} \langle a_i \rangle [(\frac{\lambda_i}{\lambda_n} - \frac{\lambda_1}{\lambda_n})a_1, \dots, \widehat{}, \dots, (\frac{\lambda_i}{\lambda_n} - \frac{\lambda_{n-1}}{\lambda_n})a_{n-1}, 1 - \frac{\lambda_i}{\lambda_n}, \lambda_n] \\
\stackrel{(4)}{=} & [\lambda_1 a_1, \dots, \lambda_n a_n] - [a_1, \dots, a_n] - \langle a_n \rangle [a_1, \dots, a_{n-1}, \lambda_n] \\
& + \varepsilon \langle \lambda_n a_n \rangle \left([(1 - \frac{\lambda_1}{\lambda_n})a_1, \dots, (1 - \frac{\lambda_{n-1}}{\lambda_n})a_{n-1}, \lambda_n] - [a_1, \dots, a_{n-1}, \lambda_n] \right) \\
\stackrel{(5)}{=} & [\lambda_1 a_1, \dots, \lambda_n a_n] - [a_1, \dots, a_n] - \langle a_n \rangle [a_1, \dots, a_{n-1}, \lambda_n] \\
& - \langle a_n \rangle [(\lambda_1 - \lambda_n)a_1, \dots, (\lambda_{n-1} - \lambda_n)a_{n-1}, \lambda_n] + \langle a_n \rangle [a_1, \dots, a_{n-1}, \lambda_n] \\
= & [\lambda_1 a_1, \dots, \lambda_n a_n] - [a_1, \dots, a_n] - \langle a_n \rangle [(\lambda_1 - \lambda_n)a_1, \dots, (\lambda_{n-1} - \lambda_n)a_{n-1}, \lambda_n].
\end{aligned}$$

Here, equation (1) follows from

$$[(\lambda_n - \lambda_i)a_n] = [(1 - \lambda_i/\lambda_n) \cdot \lambda_n a_n] = [\lambda_n a_n] + \langle \lambda_n a_n \rangle [1 - \lambda_i/\lambda_n],$$

equation (2) follows from $[\lambda_i] = [\lambda_n \frac{\lambda_i}{\lambda_n}] = [\lambda_n] + \langle \lambda_n \rangle [\frac{\lambda_i}{\lambda_n}]$ together with the Steinberg relation which yields

$$[1 - \lambda_i/\lambda_n, \lambda_i] = [1 - \lambda_i/\lambda_n, \lambda_n].$$

Equation (3) follows from the induction hypothesis and

$$[-a/\lambda, \lambda] = [a, \lambda] + \langle a \rangle [-1/\lambda, \lambda] = [a, \lambda] + \langle -a/\lambda \rangle [-\lambda, \lambda] = [a, \lambda].$$

Equation (4) follows from the induction hypothesis. Equation (5) follows from $[-a/\lambda, \lambda] = [a, \lambda]$ and $\langle -1 \rangle [a, \lambda] = \langle -1 \rangle [(-\lambda)(-\frac{a}{\lambda})][\lambda] = \langle -1 \rangle ([-\lambda] + \langle -\lambda \rangle [-a/\lambda])[\lambda] = \langle \lambda \rangle [-a/\lambda, \lambda] = \langle \lambda \rangle [a, \lambda]$. \square

5. THE OBSTRUCTION TO FURTHER STABILITY

The purpose of this section is to prove Theorems 1.2 and 1.5 from the Introduction. In subsection 5.1, A can be any commutative ring, unless otherwise stated. In the remaining subsections, A will be a local commutative ring with infinite residue field k . In this case, A has many units, its stable rank is $\text{sr}(A) = 1$, and we write $\bar{a} \in k$ for the reduction of $a \in A$ modulo the maximal ideal in A .

5.1. Multiplicative properties of the spectral sequence. Let A be a commutative ring. For integers $n \geq 0$, we consider complexes $C(A^n)$ where $C_r(A^n)$ is the free abelian group generated by the set $U_r(A^n)$ of left invertible $n \times r$ -matrices with entries in A and differential given by the formula (3.1). So, $C(A^n)$ is concentrated in degrees between 0 and n , and $C_0(A^n) = \mathbb{Z}$ and $C_n(A^n) = \mathbb{Z}[GL_n(A)]$. If A is local, then this complex is the same as the one considered in Section 3. Matrix multiplication makes $C(A^n)$ into a complex of left $GL_n(A)$ -modules.

As in Section 3, we have a spectral sequence

$$(5.1) \quad E_{p,q}^1(A^n) = \mathrm{Tor}_p^{GL_n}(\mathbb{Z}[A^*], C_q(A^n)) \Rightarrow H_{p+q}(\mathbb{Z}[A^*] \overset{L}{\otimes}_{GL_n} C(A^n))$$

with differential d^r of bidegree $(r-1, -r)$ where $\mathbb{Z}[A^*]$ is considered a right $GL_n(A)$ -module via the determinant map $GL_n A \rightarrow A^*$. This is the spectral sequence $E_{p,q}^1(A^n) \Rightarrow G_{p+q}(A^n)$ of the exact couple

$$(5.2) \quad \begin{array}{ccccc} \bigoplus_{p,q} E_{p,q}^r(A^n) & \xrightarrow{k} & \bigoplus_{p,q} D_{p,q}^r(A^n) & \xrightarrow{\rho} & \bigoplus_{p+q} G_{p+q}(A^n) \\ & \searrow j & \downarrow i & \nearrow \rho & \\ & & \bigoplus_{p,q} D_{p,q}^r(A^n)k & & \end{array}$$

where

$$\begin{aligned} D_{p,q}^1(A^n) &= H_{p+q}(\mathbb{Z}[A^*] \overset{L}{\otimes}_{GL_n} C_{\leq q}(A^n)) \\ E_{p,q}^1(A^n) &= H_{p+q}(\mathbb{Z}[A^*] \overset{L}{\otimes}_{GL_n} C_q(A^n)[q]) = H_p(\mathbb{Z}[A^*] \overset{L}{\otimes}_{GL_n} C_q(A^n)) \\ G_{p+q}(A^n) &= H_{p+q}(\mathbb{Z}[A^*] \overset{L}{\otimes}_{GL_n} C(A^n)). \end{aligned}$$

For $r = 1$, the maps i, j, k are the maps of the long exact sequence of homology groups associated with the exact sequence of complexes

$$0 \rightarrow C_{\leq q-1}(A^n) \rightarrow C_{\leq q}(A^n) \rightarrow C_q(A^n)[q] \rightarrow 0$$

and the map ρ is induced by the inclusion $C_{\leq q}(A^n) \subset C(A^n)$. The derived couple is obtained by keeping G and replacing D with $\mathrm{Im}(j)$, E with the homology of $(E, j \circ k)$, and i, j, k, ρ with certain induced maps.

The spectral sequence comes with a filtration of the abutment

$$(5.3) \quad 0 \subset F_{p+q,0}(A^n) \subset F_{p+q-1,1}(A^n) \subset \cdots \subset F_{0,p+q}(A^n) = G_{p+q}(A^n)$$

where $F_{p,q}(A^n)$ is the image of $\rho : D_{p,q}(A^n) \rightarrow G_{p+q}(A^n)$ (that image is independent of r) and an exact sequence

$$(5.4) \quad 0 \rightarrow F_{p+1,q-1}(A^n) \rightarrow F_{p,q}(A^n) \rightarrow E_{p,q}^\infty(A^n) \rightarrow 0.$$

For large r (depending on (p, q)), the map k is zero, $E_{p,q}^r = E_{p,q}^\infty$, and the exact sequence (5.4) is the exact sequence

$$0 \rightarrow D_{p+1,q-1}^r(A^n) \rightarrow D_{p,q}^r(A^n) \rightarrow E_{p,q}^r(A^n) \rightarrow 0.$$

In this section, we are interested in the degree n -part of the spectral sequence $E(A^n)$, and we set

$$S_n(A) = H_n(\mathbb{Z}[A^*] \overset{L}{\otimes}_{GL_n} C(A^n)) \quad (= G_n(A^n)).$$

For instance, we have

$$S_0(A) = \mathbb{Z}[A^*], \quad S_1(A) = I[A^*]$$

since $\mathbb{Z}[GL_0(A)] = \mathbb{Z}$, and $C(A^1)$ is the augmentation complex $\varepsilon : \mathbb{Z}[A^*] \rightarrow \mathbb{Z}$ with \mathbb{Z} placed in degree 0. We will denote by $Z_n(A^n)$ the $GL_n(A)$ -module

$$Z_n(A^n) = \ker(d_n : C_n(A^n) \rightarrow C_{n-1}(A^n)).$$

Since $C(A^n)$ is concentrated in degrees between 0 and n , we have an inclusion of complexes $Z_n(A^n)[n] \rightarrow C(A^n)$ of $GL_n A$ -modules and hence a canonical map

$$(5.5) \quad \mathbb{Z}[A^*] \otimes_{GL_n} Z_n(A^n) \longrightarrow S_n(A).$$

Remark 5.1. If A is a local ring with infinite residue field, the canonical map $Z_n(A^n)[n] \rightarrow C(A^n)$ is a quasi-isomorphism of complexes of $GL_n(A)$ -modules, by Lemma 3.1 or Lemma 5.6 below. In this case, the map (5.5) is therefore an isomorphism and we have

$$(5.6) \quad S_n(A) = \mathbb{Z}[A^*] \otimes_{GL_n} Z_n(A^n) = \begin{cases} \mathbb{Z} \otimes_{SL_n} Z_n(A^n) = SL_n \setminus Z_n(A^n), & n \geq 1 \\ \mathbb{Z}[A^*], & n = 0. \end{cases}$$

Let A be a commutative ring. Since the complex $C(A^n)$ is concentrated in degrees between 0 and n , the spectral sequence $E(A^n)$ has two edge maps. Set

$$(5.7) \quad \mathcal{B}_n(A) = H_n(\mathbb{Z}[A^*] \otimes_{GL_n}^L \mathbb{Z}) = \begin{cases} H_n(SL_n A, \mathbb{Z}), & n \geq 1 \\ \mathbb{Z}[A^*], & n = 0 \end{cases}$$

$$(\quad (= D_{n,0}^1(A^n) = E_{n,0}^1(A^n))).$$

Then the *incoming edge map* is the map $\mathcal{B}_n(A) \rightarrow S_n(A)$ induced by the inclusion $\mathbb{Z} = C_0(A^n) \subset C(A^n)$. The *outgoing edge map* is the map

$$S_n(A) \rightarrow H_n(\mathbb{Z}[A^*] \otimes_{GL_n}^L \mathbb{Z}[GL_n][n]) = \mathbb{Z}[A^*]$$

induced by the projection $C(A^n) \rightarrow C_n(A^n)[n] = \mathbb{Z}[GL_n(A)][n]$.

Remark 5.2. Let A be a local ring with infinite residue field. In the description (5.6) of $S_n(A)$, the outgoing edge map is the determinant map

$$\det : S_n(A) \rightarrow \mathbb{Z}[A^*] : z \mapsto \det(z)$$

sending the class of a cycle $z = \sum_i n_i \cdot [\alpha_i] \in Z_n(A^n)$, $\alpha_i \in GL_n(A)$, $n_i \in \mathbb{Z}$, to its determinant

$$\det(z) = \sum_i n_i \langle \det \alpha_i \rangle \in \mathbb{Z}[A^*].$$

Let A be a commutative ring. We will need some multiplicative properties of the spectral sequence which we explain now. For a group G , denote by $E(G)$ the standard contractible simplicial set with free G -action on the right, and by $\mathbb{Z}E(G)$ the chain complex associated with the free simplicial abelian group generated by $E(G)$. So, $\mathbb{Z}E(G)$ is the standard free $\mathbb{Z}[G]$ -resolution of the trivial G -module \mathbb{Z} with $\mathbb{Z}E_q(G) = \mathbb{Z}[G^{q+1}]$ and differential $d_q = \sum_{i=0}^q (-1)^i \delta^i$ defined by

$\delta^i(g_0, \dots, g_q) = (g_0, \dots, \hat{g}_i, \dots, g_q)$. The group G acts from the right on $\mathbb{Z}E(G)$ by the formula $(g_0, \dots, g_q)g = (g_0g, \dots, g_qg)$. Recall that the shuffle map

$$\nabla : \mathbb{Z}E(G_1) \otimes \mathbb{Z}E(G_2) \rightarrow \mathbb{Z}E(G_1 \times G_2)$$

is a quasi-isomorphism and a lax monoidal transformation and thus makes the usual unit and associativity diagrams commute; see for instance [SS03, §2.2 and 2.3].

Let Λ be a commutative ring which is flat over \mathbb{Z} . Let G be a group equipped with a group homomorphism $G \rightarrow \Lambda^*$ to the group of units of Λ , that is, a ring homomorphism $\mathbb{Z}[G] \rightarrow \Lambda$. We consider Λ as a right G -module via this ring homomorphism. We let G act on $\Lambda E(G) = \Lambda \otimes \mathbb{Z}E(G)$ via the formula $(\lambda \otimes (g_0, \dots, g_q)) \cdot g = \lambda g \otimes (g_0g, \dots, g_qg)$. Then $\Lambda E(G)$ is a free resolution of Λ in the category of $\Lambda[G]$ -modules. Let M be a bounded complex of G -modules. Then $\Lambda \otimes M$ is a bounded complex of $\Lambda[G]$ -modules. Since Λ is flat over \mathbb{Z} , the canonical map

$$\Lambda \otimes_{\Lambda G}^L (\Lambda \otimes M) \longrightarrow \Lambda \otimes_G^L M$$

induced by the isomorphism

$$\Lambda \otimes_{\Lambda G} (\Lambda \otimes M) \xrightarrow{\cong} \Lambda \otimes_G M : \lambda_1 \otimes (\lambda_2 \otimes m) \mapsto \lambda_1 \lambda_2 \otimes m$$

is a quasi-isomorphism. This can be checked using a $\mathbb{Z}[G]$ -projective resolution of M . In particular,

$$\Lambda E(G) \otimes_{\Lambda[G]} (\Lambda \otimes M) \cong \Lambda E(G) \otimes_G M$$

represents $\Lambda \otimes_G^L M$.

Let G_1, G_2 be groups equipped with group homomorphisms $G_1, G_2 \rightarrow \Lambda^*$. Since Λ is commutative, we obtain a group homomorphism $G_1 \times G_2 \rightarrow \Lambda^*$. Let M_1, M_2 be bounded complexes of G_1 and G_2 -modules, respectively. Then $M_1 \otimes M_2$ is a complex of $G_1 \times G_2$ -modules. The cross product

$$\times : \mathrm{Tor}_p^{G_1}(\Lambda, M_1) \otimes_{\Lambda} \mathrm{Tor}_q^{G_2}(\Lambda, M_2) \longrightarrow \mathrm{Tor}_{p+q}^{G_1 \times G_2}(\Lambda, M_1 \otimes M_2)$$

is the map on homology induced by the map of complexes

$$\begin{aligned} & \{\Lambda E(G_1) \otimes_{G_1} M_1\} \otimes_{\Lambda} \{\Lambda E(G_2) \otimes_{G_2} M_2\} \\ & \cong \{\Lambda E(G_1) \otimes_{\Lambda} \Lambda E(G_2)\} \otimes_{G_1 \times G_2} (M_1 \otimes M_2) \\ & \xrightarrow{\nabla \otimes 1} \Lambda E(G_1 \times G_2) \otimes_{G_1 \times G_2} (M_1 \otimes M_2). \end{aligned}$$

Since the shuffle map is unital and associative, so is the cross product.

We apply the previous considerations to $\Lambda = \mathbb{Z}[A^*]$, $G_1 = GL_m(A)$, $G_2 = GL_n(A)$, $M_1 = C(A^m)$, and $M_2 = C(A^n)$. The cross product together with the $GL_m(A) \times GL_n(A)$ -equivariant map of complexes [NS89, §3]

$$(5.8) \quad C(A^m) \otimes_{\mathbb{Z}} C(A^n) \rightarrow C(A^{m+n}) : x \otimes y \mapsto (x, y)$$

given on basis elements by concatenation of sequences, then defines a product

$$S_m(A) \otimes_{A^*} S_n(A) \rightarrow S_{m+n}(A)$$

which makes

$$S(A) = \bigoplus_{n \geq 0} S_n(A)$$

into an associative and unital $\mathbb{Z}[A^*]$ -algebra.

Remark 5.3. Let A be a local ring with infinite residue field. In terms of cycles (5.6), the product is given by

$$\left(\sum_i m_i [\alpha_i] \right) \cdot \left(\sum_j n_j [\beta_j] \right) = \sum_{i,j} m_i n_j \begin{bmatrix} \alpha_i & 0 \\ 0 & \beta_j \end{bmatrix}$$

where $\alpha_i \in GL_m(A)$ and $\beta_j \in GL_n(A)$, $m_i, n_j \in \mathbb{Z}$. In particular, the outgoing edge map $\det : S(A) \rightarrow \mathbb{Z}[A^*]$ is a ring homomorphism.

Let A be a commutative ring. The map of complexes (5.8) restricts to maps

$$C_{\leq r}(A^m) \otimes C_{\leq s}(A^n) \rightarrow C_{\leq r+s}(A^{m+n})$$

which, together with the cross products, define pairings

$$D_{m-r,r}^1(A^m) \otimes_{A^*} D_{n-s,s}^1(A^n) \rightarrow D_{m+n-r-s,r+s}^1(A^{m+n})$$

that are suitably associative and unital. In particular,

$$\mathcal{B}(A) = \bigoplus_{n \geq 0} \mathcal{B}_n(A) = \bigoplus_{n \geq 0} D_{n,0}^1(A^n)$$

is a unital and associative $\mathbb{Z}[A^*]$ -algebra, and for all $r \geq 0$ the graded A^* -module

$$\bigoplus_{n \geq 0} D_{n-r,r}^1(A^n)$$

is a graded $\mathcal{B}(A)$ bimodule. The map $\mathcal{B}(A) \rightarrow S(A)$ is a map of graded $\mathbb{Z}[A^*]$ -algebras, and the maps

$$\mathcal{B}(A) \rightarrow \bigoplus_{n \geq 0} D_{n-r,r}^1(A^n) \twoheadrightarrow \bigoplus_{n \geq 0} F_{n-r,r}(A^n) \hookrightarrow S(A)$$

are $\mathcal{B}(A)$ -bimodule maps. Since all maps in (5.2) are $\mathcal{B}(A)$ -bimodule maps, the resulting spectral sequence

$$\bigoplus_{n \geq 0} E_{p,q}^1(A^n) \Rightarrow \bigoplus_{n \geq 0} G_{p+q}(A^n)$$

is a spectral of $\mathcal{B}(A)$ -bimodules.

For a group G and a subgroup $H \leq G$, write $C(G, H)$ for the complex of G -modules $\mathbb{Z}[G/H] \rightarrow \mathbb{Z} : gH \mapsto 1$ with \mathbb{Z} placed in degree 0. By Shapiro's Lemma, we have a canonical isomorphism $H_n(G, H; \mathbb{Z}) \cong H_n(G, C(G, H))$.

For $n \geq 2$, the inclusion $SL_{n-1}(A) \subset \text{Aff}_{1,n-1}^{SL}(A)$ defines a map of complexes of $SL_n(A)$ -modules

$$(5.9) \quad C(SL_n(A), SL_{n-1}(A)) \rightarrow C(SL_n A, \text{Aff}_{1,n-1}^{SL} A) = C_{\leq 1}(A^n).$$

Lemma 5.4. *Let A be a commutative local ring with infinite residue field. Let $2 \leq n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*).*

(1) *The map (5.9) induces an isomorphism of A^* -modules*

$$H_n(SL_n A, SL_{n-1} A) \xrightarrow{\cong} \sigma^{-1} D_{n-1,1}^1(A).$$

(2) *The natural surjection $D_{n-1,1}^1(A) \rightarrow F_{n-1,1}(A)$ induces an isomorphism of A^* -modules*

$$\sigma^{-1} D_{n-1,1}^1(A) \xrightarrow{\cong} \sigma^{-1} F_{n-1,1}(A).$$

Proof. It follows from Theorem 2.5 and the five lemma that the map (5.9) induces an isomorphism

$$H_p(SL_n(A), SL_{n-1}(A)) \rightarrow \sigma^{-1}H_p(SL_n A, \text{Aff}_{1,n-1}^{SL} A)$$

for all $p \leq n_0$. This proves (1).

For part (2), consider the exact sequence

$$D_{n,0}^1 \xrightarrow{j} D_{n-1,1}^1 \xrightarrow{k} E_{n-1,1}^1 \xrightarrow{i} D_{n-1,0}^1$$

and the isomorphism $k : D_{n-1,0}^1 \cong E_{n-1,0}^1$. By Lemma 3.4, the differential $d = ki : \sigma^{-1}E_{n-1,2}^1 \rightarrow \sigma^{-1}E_{n-1,1}^1$ is zero. Hence, $\sigma^{-1}E_{n-1,1}^2$ is the kernel of the map $i : \sigma^{-1}E_{n-1,1}^1 \rightarrow \sigma^{-1}D_{n-1,0}^1$. The exact sequence above induces the exact sequence $0 \rightarrow \text{Im}(j) \rightarrow D_{n-1,1}^1 \rightarrow \ker(d) \rightarrow 0$ which, after localization at σ , is $0 \rightarrow \sigma^{-1}D_{n,0}^2 \rightarrow \sigma^{-1}D_{n-1,1}^1 \rightarrow \sigma^{-1}E_{n-1,1}^2 \rightarrow 0$. By Proposition 3.5, we have $\sigma^{-1}D_{n,0}^2 = \sigma^{-1}F_{n,0}$ and $\sigma^{-1}E_{n-1,1}^2 = \sigma^{-1}E_{n-1,1}^\infty$. Now, the last exact sequence maps to $0 \rightarrow \sigma^{-1}F_{n,0} \rightarrow \sigma^{-1}F_{n-1,1} \rightarrow \sigma^{-1}E_{n-1,1}^\infty \rightarrow 0$. By the five lemma, we are done. \square

5.2. Presentation and decomposability.

In this subsection, A is a commutative local ring with infinite residue field.

In order to obtain a presentation of $S_n(A)$, we need to recall from [NS89] the definition of the complex of $GL_n(A)$ -modules $\tilde{C}(A^n)$. A sequence (v_1, \dots, v_r) of r vectors in A^n is said to be in general position if any $\min(r, n)$ of the vectors v_1, \dots, v_r span a free submodule of rank $\min(r, n)$. A *rank r general position sequence* in A^n is a sequence (v_1, \dots, v_r) of r vectors in A^n which are in general position. Note that (v_1, \dots, v_r) is in general position in A^n if and only if their reduction $(\bar{v}_1, \dots, \bar{v}_r)$ modulo the maximal ideal of A is in general position in k^n . This is because a set of vectors v_1, \dots, v_s spans a free submodule of rank s if and only if the matrix (v_1, \dots, v_s) has a left inverse.

Let $V = (v_1, \dots, v_r)$ be a general position sequence, we call a vector $w \in A^n$ *transversal to V* if $(V, w) = (v_1, \dots, v_r, w)$ is also in general position.

Lemma 5.5. *Let (A, m, k) be a local ring with infinite residue field k . Let V^1, \dots, V^s be a finite set of rank r general position sequences in A^n . Then there is an element $e \in A^n$ which is transversal to V^1, \dots, V^s .*

Proof. Since a set of vectors is in general position in A^n if and only if it is modulo m , we can assume $A = k$ is an infinite field. Let \mathcal{V} be the union of the vectors occurring in the sequences V^1, \dots, V^s . Let $r_0 = \min(r, n) - 1$. Each subset of \mathcal{V} of cardinality r_0 generates a k -linear subspace of k^n of dimension $\leq r_0$. Since k is infinite and $r_0 < n$ there is $e \in A^n$ which is not in any of these finitely many subspaces. Any such e is transversal to V^1, \dots, V^s . \square

Let $\tilde{U}_r(A^n)$ be the set of sequences (v_1, \dots, v_r) of vectors v_1, \dots, v_r which are in general position in A^n . For integers r, n with $n \geq 0$, let $\tilde{C}_r(A^n) = \mathbb{Z}[\tilde{U}_r^n(A)]$ be the free abelian group with basis the rank r general position sequences (v_1, \dots, v_r) in A^n . For instance, $\tilde{C}_r(A^n) = 0$ for $r < 0$, $\tilde{C}_0(A^n) = \mathbb{Z}$ generated by the empty sequence, and $\tilde{C}_n(A^n) = \mathbb{Z}[GL_n(A)]$. For $i = 1, \dots, r$ one has maps $\delta_r^i : \tilde{C}_r^n \rightarrow \tilde{C}_{r-1}^n$ defined on basis elements by $\delta_r^i(v_1, \dots, v_r) = (v_1, \dots, \hat{v}_i, \dots, v_r)$ omitting the i -th entry. We set $d_r = \sum_{i=1}^r (-1)^{i-1} \delta_r^i : \tilde{C}_r \rightarrow \tilde{C}_{r-1}$, and it is standard that $d_r d_{r+1} = 0$. This

defines the chain complex $\tilde{C}(A^n)$. The group $GL_n(A)$ acts on this complex by left matrix multiplication.

Lemma 5.6. *Let A be a local ring with infinite residue field. Then the complex $\tilde{C}(A^n)$ is acyclic, that is, for all $r \in \mathbb{Z}$ we have*

$$H_r(\tilde{C}(A^n)) = 0.$$

Proof. Let $\xi = \sum_{i=1}^s n_i V^i \in \tilde{C}_r(A^n)$ where V^i are rank r general position sequences. By Lemma 5.5 we can choose $e \in A^n$ which is transversal to V^1, \dots, V^s . Set $(\xi, e) = \sum_{i=1}^s n_i (V^i, e) \in \tilde{C}_{r+1}(A^n)$. If $d_r \xi = 0$ then

$$\begin{aligned} d_{r+1}(\xi, e) &= \sum_{j=1}^{r+1} (-1)^{j-1} \delta_{r+1}^j(\xi, e) = \sum_{j=1}^r (-1)^{j-1} \delta_{r+1}^j(\xi, e) + (-1)^r \delta_{r+1}^{r+1}(\xi, e) \\ &= (d_r \xi, e) + (-1)^r \delta_{r+1}^{r+1}(\xi, e) = (-1)^r \xi. \end{aligned}$$

This shows that ξ is a boundary. \square

In the following proposition, we consider the empty symbol \square as a symbol, the unique symbol of length zero.

Proposition 5.7. *For $n \geq 0$, the $\mathbb{Z}[A^*]$ -module $S_n(A)$ has the following presentation. Generators are the symbols $[a_1, \dots, a_n]$ with $a_i \in A^*$. A system of defining relations has the form*

$$\begin{aligned} &[\lambda_1 a_1, \dots, \lambda_n a_n] - [a_1, \dots, a_n] \\ &= \sum_{i=1}^n \varepsilon^{i+n} \cdot \langle a_i \rangle \cdot [(\lambda_1 - \lambda_i) a_1, \dots, (\widehat{\lambda_i - \lambda_i}) a_i, \dots, (\lambda_n - \lambda_i) a_n, \lambda_i] \end{aligned}$$

where $\lambda_i \in A^*$ and $\bar{\lambda}_i \neq \bar{\lambda}_j \in k$ for $i \neq j$ and $\varepsilon = -\langle -1 \rangle \in \mathbb{Z}[A^*]$.

Proof. For $n = 0$, the module given by the presentation is generated by the empty symbol \square subject to the trivial relation $\square - \square = 0$. Hence, this module is $\mathbb{Z}[A^*]$ which is $S_0(A)$. For $n = 1$, the module given by the presentation is generated by symbols $[a]$ for $a \in A^*$ subject to the relation $[\lambda a] - [a] = \langle a \rangle [\lambda]$ for $a, \lambda \in A^*$. By Lemma 4.1, this module is the augmentation ideal $I[A^*]$ which is $S_1(A)$.

For $n \geq 2$, the proof is the same as in [HT10, Theorem 3.3]. We have

$$S_n(A) = H_0(\mathbb{Z}[A^*] \otimes_{GL_n}^L Z_n(A^n)) = H_0(SL_n, Z_n(A^n))$$

since $\det : GL_n(A) \rightarrow A^*$ is surjective with kernel $SL_n(A)$ (Shapiro's Lemma). In view of Lemma 5.6 we have an exact sequence of $GL_n(A)$ -modules

$$\tilde{C}_{n+2}(A^n) \xrightarrow{d_{n+2}} \tilde{C}_{n+1}(A^n) \rightarrow Z_n(A^n) \rightarrow 0.$$

Taking $SL_n(A)$ -coinvariants yields a presentation of $S_n(A)$ as $A^* = GL_n(A)/SL_n(A)$ -module.

As $GL_n(A)$ -sets we have equalities

$$\tilde{U}_{n+1}(A^n) = \bigsqcup_{a_1, \dots, a_n \in A^*} GL_n(A) \cdot (e_1, \dots, e_n, a)$$

where $a = a_1 e_n + \dots + a_n e_n$ and

$$\tilde{U}_{n+2}(A^n) = \bigsqcup_{\substack{a_1, \dots, a_n \\ b_1, \dots, b_n \in A^*}} GL_n(A) \cdot (e_1, \dots, e_n, a, b)$$

where $a = a_1 e_n + \cdots + a_n e_n$, $b = b_1 e_n + \cdots + b_n e_n$ with $\bar{a}_i \bar{b}_j \neq \bar{a}_j \bar{b}_i \in k$ for $i \neq j$. In other words $b_i = \lambda_i a_i$ for some $\lambda_i \in A^*$ such that $\bar{\lambda}_i \neq \bar{\lambda}_j$.

For $i = 1, \dots, n$ we have

$$d_{n+2}^i(e_1, \dots, e_n, a, b) = (e_1, \dots, \hat{e}_i, \dots, e_n, a, b) = M^i(a) \cdot (e_1, \dots, e_n, c)$$

where $M^i(a)$ is the matrix $(e_1, \dots, \hat{e}_i, \dots, e_n, a)$ and

$$c = M^i(a)^{-1}b = ((\lambda_1 - \lambda_i)a_1, \dots, (\lambda_i - \lambda_i)a_i, \dots, (\lambda_n - \lambda_i)a_n, \lambda_i).$$

Since $\det M^i(a) = (-1)^{n+i}a_i$ we have the following presentation for $S_n(A)$ as $\mathbb{Z}[A^*]$ -module. Generators are the symbols $[a] = [a_1, \dots, a_n]$ with $a = a_1 e_1 + \cdots + a_n e_n$ where $a_i \in A^*$, the symbol $[a]$ representing the $SL_n(A)$ -orbit of (e_1, \dots, e_n, a) . The relations are

$$0 = d_{n+2}(e_1, \dots, e_n, a, b) = (-1)^n [b] - (-1)^n [a] + \sum_{i=1}^n (-1)^{i-1} \langle (-1)^{n+i} a_i \rangle [M^i(a)^{-1}b]$$

where $b_i = \lambda_i a_i$ with $\bar{\lambda}_i \neq \bar{\lambda}_j$ for $i \neq j$. This can be written as

$$[b] - [a] = \sum_{i=1}^n \varepsilon^{i+n} \langle a_i \rangle [M^i(a)^{-1}b].$$

□

Remark 5.8. We compute the determinant of $[a_1, \dots, a_n]$ as

$$\begin{aligned} \det[a_1, \dots, a_n] &= (-1)^n \langle 1 \rangle + \sum_{i=1}^n (-1)^{i+1} \langle \det(e_1, \dots, \hat{e}_i, \dots, e_n, a) \rangle \\ &= (-1)^n \langle 1 \rangle + \sum_{i=1}^n (-1)^{i+1} \langle (-1)^{n+i} a_i \rangle \end{aligned}$$

where $a = a_1 e_1 + \cdots + a_n e_n$. For instance,

$$\begin{aligned} \det[a, b] &= \langle -a \rangle - \langle b \rangle + \langle 1 \rangle \\ \det[a, b, c] &= \langle a \rangle - \langle -b \rangle + \langle c \rangle - \langle 1 \rangle. \end{aligned}$$

Let $S_{\geq 1}(A) = \bigoplus_{n \geq 1} S_n(A) \subset S(A)$ be the ideal of positive degree elements, and define the graded ideal $S(A)^{dec}$ of decomposable elements as

$$S(A)^{dec} = S_{\geq 1}(A) \cdot S_{\geq 1}(A) \subset S(A).$$

The algebra of indecomposable elements is the quotient

$$S(A)^{ind} = S(A)/S(A)^{dec}.$$

We will show in Proposition 5.14 that $\sigma^{-1}S_n^{ind}(A) = 0$ for $n \geq 3$. The proof requires the complexes $C(V, W)$ from [NS89, §3] which we recall now. Let V and W be finitely generated free A -modules. Let $U_m(V, W) = U_m(V) \times W^m \subset U_m(V \oplus W)$ be the set of sequences

$$\begin{pmatrix} v_1 \\ w_1 \end{pmatrix}, \begin{pmatrix} v_2 \\ w_2 \end{pmatrix}, \dots, \begin{pmatrix} v_m \\ w_m \end{pmatrix}$$

with $v_i \in V$ and $w_i \in W$ such that (v_1, v_2, \dots, v_m) can be completed to a basis of V . Let $C_m(V, W) = \mathbb{Z}[U_m(V, W)]$ be the free abelian group generated by $U_m(V, W)$, and define a differential $d : C_m(V, W) \rightarrow C_{m-1}(V, W)$ as in (3.1). In particular, $C(V, W)$ is a subcomplex of $C_m(V \oplus W)$, and $C(V, 0) = C(V)$. Note that

$C_m(V, W) = 0$ for $m > \text{rk } V$ where $\text{rk } V$ denotes the rank of the free A -module V . The group

$$\text{Aff}(V, W) = \begin{pmatrix} GL(V) & 0 \\ \text{Hom}(V, W) & 1_W \end{pmatrix}$$

acts on $U_m(V, W)$ by multiplication from the left making the complex $C(V, W)$ into a complex of left $\text{Aff}(V, W)$ -modules. For $\text{rk } V = n$ let

$$Z_n(V, W) = \ker(d : C_n(V, W) \rightarrow C_{n-1}(V, W)).$$

Lemma 5.9. *If $n = \text{rk } V$, then the canonical map of complexes of $\text{Aff}(V, W)$ -modules*

$$Z_n(V, W)[n] \rightarrow C(V, W)$$

is a quasi-isomorphism.

Proof. This is [NS89, Corollary 3.6]. \square

The determinant map $\text{Aff}(V, W) \rightarrow A^*$ makes $\mathbb{Z}[A^*]$ into a right $\text{Aff}(V, W)$ -module. Write $S(V, W)$ for the group

$$S(V, W) = \text{Tor}_n^{\text{Aff}(V, W)}(\mathbb{Z}[A^*], C(V, W)) = \mathbb{Z}[A^*] \otimes_{\text{Aff}(V, W)} Z_n(V, W).$$

Lemma 5.10. *Let V, W be finitely generated free A -modules with $n = \text{rk } V$. Let $n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then for $0 \leq p \leq n_0$, the inclusion $C(V) = C(V, 0) \subset C(V, W)$ induces isomorphisms*

$$\sigma^{-1} \text{Tor}_p^{GL(V)}(\mathbb{Z}[A^*], C(V)) \xrightarrow{\cong} \sigma^{-1} \text{Tor}_p^{\text{Aff}(V, W)}(\mathbb{Z}[A^*], C(V, W)).$$

In particular, the inclusion $0 \subset W$ induces an isomorphism of A^ -modules*

$$\sigma^{-1} S(V) \xrightarrow{\cong} \sigma^{-1} S(V, W).$$

Proof. The proof is the same as in [NS89, Proposition 3.9] replacing [NS89, Theorem 1.11] with Theorem 2.5. \square

If $n = \text{rk } V$ and $m = \text{rk } W$, then concatenation of sequences defines a map

$$\begin{aligned} Z_n(V, W) \otimes Z_m(W) &\rightarrow Z_{n+m}(V \oplus W) \\ \begin{pmatrix} v_1 & \dots & v_n \\ w_1 & \dots & w_n \end{pmatrix} \otimes (z_1, \dots, z_m) &\mapsto \begin{pmatrix} v_1 & \dots & v_n & 0 & \dots & 0 \\ w_1 & \dots & w_n & z_1 & \dots & z_m \end{pmatrix} \end{aligned}$$

which induces a well-defined multiplication

$$S(V, W) \otimes Z_m(W) \rightarrow S(V \oplus W).$$

Similarly, concatenation defines a product

$$\begin{aligned} Z_m(W) \otimes S(V, W) &\rightarrow S(V \oplus W) \\ (z_1, \dots, z_m) \otimes \begin{pmatrix} v_1 & \dots & v_n \\ w_1 & \dots & w_n \end{pmatrix} &\mapsto \begin{pmatrix} 0 & \dots & 0 & v_1 & \dots & v_n \\ z_1 & \dots & z_m & w_1 & \dots & w_n \end{pmatrix}. \end{aligned}$$

Proposition 5.11. *Let V, W be finitely generated free A -modules with $\text{rk } W = m$. If $\text{rk } V \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*), then the map $p : S(V, W) \rightarrow S(V)$ induced by the unique map of A -modules $W \rightarrow 0$ yields commutative diagrams*

$$\begin{array}{ccc} \sigma^{-1} S(V, W) \otimes Z_m(W) & \longrightarrow & \sigma^{-1} S(V \oplus W) \longleftarrow Z_m(W) \otimes \sigma^{-1} S(V, W) \\ \downarrow p \otimes 1 \cong & \nearrow & \nwarrow \cong \downarrow 1 \otimes p \\ \sigma^{-1} S(V) \otimes Z_m(W) & & Z_m(W) \otimes \sigma^{-1} S(V) \end{array}$$

Proof. This is because the diagrams obviously commutes when the isomorphism $p : \sigma^{-1}S(V, W) \rightarrow \sigma^{-1}S(V)$ is replaced with its inverse given in Lemma 5.10. \square

Lemma 5.12. *Let $n_0 \geq n \geq 2$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then for $a_1, \dots, a_n, b \in A^*$ and $1 \leq i \leq n$ we have in $\sigma^{-1}S(A)^{ind}$*

$$[a_1, \dots, ba_i, \dots, a_n] = \langle b \rangle [a_1, \dots, a_n].$$

Proof. Using Proposition 5.11, the proof is the same as in [HT10, Theorem 6.2] and [NS89, Proposition 3.19]. We omit the details; the case $n = 3$ is explained in more details in Lemma 5.13 below. \square

We will need a slightly more precise version of Lemma 5.12 when $n = 3$.

Lemma 5.13. *Let $n_0 \geq 3$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then for all $a, b, c, \lambda \in A^*$, the following holds in $\sigma^{-1}S_3(A)$ modulo $S_1(A) \cdot S_2(A)$.*

- (1) $[\lambda a, b, c] = \langle \lambda \rangle [a, b, c]$
- (2) $[a, \lambda b, c] = \langle \lambda \rangle [a, b, c] + S_2(A) \cdot [c]$
- (3) $[a, b, \lambda c] = \langle \lambda \rangle [a, b, c] + S_2(A) \cdot [\lambda]$
- (4) $[a, b, c] = \langle abc \rangle [1, 1, 1] + S_2(A) \cdot [c]$.

Proof. To prove (1), write $x = \lambda a e_1 + b e_2 + c e_3$ and note that

$$\begin{aligned} [\lambda a, b, c] - \langle \lambda \rangle [a, b, c] &= d\{(e_1, e_2, e_3, x) - (\lambda e_1, e_2, e_3, x)\} \\ &= (e_1 - \lambda e_1) \cdot d(e_2, e_3, x) \\ &= -[\lambda] \cdot [b, c] \in S_1 \cdot S_2 \end{aligned}$$

where the last equality follows from Proposition 5.11.

To prove (2), write $x = a e_1 + \lambda b e_2 + c e_3$, $u = a e_1 + \lambda b e_2$ and note that

$$\begin{aligned} [a, \lambda b, c] - \langle \lambda \rangle [a, b, c] &= d\{(e_1, e_2, e_3, x) - (e_1, \lambda e_2, e_3, x)\} \\ &= (e_2 - \lambda e_2) \cdot d(u, e_3, x) - d(e_1, u)(e_2 - \lambda e_2) \cdot d(e_3, x) \\ &\quad + \{u(e_2 - \lambda e_2) + (e_2 - \lambda e_2)u\} \cdot d(e_3, x) \\ &\in S_1 \cdot S_2 + S_2 \cdot [c] + S_2 \cdot [c] \end{aligned}$$

where the last line follows from Proposition 5.11.

To prove (3), write $x = a e_1 + b e_2 + \lambda c e_3$ and note that

$$\begin{aligned} [a, b, \lambda c] - \langle \lambda \rangle [a, b, c] &= d\{(e_1, e_2, e_3, x) - (e_1, e_2, \lambda e_3, x)\} \\ &= (e_2 - e_1) \cdot \{x(e_3 - \lambda e_3) + (e_3 - \lambda e_3)x\} \\ &\quad - \{(e_1, e_2) + (e_2 - e_1)x\} \cdot (e_3 - \lambda e_3) \\ &\in S_1 \cdot S_2 + S_2 \cdot [\lambda] \end{aligned}$$

where the last line follows from Proposition 5.11.

Finally, to prove (4), we note that modulo $S_1 \cdot S_2$ we have

$$\begin{aligned} [a, b, c] &= \langle a \rangle [1, b, c] \\ &= \langle ab \rangle [1, 1, c] + S_2 \cdot [c] \\ &= \langle abc \rangle [1, 1, 1] + S_2 \cdot [c] \end{aligned}$$

in view of (1), (2) and (3). \square

The rest of the section is devoted to the proof of the following.

Proposition 5.14. *For $3 \leq n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*) we have*

$$\sigma^{-1}S_n(A)^{dec} = \sigma^{-1}S_n(A).$$

Keep the hypothesis of Proposition 5.14. Let $\Sigma_1(A)$ be the free $\mathbb{Z}[A^*]$ -module generated by the set of units $a \in A^*$ with $\bar{a} \neq 1$. Define $\Sigma(A)$ as the tensor algebra of $\Sigma_1(A)$ over $\mathbb{Z}[A]$ with $\Sigma_1(A)$ placed in degree 1. So, $\Sigma_n(A)$ is the free $\mathbb{Z}[A^*]$ -module generated by symbols $[a_1, \dots, a_n]$ with $a_i \in A^*$ such that $\bar{a}_i \neq 1$, and multiplication is given by concatenation of symbols. Similarly, let $\tilde{\Sigma}_1(A)$ be the free $\mathbb{Z}[A^*]$ -module generated by the set of all units $a \in A^*$. Define $\tilde{\Sigma}(A)$ as the tensor algebra of $\tilde{\Sigma}_1(A)$ over $\mathbb{Z}[A]$ with $\tilde{\Sigma}_1(A)$ placed in degree 1. Consider the diagram of graded $\mathbb{Z}[A^*]$ -module maps

$$(5.10) \quad \begin{array}{ccccc} \Sigma(A) & \xrightarrow{L} & \tilde{\Sigma}_{2*}(A) & \xrightarrow{\Pi} & \mathbb{Z}[A^*][T^2] \\ \parallel & & \downarrow p & & \downarrow q \\ \Sigma(A) & \xrightarrow{R} & S_{2*}(A) & \longrightarrow & S_{2*}(A)^{ind} \end{array}$$

where the maps are defined as follows. The map $L : \Sigma(A) \rightarrow \tilde{\Sigma}_{2*}(A)$ is the $\mathbb{Z}[A^*]$ -algebra map induced by the $\mathbb{Z}[A^*]$ -module homomorphism $\Sigma_1(A) \rightarrow \tilde{\Sigma}_2(A)$ defined on generators $a \in A^*$ of $\Sigma_1(A)$ by

$$L(a) = \langle -1 \rangle [1 - a, 1] - \langle a \rangle [1 - a^{-1}, a^{-1}] + [1, 1].$$

The map $R : \Sigma(A) \rightarrow S_{2*}(A)$ is the $\mathbb{Z}[A^*]$ -algebra homomorphism induced by the $\mathbb{Z}[A^*]$ -module homomorphism

$$\Sigma_1(A) \rightarrow S_2(A) : [a] \mapsto R(a) = [1, a].$$

The map $\Pi : \tilde{\Sigma}_{2*}(A) \rightarrow \mathbb{Z}[A^*][T^2]$ is the $\mathbb{Z}[A^*]$ -algebra homomorphism which is the even part of

$$\tilde{\Sigma}(A) \rightarrow \mathbb{Z}[A^*][T] : [a_1, \dots, a_n] \mapsto \langle a_1 \cdots a_n \rangle T^n.$$

The middle and right vertical maps are the $\mathbb{Z}[A^*]$ -module homomorphisms

$$p : \tilde{\Sigma}_n(A) \rightarrow S_n(A) : [a_1, \dots, a_n] \mapsto [a_1, \dots, a_n]$$

and

$$q : \mathbb{Z}[A^*] \cdot T^n \rightarrow S_n(A)^{ind} : T^n \mapsto [1, \dots, 1].$$

Lemma 5.15. *The diagram (5.10) commutes in degrees $\leq n_0$ after inverting $\sigma \in \mathbb{Z}[A^*]$.*

Proof. Commutativity of the right hand square follows from Lemma 5.12. The proof of the commutativity of the left hand square is the same as in [HT10, Lemma 6.6] and we omit the details. \square

Lemma 5.16. *Proposition 5.14 is true for $n \geq 4$ even.*

Proof. Set $d = n/2$. So $d \geq 2$. The composition of the two lower horizontal maps in diagram (5.10) is zero in degree d since Σ is decomposable in degrees ≥ 2 . The right vertical map in that diagram is surjective in degree d after inverting σ , by Lemma 5.12. For any $r \geq 1$, when extending scalars along $\mathbb{Z}[A^*] \rightarrow \mathbb{Z}[A^*/A^{r*}]$, the

composition of the top two horizontal arrows in the diagram becomes surjective. For if $a \in A$ is a unit with $\bar{a}^r \neq 1$ then $1 - a^{-r} = a^r - 1$ in A^*/A^{r*} , and hence

$$\Pi L(a^r) = \langle -1 \rangle \langle 1 - a^r \rangle - \langle a^r \rangle \langle 1 - a^{-r} \rangle \langle a^{-r} \rangle + 1 = 1 \in \mathbb{Z}[A^*/A^{r*}].$$

It follows that $(\Pi \circ L) \otimes_{A^*} \mathbb{Z}[A^*/A^{r*}]$ is surjective in degree 1 which implies surjectivity in degrees $n_0 \geq n \geq 1$. We have thus shown that

$$(5.11) \quad \sigma^{-1} S_n(A)^{ind} \otimes_{A^*} \mathbb{Z}[A^*/A^{r*}] = 0$$

for all $r \geq 1$.

Taking the image of the filtration (5.3) in $\sigma^{-1} S_n(A)^{ind}$ defines the filtration $0 \subset F_{n,0}^{ind} \subset \dots \subset F_{0,n}^{ind} = \sigma^{-1} S_n(A)^{ind}$ with quotients $E_{p,q}^\infty(A^n)^{ind} = F_{p,q}^{ind}/F_{p+1,q-1}^{ind}$ subquotients of $\sigma^{-1} E_{p,q}^1(A^n)$. Since the edge map $\det : S_n(A) \rightarrow E_{0,n}^\infty = E_{0,n}^1 = \mathbb{Z}[A^*]$ sends the decomposable element $[-1, 1]^d$ to $\det([-1, 1]^d) = (\det[-1, 1])^d = \langle 1 \rangle^d = \langle 1 \rangle$, we have $E_{0,n}^\infty(A^n)^{ind} = 0$. Moreover, $E_{1,n-1}^\infty(A^n)^{ind} = 0$ as $\sigma^{-1} E_{1,n-1}^1(A^n) = 0$. For $2 \leq r \leq n$, the A^* -module $E_{r,n-r}^\infty(A^n)^{ind}$ is a subquotient of $\sigma^{-1} E_{r,n-r}^1(A^n) = H_r(SL_r A)$ and hence a $\mathbb{Z}[A^*/A^{r*}]$ -module. Using (5.11), induction on r shows that $E_{r,n-r}^\infty(A^n)^{ind} = 0$ for all r . Hence, $\sigma^{-1} S_n(A)^{ind} = 0$. \square

Lemma 5.17. *Proposition 5.14 is true for n odd.*

Proof. Let $n = 2d + 1$ be odd with $d \geq 1$. Choose $\lambda_n, \lambda_1, \dots, \lambda_d \in A^*$ such that $\bar{\lambda}_i \neq \bar{\lambda}_j$ for $i \neq j$, $\bar{\lambda}_n \neq \bar{\lambda}_i + \bar{\lambda}_j$ for $i, j = 1, \dots, d$. This is possible since A has infinite residue field. For $i = d + 1, \dots, 2d$ set $\lambda_i = \lambda_n - \lambda_{n-i}$. Then $\lambda_i \in A^*$ and $\bar{\lambda}_i \neq \bar{\lambda}_j$ for $1 \leq i \neq j \leq n$. For $1 \leq i, j < n$ we have $\lambda_j - \lambda_i = -(\lambda_{n-j} - \lambda_{n-i})$. Since n is odd, for $i \neq n$ we therefore find

$$(\lambda_n - \lambda_i) \lambda_i \prod_{j \neq i, n} (\lambda_j - \lambda_i) = -\lambda_{n-i} (\lambda_n - \lambda_{n-i}) \prod_{j \neq n-i, n} (\lambda_j - \lambda_{n-i}).$$

The equation from Proposition 5.7 with $a_1 = \dots = a_n = 1$ together with Lemma 5.12 then implies that $-1 = 0$ in the group $\sigma^{-1} S_n(A)^{ind}$. Hence this group is zero. \square

5.3. The Steinberg relation and $H_2(SL_2 A)$.

In this subsection, A is a commutative local ring with infinite residue field.

Definition 5.18. We define $\bar{S}(A)$ as the quotient left $S(A)$ -module

$$\bar{S}(A) = S(A) / (S(A) \cdot [-1, 1]).$$

Recall the canonical maps

$$(5.12) \quad H_n(SL_n(A), SL_{n-1}(A)) \rightarrow D_{n-1,1}^1(A) \rightarrow F_{n-1,1}(A^{n-1}) \subset S_n(A) \rightarrow \bar{S}_n(A).$$

all but the last of which were defined in Subsection 5.1, and the last is the natural surjection.

Lemma 5.19. *Let $2 \leq n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then the map (5.12) induces isomorphisms of A^* -modules*

$$H_n(SL_n(A), SL_{n-1}(A)) \xrightarrow{\cong} \sigma^{-1} F_{n-1,1}(A^n) \xrightarrow{\cong} \sigma^{-1} \bar{S}_n(A).$$

Proof. The first isomorphism was proved in Lemma 5.4.

For the second isomorphism, recall from the proof of Proposition 3.5 the map of complexes $\psi : C(A^{n-2})[-2] \rightarrow C(A^n)$ which induces maps of short exact sequences

$$\begin{array}{ccccc} \sigma^{-1}F_{n-s+3,s-3}(A^{n-2}) & \hookrightarrow & \sigma^{-1}F_{n-s+2,s-2}(A^{n-2}) & \rightarrow & \sigma^{-1}E_{n-s+2,s-2}^\infty(A^{n-2}) \\ & & \downarrow \psi & & \downarrow \psi \\ \sigma^{-1}F_{n-s+1,s-1}(A^n) & \hookrightarrow & \sigma^{-1}F_{n-s,s}(A^n) & \rightarrow & \sigma^{-1}E_{n-s,s}^\infty(A^n). \end{array}$$

For $s = 2$, the right vertical map is an isomorphism and the upper left corner is 0. It follows that the middle map is injective with cokernel the lower left corner $\sigma^{-1}F_{n-1,1}(A^n)$. Since the right vertical map is an isomorphism for $s \geq 2$, it follows by induction on s that $\psi : \sigma^{-1}F_{n-s+2,s-2}(A^{n-2}) \rightarrow \sigma^{-1}F_{n-s,s}(A^n)$ is injective with cokernel $\sigma^{-1}F_{n-1,1}(A^n)$. The case $s = n$ is the isomorphism $\sigma^{-1}F_{n-1,1}(A^n) \rightarrow \sigma^{-1}S_n/\psi(S_{n-2})$. Since the map ψ is right multiplication with $[-1, 1]$, we have $\psi(S_{n-2}) = S_{n-2} \cdot [-1, 1]$ and we are done. \square

Remark 5.20. [NS89, Remark 3.14]. Let $n \geq 1$. We describe a standard procedure which allows us to represent an arbitrary element in $S_n(A)$ as a sum of generators $[a_1, \dots, a_n]$. Take an arbitrary cycle

$$x = \sum_i n_i(\alpha_i) \in Z_n(A^n)$$

with $n_i \in \mathbb{Z}[A^*]$ and $\alpha_i \in GL_n(A)$ and find a vector $v \in A^n$ in general position with the column vectors of α_i for all i . Then

$$x = (-1)^n d(x, v) = (-1)^n \sum n_i d(\alpha_i, v).$$

We have

$$\langle \alpha_i, v \rangle = \alpha_i \cdot (e_1, \dots, e_n, \alpha_i^{-1}v) \equiv \langle \alpha_i \rangle \cdot (e_1, \dots, e_n, \alpha_i^{-1}v) \pmod{SL_n(A)}$$

where $\langle \alpha_i \rangle \in \mathbb{Z}[A^*]$ denotes the determinant of α_i . Hence,

$$x = \sum_i n_i(\alpha_i) = (-1)^n \sum_i n_i \langle \alpha_i \rangle [\alpha_i^{-1}v].$$

Proposition 5.21. *The map*

$$T_n : S_n(A) \rightarrow \hat{K}_n^{MW}(A) : [a_1, \dots, a_n] \mapsto [a_1, \dots, a_n]$$

defines a map of $\mathbb{Z}[A^]$ -algebras $T : S(A) \rightarrow \hat{K}^{MW}(A)$ sending $S(A) \cdot [-1, 1]$ to zero. In particular, it induces a map of left $S(A)$ -modules*

$$\bar{S}(A) \rightarrow \hat{K}^{MW}(A).$$

Proof. The map $S_n(A) \rightarrow \hat{K}_n^{MW}(A)$ given by the formula in the proposition is a well-defined map of $\mathbb{Z}[A^*]$ -modules in view of Propositions 5.7 and 4.19. In order to check multiplicativity of this map, take $x = \sum_i m_i(\alpha_i) \in Z_m(A^m)$ and $y = \sum_j n_j(\beta_j) \in Z_n(A^n)$ with $m_i, n_j \in \mathbb{Z}[A^*]$ and $\alpha_i \in GL_m(A)$ and $\beta_j \in GL_n(A)$. Choose vectors $v \in A^m$ (and $w \in A^n$) which are in general position with respect to

α_i (and β_j respectively). Then $(v, w) \in A^{m+n}$ is in general position with respect to the frames $\alpha_i \oplus \beta_j = \begin{pmatrix} \alpha_i & 0 \\ 0 & \beta_j \end{pmatrix}$. By Remark 5.20, we have in $S(A)$

$$x \cdot y = \sum m_i n_j (\alpha_i \oplus \beta_j) = \sum m_i n_j \langle \alpha_i \oplus \beta_j \rangle [\alpha_i^{-1} v, \beta_j^{-1} w]$$

whereas

$$\begin{aligned} x &= \sum_i m_i (\alpha_i) = \sum_i m_i \langle \alpha_i \rangle [\alpha_i^{-1} v] \\ y &= \sum_j n_j (\beta_j) = \sum_j n_j \langle \beta_j \rangle [\beta_j^{-1} w]. \end{aligned}$$

This proves multiplicativity. Since $[-1, 1] = 0 \in \hat{K}_2^{MW}$, we are done. \square

Lemma 5.22. *For arbitrary $a_1, \dots, a_n \in A^*$, the following formula holds in $S_n(A)$*

$$[a_1] \cdots [a_n] = \sum_{1 \leq i_1 < \dots < i_k \leq n} (-1)^k \left\langle \prod_{s=1}^k a_{i_s} \right\rangle [a_1, \dots, 1, \dots, 1, \dots, a_n]$$

where the summand $[a_1, \dots, 1, \dots, 1, \dots, a_n]$, corresponding to the index (i_1, \dots, i_k) , is obtained from $[a_1, \dots, a_n]$ by replacing a_{i_s} with 1 for $s = 1, \dots, k$.

Proof. We have $[a] = d(e_1, ae_1) = (ae_1) - (e_1)$. Hence,

$$\begin{aligned} [a_1] \cdots [a_n] &= \prod_{i=1}^n ((a_i e_i) - (e_i)) \\ &= \sum_{1 \leq i_1 < \dots < i_k \leq n} (-1)^{n-k} (e_1, \dots, a_{i_1} e_{i_1}, \dots, a_{i_k} e_{i_k}, \dots, e_n) \end{aligned}$$

The vector $v = a_1 e_1 + \dots + a_n e_n$ is in general position with respect to this cycle. Hence,

$$[a_1] \cdots [a_n] = \sum_{1 \leq i_1 < \dots < i_k \leq n} (-1)^k \langle \alpha_i \rangle [\alpha_{i_1, \dots, i_k}^{-1} v]$$

where α_{i_1, \dots, i_k} is the matrix

$$\alpha_{i_1, \dots, i_k} = (e_1, \dots, a_{i_1} e_{i_1}, \dots, a_{i_k} e_{i_k}, \dots, e_n) \in GL_n(A).$$

Since $v = \alpha_{i_1, \dots, i_k} \cdot (a_1, \dots, 1, \dots, 1, \dots, a_n)^T$ and $\det \alpha_{i_1, \dots, i_k} = a_{i_1} \cdots a_{i_k}$ we are done. \square

Lemma 5.23. *For $\lambda \in A^*$ such that $\bar{\lambda} \neq 1$, the element*

$$s(\lambda) = 1 - \langle 1 - \lambda \rangle - \langle \lambda \rangle \in \mathbb{Z}[A^*]$$

acts as a unit on the A^ -module $H_2(SL_2(A))$.*

Proof. The proof is essentially contained in [Maz05, §2]. Let (a_1, \dots, a_r) be a sequence of units $a_i \in A^*$. Then a sequence (v_1, \dots, v_r) of vectors $v_i \in A^n$ is in general position if and only if the sequence $(a_1 v_1, \dots, a_r v_r)$ is in general position. For $r \geq 1$ we can therefore define the set $\mathbb{P}\tilde{U}_r(A)$ as the quotient of the set $\tilde{U}_r(A)$ by the equivalence relation $(v_1, \dots, v_r) \sim (a_1 v_1, \dots, a_r v_r)$. We define the complex $\mathbb{P}\tilde{C}(A^n)$ by

$$\mathbb{P}\tilde{C}_r(A^n) = \mathbb{Z}[\mathbb{P}\tilde{U}_{r+1}(A^n)]$$

for $r \geq 0$ and $\mathbb{P}\tilde{C}_r(A^n) = 0$ for $r < 0$ (note the shift in degree compared to $\tilde{C}_r(A^n)$). The differential $\mathbb{P}\tilde{C}_r(A^n) \rightarrow \mathbb{P}\tilde{C}_{r-1}(A^n)$ is given by the same formula as for $\tilde{C}_r(A^n)$.

The action of $GL_n(A)$ on A^n makes the complex $\mathbb{P}\tilde{C}(A^n)$ into a complex of $GL_n(A)$ -modules. The unique map $\mathbb{P}\tilde{U}_1 \rightarrow \text{pt}$ defines a map of complexes $\mathbb{P}\tilde{C}(A^n) \rightarrow \mathbb{Z}$ of $GL_n(A)$ -modules where pt is the one-element set. The proof of Lemma 5.6 shows that this map of complexes induces an isomorphism on homology. Hence, for $n \geq 1$ the homology of

$$(5.13) \quad \mathbb{Z} \otimes_{SL_n}^L \mathbb{P}\tilde{C}(A^n) \simeq \mathbb{Z}[A^*] \otimes_{GL_n}^L \mathbb{P}\tilde{C}(A^n)$$

computes the homology of $SL_n(A)$. Let $\mathbb{P}\tilde{C}_{\leq r}(A^n) \subset \mathbb{P}\tilde{C}(A^n)$ be the subcomplex with $\mathbb{P}\tilde{C}_{\leq r}(A^n)_i = \mathbb{P}\tilde{C}_i(A^n)$ for $i \leq r$ and zero otherwise. This defines a filtration on (5.13) by the complexes $\mathbb{Z}[A^*] \otimes_{GL_n}^L \mathbb{P}\tilde{C}_{\leq r}(A^n)$ of $\mathbb{Z}[A^*]$ -modules and thus a spectral sequence of $\mathbb{Z}[A^*]$ -modules

$$E_{p,q}^1(A^n) = H_p(\mathbb{Z}[A^*] \otimes_{GL_n}^L \mathbb{P}\tilde{C}_q(A^n)) \Rightarrow H_{p+q}(SL_n(A))$$

with differentials d^r of bidegree $(r-1, -r)$. For $1 \leq q \leq n$, the group $SL_n(A)$ acts transitively on the set $\mathbb{P}\tilde{U}_q(A^n)$ with stabilizer at (e_{n-q+1}, \dots, e_n) the group

$$\mathbb{P}\text{Aff}_{q,n-q}^{SL}(A) =$$

$$\left\{ \begin{pmatrix} M & 0 \\ N & D \end{pmatrix} \mid M \in GL_{n-q}(A), D \in T_q(A), N \in M_{q,n-q}(A), \det M \det D = 1 \right\}$$

where $(A^*)^q = T_q(A) \subset GL_q(A)$ is the subgroup of diagonal matrices. By [Hut90, Lemma 9] (whose proof works for local rings with infinite residue field), the inclusion of groups

$$\left\{ \begin{pmatrix} M & 0 \\ 0 & D \end{pmatrix} \mid M \in GL_{n-q}(A), D \in T_q(A), \det M \det D = 1 \right\} \subset \mathbb{P}\text{Aff}_{q,n-q}^{SL}(A)$$

induces an isomorphism on integral homology groups. For $n = 2$ and $q = 1, 2$, the left hand side is A^* and thus, its homology has trivial A^* -action. Thus, A^* acts trivially on $E_{p,q}^1(A^2)$ for $q \leq 1$. It follows that A^* acts trivially on $E_{p,q}^\infty(A^2)$ for $q \leq 1$. In particular, the element $s(\lambda)$ acts as -1 , hence as a unit on $E_{p,q}^\infty(A^2)$ for $q \leq 1$. To finish the proof of the lemma, it suffices to show that $s(\lambda)$ acts as a unit on the cokernel of the $\mathbb{Z}[A^*]$ -module map

$$(5.14) \quad d^1 : E_{0,3}^1(A^2) \rightarrow E_{0,2}^1(A^2).$$

As a $GL_2(A)$ -set we have

$$\mathbb{P}\tilde{U}_3(A^2) = GL_2(A)/D_2(A) \cdot (e_1, e_2, e_1 + e_2)$$

where $D_2(A) = A^* \cdot 1$ is the group of invertible scalar matrices. It follows that we have isomorphisms of A^* -modules

$$E_{0,2}^1(A^2) = \mathbb{Z}[A^*] \otimes_{GL_2} \mathbb{Z}[\mathbb{P}\tilde{U}_3(A^2)] \cong \mathbb{Z}[A^*] \otimes_{D_2} \mathbb{Z} \cong \mathbb{Z}[A^*/A^{2*}]$$

where $1 \in \mathbb{Z}[A^*/A^{2*}]$ corresponds to $1 \otimes (e_1, e_2, e_1 + e_2) \in \mathbb{Z}[A^*] \otimes_{GL_2} \mathbb{Z}[\mathbb{P}\tilde{U}_3(A^2)]$. As a $GL_2(A)$ -set we have

$$\mathbb{P}\tilde{U}_4(A^2) = \bigsqcup_{a,b \in A^*, \bar{a} \neq \bar{b}} GL_2(A)/D_2(A) \cdot (e_1, e_2, e_1 + e_2, ae_1 + be_2).$$

The map (5.14) sends the element $(e_1, e_2, e_1 + e_2, ae_1 + be_2)$ to the following element in $E_{0,2}^1(A^2)$

$$\begin{aligned}
& \begin{pmatrix} 0 & 1 & a \\ 1 & 1 & b \end{pmatrix} - \begin{pmatrix} 1 & 1 & a \\ 0 & 1 & b \end{pmatrix} + \begin{pmatrix} 1 & 0 & a \\ 0 & 1 & b \end{pmatrix} - \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \\
&= \langle -1 \rangle \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & a \\ 1 & 1 & b \end{pmatrix} - \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & a \\ 0 & 1 & b \end{pmatrix} + \begin{pmatrix} 1 & 0 & a \\ 0 & 1 & b \end{pmatrix} - \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \\
&= \langle -1 \rangle \begin{pmatrix} 1 & 0 & b-a \\ 0 & 1 & a \end{pmatrix} - \begin{pmatrix} 1 & 0 & a-b \\ 0 & 1 & b \end{pmatrix} + \begin{pmatrix} 1 & 0 & a \\ 0 & 1 & b \end{pmatrix} - \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \\
&= (\langle -(b-a)a \rangle - \langle (a-b)b \rangle + \langle ab \rangle - \langle 1 \rangle) \cdot (e_1, e_2, e_1 + e_2).
\end{aligned}$$

It follows that the cokernel of the map (5.14) is the quotient of $\mathbb{Z}[A^*/A^{2*}]$ modulo the A^* -submodule generated by $\langle (a-b)a \rangle - \langle (a-b)b \rangle + \langle ab \rangle - \langle 1 \rangle$ whenever $a, b \in A^*$ with $\bar{a} \neq \bar{b}$. Setting $a = 1$ and $b = \lambda$, we see that $s(\lambda) = -\langle \lambda(1-\lambda) \rangle$ acts invertibly on the cokernel of (5.14). \square

The following proposition shows that the Steinberg relation holds in $\sigma^{-1}\bar{S}_2(A)$.

Proposition 5.24. *Let $2 \leq n_0$ and let $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then for any $\lambda \in A^*$ such that $\bar{\lambda} \neq 1$ we have in the A^* -module $\sigma^{-1}S_2(A)$ the following equality*

$$[\lambda][1-\lambda] = \langle \langle \lambda \rangle \rangle \langle \langle 1-\lambda \rangle \rangle \cdot [-1, 1].$$

Proof. By Lemma 5.22, we have

$$[\lambda][1-\lambda] = [\lambda, 1-\lambda] - \langle 1-\lambda \rangle [\lambda, 1] - \langle \lambda \rangle [1, 1-\lambda] + \langle \lambda(1-\lambda) \rangle [1, 1].$$

Recall from Proposition 5.7 that for $\alpha, \beta, s, t \in A^*$ with $\bar{\alpha} \neq \bar{\beta}$ we have

$$(5.15) \quad [\alpha s, \beta t] - [s, t] = \langle t \rangle [(\alpha - \beta)s, \beta] - \langle -s \rangle [(\beta - \alpha)t, \alpha].$$

Setting $\alpha = 1 - b\lambda^{-1}$, $\beta = 1$, $s = a\lambda$, $t = b$ (where $\bar{\lambda} \neq \bar{b}$) in (5.15), we obtain

$$(5.16) \quad [a\lambda - ab, b] - [a\lambda, b] = \langle b \rangle [-ab, 1] - \langle -a\lambda \rangle [b^2\lambda^{-1}, 1 - b\lambda^{-1}].$$

Setting $\alpha = \lambda$, $\beta = b$, $s = a$ and $t = 1$ (where $\bar{\lambda} \neq \bar{b}$) in (5.15), we obtain

$$(5.17) \quad [a\lambda, b] - [a, 1] = [a\lambda - ab, b] - \langle -a \rangle [b - \lambda, \lambda].$$

Adding equations (5.16) and (5.17), cancelling common summands and multiplying with $\langle -a^{-1} \rangle$ we obtain

$$(5.18) \quad \langle -a^{-1}b \rangle [-ab, 1] + \langle -a^{-1} \rangle [a, 1] = \langle \lambda \rangle [b^2\lambda^{-1}, 1 - b\lambda^{-1}] + [b - \lambda, \lambda].$$

for any $a, b, \lambda \in A^*$ with $\bar{\lambda} \neq \bar{b}$. Note that the right hand side of (5.18) is independent of $a \in A^*$. Hence, so is the left hand side and thus it equals its evaluation at $a = -1$, that is, we have

$$\langle -a^{-1}b \rangle [-ab, 1] + \langle -a^{-1} \rangle [a, 1] = \langle b \rangle [b, 1] + [-1, 1].$$

Therefore,

$$(5.19) \quad [-ab, 1] = -\langle b^{-1} \rangle [a, 1] + \langle -a \rangle [b, 1] + \langle -ab^{-1} \rangle [-1, 1],$$

for any $a, b \in A^*$ (as we can always choose a $\lambda \in A^*$ with $\bar{\lambda} \neq \bar{b}$). Replacing b with $-b$ in equation (5.18) we see that the expression $\langle a^{-1}b \rangle [ab, 1] + \langle -a^{-1} \rangle [a, 1]$ does not depend on a . In particular, it equals its evaluation at $a = 1$, and we have

$$\langle a^{-1}b \rangle [ab, 1] + \langle -a^{-1} \rangle [a, 1] = \langle b \rangle [b, 1] + \langle -1 \rangle [1, 1],$$

that is,

$$(5.20) \quad [ab, 1] = -\langle -b^{-1} \rangle [a, 1] + \langle a \rangle [b, 1] + \langle -ab^{-1} \rangle [1, 1].$$

For $b = 1$ this yields

$$(5.21) \quad [a, 1] = -\langle -1 \rangle [a, 1] + \langle a \rangle [1, 1] + \langle -a \rangle [1, 1].$$

Setting $s = a$, $\alpha = \lambda$, $t = \beta = 1$ (where $\bar{\lambda} \neq 1$) in (5.15), we obtain

$$[\lambda a, 1] - [a, 1] = [(\lambda - 1)a, 1] - \langle -a \rangle [(1 - \lambda), \lambda]$$

that is,

$$(5.22) \quad \langle -a \rangle [1 - \lambda, \lambda] = [(\lambda - 1)a, 1] - [\lambda a, 1] + [a, 1].$$

Together with equations (5.19) and (5.20) this yields

$$\begin{aligned} & \langle -a \rangle [1 - \lambda, \lambda] \\ = & -\langle (1 - \lambda)^{-1} \rangle [a, 1] + \langle -a \rangle [1 - \lambda, 1] + \langle -a(1 - \lambda)^{-1} \rangle [-1, 1] \\ & + \langle -\lambda^{-1} \rangle [a, 1] - \langle a \rangle [\lambda, 1] - \langle -a\lambda^{-1} \rangle [1, 1] + [a, 1] \\ \stackrel{(5.21)}{=} & (1 - \langle \lambda^{-1} \rangle - \langle (1 - \lambda)^{-1} \rangle) \cdot [a, 1] \\ & + \langle -a \rangle \cdot ([1 - \lambda, 1] + \langle (1 - \lambda)^{-1} \rangle [-1, 1] - \langle -1 \rangle [\lambda, 1] + \langle -\lambda^{-1} \rangle [1, 1]). \end{aligned}$$

It follows that the expression

$$(1 - \langle (1 - \lambda)^{-1} \rangle - \langle \lambda^{-1} \rangle) \cdot \langle -a^{-1} \rangle [a, 1]$$

does not depend on a . In particular,

$$(5.23) \quad \{1 - \langle (1 - \lambda)^{-1} \rangle - \langle \lambda^{-1} \rangle\} \cdot \{\langle -a^{-1} \rangle [a, 1] - [-1, 1]\} = 0 \in S_2(A).$$

Since $\det[a, 1] = \langle -a \rangle$, we have

$$\langle -a^{-1} \rangle [a, 1] - [-1, 1] \in \sigma^{-1} \ker(\det : S_2 \rightarrow \mathbb{Z}[A^*]) = \sigma^{-1} F_{2,0} = H_2(SL_2 A),$$

where the equality $\sigma^{-1} F_{2,0}(A^2) = H_2(SL_2 A)$ follows from Lemma 5.19.

By Lemma 5.23, the first factor in (5.23) is invertible in $H_2(SL_2 A)$ as square units act trivially on that group. Hence, we have $\langle -a^{-1} \rangle [a, 1] - [-1, 1] = 0 \in \sigma^{-1} S_2(A)$ for all $a \in A^*$ and therefore,

$$(5.24) \quad [a, 1] = \langle a \rangle [1, 1] \in \sigma^{-1} S_2(A).$$

Setting $s = t = \alpha = 1$, $\beta = \lambda$ with $\bar{\lambda} \neq 1$ in (5.15) yields

$$[1, \lambda] - [1, 1] = [(1 - \lambda), \lambda] - \langle -1 \rangle [\lambda - 1, 1].$$

Putting this into (5.22) with $a = 1$ yields the equation in $\sigma^{-1} S_2(A)$

$$(5.25) \quad [1, \lambda] = -\langle -1 \rangle [\lambda, 1] + [1, 1] + \langle -1 \rangle [1, 1].$$

Finally, putting $a = -1$ in (5.22) and using (5.25) we find for $\lambda \in A^*$ with $\bar{\lambda} \neq 1$ the equation in $\sigma^{-1}S_2(A)$

$$\begin{aligned}
& [1 - \lambda, \lambda] - \langle \lambda \rangle [1 - \lambda, 1] - \langle 1 - \lambda \rangle [1, \lambda] + \langle (1 - \lambda)\lambda \rangle [1, 1] \\
\stackrel{(5.25)}{=} & [1 - \lambda, 1] + [-1, 1] - [-\lambda, 1] - \langle \lambda \rangle [1 - \lambda, 1] \\
& - \langle 1 - \lambda \rangle (-\langle -1 \rangle [\lambda, 1] + [1, 1] + \langle -1 \rangle [1, 1]) + \langle (1 - \lambda)\lambda \rangle [1, 1] \\
\stackrel{(5.24)}{=} & \langle -1 \rangle (1 - \langle \lambda \rangle + \langle \lambda(1 - \lambda) \rangle - \langle 1 - \lambda \rangle) \cdot [1, 1] \\
= & (1 - \langle \lambda \rangle + \langle \lambda(1 - \lambda) \rangle - \langle 1 - \lambda \rangle) \cdot [-1, 1] \\
= & \langle \langle \lambda \rangle \rangle \langle \langle 1 - \lambda \rangle \rangle \cdot [-1, 1].
\end{aligned}$$

Replacing λ with $1 - \lambda$ yields the desired result. \square

Lemma 5.25. *Let $2 \leq n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then the localization map induces an isomorphism*

$$\hat{K}_n^{MW}(A) \xrightarrow{\cong} \sigma^{-1} \hat{K}_n^{MW}(A).$$

Proof. Since $n_0 \geq 2$, the number t in (*) is even and $\sigma = s_{m,-t}$ acts as 1 on $\hat{K}_n^{MW}(A)$ for $n \geq 2$ since square units act as 1 on it (Lemma 4.4 (6)) and $\varepsilon(\sigma) = 1$. Therefore, $\hat{K}_n^{MW}(A) = \sigma^{-1} \hat{K}_n^{MW}(A)$. \square

Corollary 5.26. *Let $2 \leq n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). The following map is well-defined and an isomorphism of A^* -modules*

$$\hat{K}_2^{MW}(A) \xrightarrow{\cong} \sigma^{-1} \bar{S}_2(A) : [a, b] \mapsto [a] \cdot [b].$$

The inverse isomorphism is the map $T_2 : \sigma^{-1} \bar{S}_2(A) \rightarrow \sigma^{-1} K_2^{MW}(A) = K_2^{MW}(A)$ from Proposition 5.21.

Proof. The map is well-defined, by Proposition 5.24. It is surjective, by Lemma 5.12. It follows from the multiplicativity of the map in Proposition 5.21 that the composition $\hat{K}_2^{MW}(A) \rightarrow \sigma^{-1} \bar{S}_2(A) \rightarrow \hat{K}_2^{MW}(A)$ is the identity. This proves the claim. \square

We have proved the following presentation of $H_2(SL_2A)$. Different presentations were given in [Mat69, Corollaire 5.11], [Moo68, Theorem 9.2], [vdK77, Theorem 3.4].

Theorem 5.27. *Let A be a commutative local ring with infinite residue field. Let $I[A^*] \subset \mathbb{Z}[A^*]$ be the augmentation ideal, and write $[a]$ for $\langle a \rangle - 1 \in I[A^*]$. Then there is an isomorphism of A^* -modules*

$$H_2(SL_2A, \mathbb{Z}) \cong I[A^*] \otimes_{A^*} I[A^*] / \{[a] \otimes [1 - a] \mid a, 1 - a \in A^*\}.$$

Proof. Recall that the right hand side is $\hat{K}_2^{MW}(A)$, by definition. The isomorphism in the theorem is the composition of isomorphisms

$$H_2(SL_2A) \xrightarrow{\cong} \sigma^{-1} F_{1,1}(A^2) \xrightarrow{\cong} \sigma^{-1} \bar{S}_2 \xrightarrow{\cong} \hat{K}_2^{MW}(A)$$

from Lemma 5.19 and Corollary 5.26. \square

Remark 5.28. For any $a, b \in A^*$ we have in $\sigma^{-1}S_2(A)$ the following equality

$$[a, b] = [a] \cdot [b] + (\langle -a \rangle + \langle a \rangle - \langle ab \rangle)[-1, 1].$$

This follows from the isomorphism

$$(\det, T_2) : \sigma^{-1}S_2(A) \xrightarrow{\cong} \sigma^{-1}\mathbb{Z}[A^*] \oplus \hat{K}_2^{MW}(A).$$

5.4. Centrality of $[-1, 1]$ and $H_n(SL_n A, SL_{n-1} A)$.

In this subsection, A is a commutative local ring with infinite residue field.

Definition 5.29. For $\lambda \in A^*$ with $\bar{\lambda} \neq 1$, we define the element $\beta_\lambda \in S_3(A)$ as

$$\begin{aligned} \beta_\lambda &= [1, 1 - \lambda, \lambda] - [1, 1 - \lambda, 1] + [1, -\lambda, 1] \\ &\quad - [1 - \lambda, \lambda, 1] + [1 - \lambda, 1, 1] - [-\lambda, 1, 1]. \end{aligned}$$

Note that $\det \beta_\lambda = 0$, by Remark 5.8.

Lemma 5.30. For all $a, \lambda \in A^*$ such that $\bar{\lambda} \neq 1$ we have in $S_3(A)$ the equality

$$[-1, 1] \cdot [a] - [a] \cdot [-1, 1] = \langle \langle a \rangle \rangle \cdot \beta_\lambda.$$

Moreover, $\beta_\lambda = \beta_\mu$ for all $\lambda, \mu \in A^*$ with $\bar{\lambda}, \bar{\mu} \neq 1$.

Proof. Let u be the element $u = d(e_2, e_3, e_3 - e_2) \in Z_2(Ae_2 + Ae_3)$. We have

$$\begin{aligned} &[-1, 1] \cdot [a] - [a] \cdot [-1, 1] \\ &= d(e_1, e_2, e_2 - e_1) \cdot (ae_3 - e_3) - (ae_1 - e_1) \cdot d(e_2, e_3, e_3 - e_2) \\ &= u \cdot (ae_1 - e_1) - (ae_1 - e_1) \cdot u \\ &= \{(e_1) \cdot u - u \cdot (e_1)\} - \{(ae_1) \cdot u - u \cdot (ae_1)\} \\ &= \langle \langle a \rangle \rangle \{u \cdot (e_1) - (e_1) \cdot u\}. \end{aligned}$$

We need to show that $\beta_\lambda = u \cdot (e_1) - (e_1) \cdot u$. Note that the right hand side is independent of λ . The vector $v = (1, -\lambda, 1)^T$ is in general position with respect to the vectors $e_1, e_2, e_3, e_3 - e_2$ occurring in $u(e_1) - (e_1)u$. Therefore, we obtain the equality $u(e_1) - (e_1)u = d\{(e_1, u, v) - (u, e_1, v)\}$. In $\tilde{C}_4(A^3)/SL_3(A)$ we have

$$\begin{aligned} (e_1, u, v) &= (e_1, e_3, e_3 - e_2, v) - (e_1, e_2, e_3 - e_2, v) + (e_1, e_2, e_3, v) \\ &= (e_1, e_2, e_3, v_1) - (e_1, e_2, e_3, v_2) + (e_1, e_2, e_3, v_3) \end{aligned}$$

where $v_1 = (1, 1 - \lambda, \lambda)^T$, $v_2 = (1, 1 - \lambda, 1)^T$ and $v_3 = v$ since matrix multiplication with $(e_1, e_3, e_3 - e_2), (e_1, e_2, e_3 - e_2) \in SL_3(A)$ yields $v = (e_1, e_3, e_3 - e_2)v_1 = (e_1, e_2, e_3 - e_2)v_2$. Similarly, we have in $\tilde{C}_4(A^3)/SL_3(A)$

$$\begin{aligned} (u, e_1, v) &= (e_3, e_3 - e_2, e_1, v) - (e_2, e_3 - e_2, e_1, v) + (e_2, e_3, e_1, v) \\ &= (e_1, e_2, e_3, v_4) - (e_1, e_2, e_3, v_5) + (e_1, e_2, e_3, v_6) \end{aligned}$$

where $v_4 = (1 - \lambda, \lambda, 1)^T$, $v_5 = (1 - \lambda, 1, 1)^T$ and $v_6 = (-\lambda, 1, 1)^T$ since matrix multiplication with $(e_3, e_3 - e_2, e_1), (e_2, e_3 - e_2, e_1), (e_2, e_3, e_1) \in SL_3(A)$ yields $v = (e_3, e_3 - e_2, e_1)v_4 = (e_2, e_3 - e_2, e_1)v_5 = (e_2, e_3, e_1)v_6$. The result follows. \square

In view of the independence of λ we will write β for β_λ . Write $S_n^0(A)$ for the kernel of the determinant map $\det : S_n(A) \rightarrow \mathbb{Z}[A^*]$.

Lemma 5.31. *Let $n_0 \geq 3$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then*

$$\beta \in \sigma^{-1}S_1(A) \cdot S_2^0(A) \subset \sigma^{-1}S_3(A).$$

Proof. To simplify notation I will write S_n for $\sigma^{-1}S_n(A)$. From the definition of β and Lemma 5.13 (4) we have in S_3 modulo $S_2[\lambda] + S_1S_2$ the equality

$$\beta = \{ \langle (1-\lambda)\lambda \rangle - \langle 1-\lambda \rangle + \langle -\lambda \rangle - \langle (1-\lambda)\lambda \rangle + \langle 1-\lambda \rangle - \langle -\lambda \rangle \} \cdot [1, 1, 1] = 0.$$

Therefore, $\beta \in S_2[\lambda] + S_1S_2$. By Remark 5.28, we have $S_2 = \mathbb{Z}[A^*] \cdot [-1, 1] + S_1S_1$, and hence, $\beta \in \mathbb{Z}[A^*] \cdot [-1, 1][\lambda] + S_1S_2$. Since $\det[-1, 1] = 1$, we have the decomposition $S_2 = S_2^0 + \mathbb{Z}[A^*] \cdot [-1, 1]$, and thus, $\beta = r \cdot [-1, 1][\lambda] + c \cdot [-1, 1]$ modulo $S_1S_2^0$ for some $r \in \mathbb{Z}[A^*]$ and $c \in S_1$ (depending on λ). Since $\det \beta = 0$ we can compare determinants and use $\det(S_1S_2^0) = \det(S_1)\det(S_2^0) = 0$ to find $c = -r[\lambda]$. Hence, for all $\lambda \in A^*$ with $\bar{\lambda} \neq 1$ there is $r \in \mathbb{Z}[A^*]$ such that

$$(5.26) \quad \beta = r \cdot \langle \lambda \rangle \cdot \beta \quad \text{mod } S_1S_2^0.$$

Since $\det \beta = 0$, we have $\beta \in S_3^0 = \sigma^{-1}F_{2,1}(A^3) \cong H_3(SL_3A, SL_2A)$. Since square units act trivially on $H_2(SL_2A)$ and cube units act trivially on $H_3(SL_3A)$, the exact sequence

$$H_3(SL_3A) \rightarrow H_3(SL_3A, SL_2A) \rightarrow H_2(SL_2A)$$

implies that for all $a, b \in A^*$ we have $\langle \langle a^2 \rangle \rangle \langle \langle b^3 \rangle \rangle \cdot H_3(SL_3A, SL_2A) = 0$. In particular, $\langle \langle a^2 \rangle \rangle \langle \langle b^3 \rangle \rangle \cdot \beta = 0$. Now choose $a, b \in A^*$ such that $\bar{a}^2, \bar{b}^3 \neq 1$. This is possible since A has infinite residue field. From (5.26) we infer that

$$\beta = r_1 r_2 \langle \langle a^2 \rangle \rangle \langle \langle b^3 \rangle \rangle \cdot \beta = 0 \quad \text{mod } S_1S_2^0.$$

Hence, $\beta \in S_1S_2^0$. □

Lemma 5.32. *Let $n_0 \geq 3$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then*

$$\beta \in \sigma^{-1}F_{3,0}(A^3).$$

Proof. To simplify, write $F_{p,q}$, $E_{p,q}^s$, $D_{p,q}^s$ and S_n also for their localizations at σ . Since $\det \beta = 0$, we have $\beta \in \ker(\det) = F_{1,2}(A^3) = F_{2,1}(A^3)$. So, we have to show that β is sent to zero under the map (when $n = 3$)

$$(5.27) \quad F_{n-1,1}(A^n) \rightarrow E_{n-1,1}^\infty(A^n) = E_{n-1,1}^2(A^n) \subset E_{n-1,1}^1(A^n)$$

which is well-defined for $n \leq n_0$, by Lemma 3.4 and Proposition 3.5. By Lemma 5.4 (2), this is also the map $D_{n-1,1}^1(A^n) \rightarrow E_{n-1,1}^1(A^n)$. For $n = 1$, this map is the map $I[A^*] \rightarrow \mathbb{Z}[A^*] : [a] \mapsto \langle \langle a \rangle \rangle$. Taking the direct sum over n , the map (5.27) is part of a $\mathcal{B}(A)$ -bimodule map

$$\bigoplus_{0 \leq n \leq n_0} F_{n-1,1}(A^n) \rightarrow \bigoplus_{0 \leq n \leq n_0} E_{n-1,1}^1(A^n) = \bigoplus_{0 \leq n \leq n_0} \sigma^{-1} \text{Tor}_n^{GL_n}(\mathbb{Z}[A^*], C_1(A^n)[1]),$$

by Subsection 5.1. Consider the embedding

$$GL_{n-1}(A) \rightarrow \begin{pmatrix} 1 & * \\ 0 & GL_{n-1}A \end{pmatrix} : M \mapsto \begin{pmatrix} 1 & 0 \\ 0 & M \end{pmatrix}$$

of $GL_{n-1}(A)$ into the stabilizer at e_1 of the $GL_n(A)$ -action on $U_1(A^n)$. By Theorem 2.5, the inclusion induces an isomorphism

$$\bigoplus_{0 \leq n \leq n_0} \sigma^{-1} \operatorname{Tor}_{n-1}^{GL_{n-1}}(\mathbb{Z}[A^*], \mathbb{Z}) \xrightarrow{\cong} \bigoplus_{0 \leq n \leq n_0} \sigma^{-1} \operatorname{Tor}_n^{GL_n}(\mathbb{Z}[A^*], C_1(A^n)[1])$$

of A^* -modules. This map is also a map of right $\mathcal{B}(A)$ -modules. The composition of the $\mathcal{B}(A)$ -bimodule map and the inverse of the right $\mathcal{B}(A)$ -module map defines a right $\mathcal{B}(A)$ -module map

$$\delta = \bigoplus_{0 \leq n \leq n_0} \delta_n : \bigoplus_{0 \leq n \leq n_0} F_{n-1,1}(A^n) \rightarrow \bigoplus_{0 \leq n \leq n_0} \sigma^{-1} \operatorname{Tor}_{n-1}^{GL_{n-1}}(\mathbb{Z}[A^*], \mathbb{Z}).$$

We have to show that $\delta_3(\beta) = 0$.

I claim that the following diagram commutes (all localized at σ):

$$(5.28) \quad \begin{array}{ccccccc} S_1 S_2^0 & \hookrightarrow & F_{2,1} & \hookrightarrow & S_3 & \xrightarrow{T_3} & K_3^{MW}(A) \\ \downarrow & & & & & & \downarrow \eta \\ F_{2,1} & \xrightarrow{\delta_3} & H_2(SL_2 A) & \hookrightarrow & S_2 & \xrightarrow{T_2} & K_2^{MW}(A) \end{array}$$

To see this, note that the A^* -module $S_1 S_2^0$ is generated by products $[a] \cdot \gamma$ where $a \in A^*$ and $\gamma \in S_2^0$. We have $S_2^0 = \mathcal{B}_2(A)$, and thus, $\gamma \in \mathcal{B}_2(A)$. Since δ is a right $\mathcal{B}(A)$ -module morphism and T is a map of rings, we have

$$T_2 \delta_3([a] \cdot \gamma) = T_2(\delta_1([a]) \cdot \gamma) = T_2(\langle\langle a \rangle\rangle \cdot \gamma) = \langle\langle a \rangle\rangle T_2(\gamma)$$

since $\delta_1([a]) = \langle\langle a \rangle\rangle$ as shown above. On the other hand

$$\eta \cdot T_3([a] \cdot \gamma) = \eta \cdot T_1([a]) \cdot T_2(\gamma) = \eta \cdot [a] \cdot T_2(\gamma) = \langle\langle a \rangle\rangle T_2(\gamma).$$

So, the diagram does indeed commute.

Since $[1] = 0 \in \hat{K}_1^{MW}(A)$ and T is multiplicative, the definition of β yields $T_3(\beta) = 0$. By commutativity of diagram (5.28), we have $T_2 \delta_3(\beta) = 0$. By (the proof of) Theorem 5.27, the map $T_2 : H_2(SL_2 A) \rightarrow K_2^{MW}(A)$ is an isomorphism. Hence, $\delta_3(\beta) = 0$. \square

Lemma 5.33. *Let $n_0 \geq 3$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*).*

- (1) A^{2*} acts trivially on $\sigma^{-1} S_1(A) S_2^0(A)$.
- (2) A^* acts trivially on $F_{3,0}(A) \cap \sigma^{-1} S_1(A) S_2^0(A)$.

In particular, for all $a \in A^$, we have*

$$\langle\langle a \rangle\rangle \cdot \beta = 0 \in \sigma^{-1} S_3(A).$$

Proof. Since square units act trivially on $\sigma^{-1} S_2^0 = H_2(SL_2 A)$, this shows part (1). The group of cube units A^{3*} acts trivially on $\sigma^{-1} F_{3,0}(A^3)$ as this group is a quotient of $H_3(SL_3 A)$. Hence, if $\gamma \in F_{3,0}(A) \cap \sigma^{-1} S_1(A) S_2^0(A)$, then $\langle a^2 \rangle \gamma = \gamma = \langle a^3 \rangle \gamma$ and thus, $\langle a \rangle \gamma = \gamma$. This proves part (2).

By Lemmas 5.31 and 5.32, we have $\beta \in F_{3,0}(A) \cap \sigma^{-1} S_1(A) S_2^0(A)$. Hence, for all $a \in A^*$, we have $\langle a \rangle \beta = \beta$, that is, $\langle\langle a \rangle\rangle \cdot \beta = 0$. \square

Proposition 5.34. *Let $n_0 \geq 3$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then for all $a \in A^*$, we have in $\sigma^{-1} S_3(A)$ the equality*

$$[a] \cdot [-1, 1] = [-1, 1] \cdot [a].$$

Proof. This follows from Lemmas 5.30 and 5.33. \square

For a graded ring $R = \bigoplus_{n \geq 0} R_n$, denote by $R_{\leq n}$ the quotient ring which is R_i in degrees $i \leq n$ and 0 otherwise.

Corollary 5.35. *Let $n_0 \geq n \geq 3$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then the element $[-1, 1]$ is central in $\sigma^{-1}S_{\leq n}(A)$. In particular,*

$$\sigma^{-1}\bar{S}_{\leq n} = \sigma^{-1}S_{\leq n}(A)/S_{\leq n-2}(A)[-1, 1]$$

is a quotient algebra of $\sigma^{-1}S(A)$.

Proof. Centrality follows from Proposition 5.34 in view of Proposition 5.14 and Lemma 5.12. The claim follows. \square

Proposition 5.36. *Let $2 \leq n \leq n_0$ and $\sigma \in \mathbb{Z}[A^*]$ as in (*). Then the following map is a well-defined isomorphism of A^* -algebras*

$$\hat{K}_{\leq n}^{MW}(A) \xrightarrow{\cong} \sigma^{-1}\bar{S}_{\leq n}(A) : [a, b] \mapsto [a] \cdot [b].$$

with inverse the map $T_{\leq n} : \sigma^{-1}\bar{S}_{\leq n}(A) \rightarrow \sigma^{-1}K_{\leq n}^{MW}(A)$ from Proposition 5.21.

Proof. The map is well-defined, by Proposition 5.24. It is surjective, by Lemma 5.12 and Proposition 5.14. It follows from the multiplicativity of the map T that the composition $\sigma^{-1}\hat{K}_{\leq n}^{MW}(A) \rightarrow \sigma^{-1}\bar{S}_{\leq n}(A) \rightarrow \sigma^{-1}\hat{K}_{\leq n}^{MW}(A)$ is the identity. This proves the claim. \square

The following proves Theorem 1.5 in view of Theorem 4.18. Recall our convention for $SL_0(A)$ from the Introduction so that $H_*(SL_n A) = H_*(GL_n A, \mathbb{Z}[A^*])$ for $n \geq 0$. We set $SL_n(A) = GL_n(A) = \emptyset$ for $n < 0$.

Theorem 5.37. *Let A be a commutative local ring with infinite residue field.*

- (1) *Then $H_i(SL_n(A), SL_{n-1}(A)) = 0$ for $i < n$ and the maps in Proposition 5.21 and Lemma 5.19 induce isomorphisms of A^* -modules for $n \geq 0$*

$$H_n(SL_n(A), SL_{n-1}(A)) \cong \hat{K}_n^{MW}(A).$$

- (2) *If n is even, then the natural map*

$$H_n(SL_n(A)) \rightarrow H_n(SL_n(A), SL_{n-1}(A))$$

is surjective and inclusion $SL_{n-1}A \subset SL_n A$ induces an isomorphism

$$H_{n-1}(SL_{n-1}A) \cong H_{n-1}(SL_n A).$$

Proof. For $i < n$, the vanishing of homology follows from Theorem 3.9 as $\text{sr}(A) = 1$ and $E_n(A) = SL_n(A)$.

The statement of the theorem is clear for $n \leq 1$. So, assume $n \geq 2$. For $2 \leq n \leq n_0$, the identification with Milnor-Witt K -theory follows from Proposition 5.36 together with Lemma 5.19.

For the second part, assume n is even. We have maps of graded $\mathbb{Z}[A^*]$ -algebras

$$\text{Tens}_{\mathbb{Z}[A^*]} H_2(SL_2(A)) \rightarrow \mathcal{B}(A) = \bigoplus_{n \geq 0} H_n(SL_n(A)) \rightarrow S(A) \xrightarrow{T} \hat{K}^{MW}(A)$$

where $H_2(SL_2(A))$ is placed in degree 2 and $H_0(SL_0 A) = \mathbb{Z}[A^*]$, by our convention for $SL_0(A)$. The composition is an isomorphism in degrees 0 and 2. Since the target ring $\hat{K}^{MW}(A)$ is generated in degree 1, its even part is generated in degree 2, and the composition is surjective in even degrees. The claim follows. \square

Theorem 5.38. *Let A be a commutative local ring with infinite residue field. Then, for $i \geq 0$, the natural homomorphism*

$$\pi_i BGL_{n-1}^+(A) \rightarrow \pi_i BGL_n^+(A)$$

is an isomorphism for $n \geq i + 2$ and surjective for $n \geq i + 1$. Moreover, for $n \geq 2$ there is an exact sequence

$$\pi_n BGL_{n-1}^+(A) \rightarrow \pi_n BGL_n^+(A) \rightarrow K_n^{MW}(A) \rightarrow \pi_{n-1} BGL_{n-1}^+(A) \rightarrow \pi_{n-1} BGL_n^+(A).$$

Proof. The theorem follows from Theorem 5.37 in view of Theorem 3.11. \square

5.5. Prestability.

In this subsection, A is a commutative local ring with infinite residue field.

This subsection is devoted to an explicit computation of the kernel and cokernel of the stabilization map in homology at the edge of stabilization as was done in [HT10] for characteristic zero fields.

Assume that A is a local ring for which the Milnor conjecture on bilinear forms holds, that is, the ring homomorphism defined by Milnor [Mil70] is an isomorphism

$$(5.29) \quad K_*^M(A)/2 \cong \bigoplus_{n \geq 0} I^n(A)/I^{n+1}(A)$$

where $I(A) \subset W(A)$ is the fundamental ideal in the Witt ring of A . By the work of Voevodsky and collaborators [OVV07] and its extension by Kerz [Ker09, Theorem 7.10], the map (5.29) is an isomorphism if A is local and contains an infinite field of characteristic not 2. The map is also an isomorphism for any henselian local ring A with $\frac{1}{2} \in A$ as both sides agree with their value at the residue field of A . Using the isomorphism (5.29) we obtain a commutative diagram

$$(5.30) \quad \begin{array}{ccc} K_n^{MW}(A) & \xrightarrow{\eta^n} & I^n(A) \\ \downarrow & & \downarrow \\ K_n^M(A) & \twoheadrightarrow & K_n^M(A)/2. \end{array}$$

In the following theorem we will assume this diagram to be cartesian. By [Mor04], this is the case for fields whose characteristic is different from 2. This was generalized in [GSZ16] to commutative local rings containing an infinite field of characteristic different from 2. So, our theorem holds in this case.

Theorem 5.39. *Let A be a commutative local ring with infinite residue field. Assume that the map (5.29) is an isomorphism and that the diagram (5.30) is cartesian for all $n \geq 0$. Then for $n \geq 3$ odd we have exact sequences*

$$H_n(SL_{n-1}A) \rightarrow H_n(SL_nA) \rightarrow 2K_n^M(A) \rightarrow 0,$$

$$0 \rightarrow I^n(A) \rightarrow H_{n-1}(SL_{n-1}A) \rightarrow H_{n-1}(SL_nA) \rightarrow 0.$$

The proof requires the following lemma.

Lemma 5.40. *Let A be a local ring with infinite residue field for which (5.29) is an isomorphism and the square (5.30) is cartesian for all $n \geq 0$. Then the following hold.*

(1) The following sequence is exact

$$K_n^{MW}(A) \xrightarrow{h_n} K_n^{MW}(A) \xrightarrow{\eta_n} K_{n-1}^{MW}(A) \rightarrow K_{n-1}^M(A) \rightarrow 0$$

where h_n and η_n are multiplication with $h = 1 + \langle -1 \rangle$ and η , respectively.

(2) For $n \geq 3$ odd, under the isomorphism of Theorem 5.37, the boundary map

$$\partial_n : H_n(SL_n A, SL_{n-1} A) \rightarrow H_{n-1}(SL_{n-1} A, SL_{n-2} A)$$

of the triple $(SL_n A, SL_{n-1} A, SL_{n-2} A)$ is multiplication with η .

(3) The following square is bicartesian for $n \geq 3$ odd

$$\begin{array}{ccc} H_{n-1}(SL_{n-1} A) & \longrightarrow & H_{n-1}(SL_n A) \\ \downarrow & & \downarrow \\ H_{n-1}(SL_{n-1} A, SL_{n-2} A) & \longrightarrow & H_{n-1}(SL_n A, SL_{n-2} A). \end{array}$$

Proof. The sequence in (1) is exact since it is isomorphic to the exact sequence

$$(5.31) \quad I^n \times_{k_n^M} K_n^M \xrightarrow{(0,2)} I^n \times_{k_n^M} K_n^M \xrightarrow{(1,0)} I^{n-1} \times_{k_{n-1}^M} K_{n-1}^M \longrightarrow K_{n-1}^M \rightarrow 0$$

in view of the cartesian square (5.30).

We prove (2). Under the isomorphism $\hat{K}_n^{MW}(A) \cong K_n^{MW}(A)$ for $n \geq 2$ proved in Theorem 4.18, multiplication by $\eta \in K_{-1}^{MW}(A)$ corresponds to the map

$$\eta_n : \hat{K}_n^{MW}(A) \rightarrow \hat{K}_{n-1}^{MW}(A) : [a_1, \dots, a_n] \mapsto \langle \langle a_n \rangle \rangle [a_1, \dots, a_{n-1}].$$

We will show that for all odd $n \geq 1$ the map in (2) is η_n . This is clear for $n = 1$. The map

$$(5.32) \quad \mathcal{B}(A) = \bigoplus_{n \geq 0} H_n(SL_n A) \rightarrow \bigoplus_{n \geq 0} \hat{K}_n^{MW}(A)$$

is a $\mathbb{Z}[A^*]$ -algebra homomorphism which is surjective in even degrees (Theorem 5.37). Moreover, the maps in (2) assemble to a map of left \mathcal{B} -modules

$$\partial : \bigoplus_{n \geq 0} H_n(SL_n A, SL_{n-1} A) \rightarrow \bigoplus_{n \geq 0} H_n(SL_n(A), SL_{n-1} A) \cong \hat{K}^{MW}(A).$$

For $n \geq 3$ odd and $[a_1, \dots, a_n] \in H_n(SL_n A, SL_{n-1} A) = \hat{K}_n^{MW}(A)$ choose a lift $b \in \mathcal{B}_{n-1}(A)$ of $[a_1, \dots, a_{n-1}]$ for the map (5.32). Then

$$\partial_n([a_1, \dots, a_n]) = b \cdot \partial_1([a_n]) = b \cdot \langle \langle a_n \rangle \rangle = \langle \langle a_n \rangle \rangle [a_1, \dots, a_{n-1}].$$

This proves (2).

We prove (3). The horizontal maps in diagram (3) are surjective because we have $H_{n-1}(SL_n A, SL_{n-1} A) = 0$. The left vertical map is surjective, by Theorem 5.37. It follows that the right vertical map is also surjective. The total complex of the square in (3) is part of a Mayer-Vietoris type long exact sequence with boundary map the composition

$$H_n(SL_n A, SL_{n-2} A) \xrightarrow{\alpha_n} H_n(SL_n A, SL_{n-1} A) \xrightarrow{\delta_n} H_{n-1}(SL_{n-1} A).$$

Thus, the square in (3) is bicartesian if and only if $\delta_n \alpha_n = 0$. From (1) and (2) we have $\text{Im}(\alpha_n) = \ker(\eta_n) = \text{Im}(h_n)$. Hence, the square in (3) is bicartesian if and

only if $\delta_n h_n = 0$. For $n = 3$, the square in (3) is bicartesian since the vertical maps are isomorphisms. In particular, $\delta_3 h_3 = 0$. Now, the map

$$\delta : \bigoplus_{n \geq 0} H_n(SL_n(A), SL_{n-1}A) \rightarrow \bigoplus_{n \geq 0} H_n(SL_n A)$$

is a left $\mathcal{B}(A)$ -module map. Take $[a_1, \dots, a_n] \in K_n^{MW}(A) = H_n(SL_n A, SL_{n-1}A)$ where $n \geq 5$ is odd. Then the element $[a_1, \dots, a_{n-3}] \in \hat{K}_{n-3}^{MW}(A)$ lifts to $b \in \mathcal{B}_{n-3}(A)$ and

$$\delta_n(h \cdot [a_1, \dots, a_n]) = b \cdot \delta_3 h_3([a_{n-2}, a_{n-1}, a_n]) = b \cdot 0 = 0$$

□

Proof of Theorem 5.39. From Lemma 5.40 we have (using the notation of that lemma)

$$\ker(H_{n-1}(SL_{n-1}A) \rightarrow H_{n-1}(SL_n A)) = \text{Im}(\partial_n) = \text{Im}(\eta_n) = I^n(A)$$

and

$$\text{coker}(H_n(SL_{n-1}A) \rightarrow H_n(SL_n A)) = \ker(\partial_n) = \ker(\eta_n) = \text{Im}(h_n) = 2K_n^M(A)$$

where the last equality follows because of the exact sequence of Lemma 5.40 (1) being isomorphic to (5.31). □

6. EULER CLASS GROUPS

Let X be a separated noetherian scheme, and denote by Open_X the category of Zariski open subsets of X and inclusions thereof as morphisms. For a simplicial presheaf $F : \text{Open}_X^{\text{op}} \rightarrow \text{sSets}$ on the small Zariski site of X , we denote by

$$[X, F]_{\text{Zar}} = \pi_0(F_{\text{Zar}}(X))$$

the set of maps from X to F in the homotopy category of simplicial presheaves on X for the Zariski-topology [BG73]. This is the set of path components of the simplicial set $F_{\text{Zar}}(X)$ where $F \rightarrow F_{\text{Zar}}$ is a map of simplicial presheaves which induces a weak equivalence of simplicial sets $F_x \rightarrow (F_{\text{Zar}})_x$ on all stalks, $x \in X$, and F_{Zar} is object-wise weakly equivalent to its fibrant model in the Zariski topology [BG73]. The latter means that $F_{\text{Zar}}(\emptyset)$ is contractible and F_{Zar} sends a square

$$(6.1) \quad \begin{array}{ccc} U \cup V & \leftarrow & U \\ \uparrow & & \uparrow \\ V & \leftarrow & U \cap V \end{array}$$

of inclusions of open subsets of X to a homotopy cartesian square of simplicial sets; see [BG73, Theorem 4].

Example 6.1. For the presheaf BGL_n defined by $U \mapsto BGL_n(\Gamma(U, O_X))$ write $B_{\text{Zar}}GL_n$ for $(BGL_n)_{\text{Zar}}$. Then $B_{\text{Zar}}GL_n(X)$ is a (functorial) model of the classifying space $B\text{Vect}_n(X)$ of the category $\text{Vect}_n(X)$ of rank n -vector bundles on X with isomorphisms as morphisms. For the inclusion of the automorphisms of O_X^n into $\text{Vect}_n(X)$ induces a map of simplicial presheaves $BGL_n \rightarrow B\text{Vect}_n$ which is a weak equivalence at the stalks of X as vector bundles over local rings are free. Moreover, $B_{\text{Zar}}GL_n = B\text{Vect}_n$ sends the squares (6.1) to homotopy cartesian squares, by an application of Quillen's Theorem B [Qui73], for instance. Hence,

$$\Phi_n(X) = [X, BGL_n]_{\text{Zar}}$$

is the set $\pi_0 B \text{Vect}_n(X)$ of isomorphism classes of rank n vector bundles on X .

Example 6.2. Similar to Example 6.1, the simplicial presheaf $B SL_n$ defined by $U \mapsto B SL_n(\Gamma(U, O_X))$ has a model $B_{Zar} SL_n$ where $B_{Zar} SL_n(X)$ is the classifying space $B \text{Vect}_n^+(X)$ of the category $\text{Vect}_n^+(X)$ of oriented rank n -vector bundles on X with isomorphisms as morphisms. Here, an oriented vector bundle of rank n is a pair (V, ω) consisting of a vector bundle V of rank n and an isomorphism $\omega : \Lambda^n V \cong O_X$ of line bundles called orientation. Morphisms of oriented vector bundles are isomorphisms of vector bundles preserving the orientation. So, the set

$$\Phi_n^+(X) = [X, B SL_n]_{Zar}$$

is the set $\pi_0 B \text{Vect}_n^+(X)$ of isomorphism classes of rank n oriented vector bundles on X .

Example 6.3. Let $n \geq 2$ be an integer, and let F be a pointed simplicial presheaf such that $\pi_i(F_x) = 0$ for $i \neq n$ and $x \in X$ where F_x denotes the stalk of F at $x \in X$. Then there is a natural bijection of pointed sets

$$[X, F]_{Zar} \cong H_{Zar}^n(X, \tilde{\pi}_n F)$$

where the right hand side denotes Zariski cohomology of X with coefficients in the sheaf of abelian groups $\tilde{\pi}_n F$ associated with the presheaf $U \mapsto \pi_n(F(U))$ [BG73, Propositions 2 and 3].

Denote by $\mathcal{H}_n(SL_n, SL_{n-1})$ the Zariski sheaf associated to the presheaf

$$U \mapsto H_n(SL_n(\Gamma(U, O_X)), SL_{n-1}(\Gamma(U, O_X))).$$

Similarly, denote by \mathcal{K}_n^{MW} the sheaf associated to the presheaf $U \mapsto K_n^{MW}(\Gamma(U, O_X))$.

Lemma 6.4. *Let X be a scheme with infinite residue fields. Then for $n \geq 2$ there is an isomorphism of sheaves of abelian groups on X*

$$\mathcal{H}_n(SL_n, SL_{n-1}) \cong \mathcal{K}_n^{MW}.$$

Proof. Let A be a commutative ring. Recall from §5.1 the graded $\mathbb{Z}[A^*]$ -algebra $S(A) = \bigoplus_{n \geq 0} S_n(A)$. It has $S_0(A) = \mathbb{Z}[A^*]$ and $S_1(A) = I[A^*]$ which allows us to define the map of graded $\mathbb{Z}[A^*]$ -algebras

$$\text{Tens}_{\mathbb{Z}[A^*]} I[A^*] \rightarrow S(A)$$

induced by the identity in degrees 0 and 1. Set

$$H_n(SL_{n/n-1}A) = \text{Tor}_n^{GL_n}(\mathbb{Z}[A^*], C_{\leq 1}(A^n)).$$

Then the inclusion $C_{\leq 1}(A^n) \rightarrow C(A^n)$ of complexes induces maps

$$H_n(SL_{n/n-1}A) \rightarrow S_n(A)$$

which are isomorphisms for $n = 0, 1$, and for $n \geq 2$, we have the identification $H_n(SL_{n/n-1}A) = H_n(SL_n A, SL_{n-1}A)$. Denote by $[-1, 1] \in S_2(A)$ the image of

$$(e_2, e_2 - e_1) - (e_1, e_2 - e_1) + (e_1, e_2) \in Z_2(A^2)/SL_2(A) = \mathbb{Z}[A^*] \otimes_{GL_2(A)} Z_2(A^2)$$

under the map (5.5). For a local ring, this is precisely the element that was denoted $[-1, 1]$ previously. Let $\bar{S}(A)$ be the graded $\mathbb{Z}[A^*]$ -algebra $S(A)$ modulo the two-sided ideal generated by $[-1, 1]$ and $[a][1 - a]$ for $a, 1 - a \in A^*$. The maps above induce a diagram of presheaves

$$H_n(SL_{n/n-1}) \rightarrow \bar{S}_n \leftarrow \hat{K}_n^{MW}$$

whose sheafification we denote by

$$(6.2) \quad \mathcal{H}_n(SL_{n/n-1}) \rightarrow \bar{\mathcal{F}}_n \leftarrow \hat{\mathcal{K}}_n^{MW}.$$

Assume now that $n \geq 2$ and let A be a local ring with infinite residue field. Then there is $\sigma \in \mathbb{Z}[A^*]$ as in (*) such that the localization map $\bar{S}_n(A) \rightarrow \sigma^{-1}\bar{S}_n(A)$ induces the diagram

$$(6.3) \quad \begin{array}{ccccc} H_n(SL_{n/n-1}A) & \longrightarrow & \bar{S}_n(A) & \longleftarrow & \hat{K}_n^{MW}(A) \\ & \searrow \cong & \downarrow & \swarrow \cong & \\ & & \sigma^{-1}\bar{S}_n(A) & & \end{array}$$

where the diagonal arrows are isomorphisms, by Lemma 5.19, Proposition 5.36, Corollary 5.35 and Proposition 5.24. In particular, the horizontal maps in (6.3) and (6.2) are injective, and the middle vertical map in (6.3) is surjective with σ -torsion kernel. From (the proof of) Lemma 5.25, σ acts on $\hat{K}_n^{MW}(A)$ as the identity for $n \geq 2$. This implies that any map from $\hat{K}_n^{MW}(A)$ to a σ -torsion module is zero. In particular, the composition

$$\hat{\mathcal{K}}_n^{MW} \rightarrow \bar{\mathcal{F}}_n \rightarrow \bar{\mathcal{F}}_n/\mathcal{H}_n(SL_{n/n-1})$$

is the zero map and we obtain an induced map of sheaves

$$\hat{\mathcal{K}}_n^{MW} \rightarrow \mathcal{H}_n(SL_{n/n-1})$$

which is an isomorphism, by diagram (6.3). Finally, by Theorem 4.18, the natural surjection of sheaves of graded algebras $\hat{\mathcal{K}}^{MW} \rightarrow \mathcal{K}^{MW} : [a] \mapsto [a]$ is an isomorphism in degrees ≥ 2 . \square

For a commutative ring R , we denote by ΔR the standard simplicial ring

$$n \mapsto \Delta_n R = R[T_0, \dots, T_n]/(-1 + T_0 + \dots + T_n).$$

For an integer $n \geq 0$, let $\tilde{E}_n(R)$ be the maximal perfect subgroup of the kernel of $GL_n(R) \rightarrow \pi_0 GL_n(\Delta R)$. Note that $\tilde{E}_n(R)$ is in fact a subgroup of $SL_n(R)$ since it maps to zero in the commutative group $GL_n(R)/SL_n(R) = R^*$.

Lemma 6.5. *Let $X = \text{Spec } R$, and $x \in X$. Let $n \geq 1$ be an integer. If $n \neq 2$ or $n = 2$ and the residue field $k(x)$ of x has more than 3 elements, then the inclusion $\tilde{E}_n \subset SL_n$ of presheaves induces an isomorphism on stalks at x*

$$(\tilde{E}_n)_x \cong SL_n(O_{X,x}).$$

Proof. The statement is trivial for $n = 1$. So we assume $n \geq 2$. Every elementary matrix $e_{i,j}(r) \in GL_n(R)$, $r \in R$, is the evaluation at $T = 1$ of the elementary matrix $e_{i,j}(Tr) \in GL_n(R[T])$ whereas the evaluation at $T = 0$ yields 1. Therefore, on fundamental groups the natural map $BGL_n(R) \rightarrow BGL_n(\Delta R)$ sends $E_n(R)$ to 1.

If $n \geq 3$, the group $E_n(R)$ is perfect. If $n = 2$, our hypothesis implies that there is $f \in R$ such that $0 \neq f \in k(x)$ and R_f has a unit u such that $u + 1$ and $u - 1$ are also units. Replacing R with R_f , we can assume that R and any localization A of R has this property. For such rings A the group $E_2(A)$ is perfect since for all $a \in A$ we have

$$\left[\begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix}, \begin{pmatrix} 1 & (1-u^2)^{-1}a \\ 0 & 1 \end{pmatrix} \right] = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix},$$

and $\begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix}$ is a product of elementary matrices.

In any case, we can assume $E_n(A)$ perfect and contained in $\tilde{E}_n(A)$ for all localizations A of R . From the inclusions $E_n(A) \subset \tilde{E}_n(A) \subset SL_n(A)$ we obtain the inclusions of corresponding stalks $(E_n)_x \subset (\tilde{E}_n)_x \subset (SL_n)_x = SL_n(O_{X,x})$. Since the composition is an isomorphism, the lemma follows. \square

We will write $BGL_n^+(R)$ and $BSL_n^+(R)$ for a functorial version of Quillen's plus-construction applied to $BGL_n(R)$ and $BSL_n(R)$ with respect to the perfect normal subgroup $\tilde{E}_n(R)$. See [BK72, VII §6] for how to make the plus-construction functorial. The canonical inclusion $GL_{n-1}(R) \subset GL_n(R)$ induces maps $BGL_{n-1}^+ \rightarrow BGL_n^+$ and $BSL_{n-1}^+ \rightarrow BSL_n^+$.

Recall from [May67, §8] that for every integer $n \geq 0$ there is an endofunctor $P_{\leq n} : \mathbf{sSets} \rightarrow \mathbf{sSets}$ of the category of simplicial sets together with natural transformations $S \rightarrow P_{\leq n}S$ such that for every choice of base point $x \in S_0$, the map $\pi_i(S, x) \rightarrow \pi_i(P_{\leq n}S, x)$ is an isomorphism for $i \leq n$ and $\pi_i(P_{\leq n}S, x) = 0$ for $i > n$. Moreover, the map $S \rightarrow P_{\leq n}S$ factors naturally as $S \rightarrow P_{\leq n+1}S \rightarrow P_{\leq n}S$. If F is a simplicial presheaf then $P_{\leq n}F$ denotes the presheaf $U \mapsto P_{\leq n}(F(U))$. For a pointed simplicial presheaf F we denote by $\tilde{\pi}_i F$ the Zariski sheaf associated with the presheaf $U \mapsto \pi_i(F(U), x_0)$ where x_0 is the base point of F . We consider the quotient B/A of an inclusion of simplicial sets $A \subset B$ and $P_{\leq n}(B/A)$ pointed at $\{A\}$.

Lemma 6.6. *Let X be a scheme with infinite residue fields. Then for $n \geq 2$ there are isomorphisms of Zariski sheaves on X*

$$\tilde{\pi}_i P_{\leq n}(BSL_n^+/BSL_{n-1}^+) \cong \begin{cases} 0 & i \neq n \\ \mathcal{K}_n^{MW} & i = n. \end{cases}$$

Proof. From the properties of $P_{\leq n}$, the statement is clear for $i > n$. So, assume $i \leq n$. We have the following string of sheaves

$$\tilde{\pi}_i P_{\leq n}(BSL_n^+/BSL_{n-1}^+) \xleftarrow{\cong} \tilde{\pi}_i(BSL_n^+/BSL_{n-1}^+) \rightarrow \mathcal{H}_i(SL_n, SL_{n-1})$$

where the right arrow is the Hurewicz homomorphism which is an isomorphism for $i \leq n$, by Theorem 5.37 and the fact that BSL_n^+R/BSL_{n-1}^+R is simply connected for local rings R (with infinite residue field). Using Lemma 6.4, the claim follows. \square

Corollary 6.7. *Let X be a noetherian separated scheme with infinite residue fields. Then for $n \geq 2$, there is a natural bijection of pointed sets*

$$[X, P_{\leq n}(BSL_n^+/BSL_{n-1}^+)]_{Zar} \cong H_{Zar}^n(X, \mathcal{K}_n^{MW}).$$

Proof. This is Example 6.3 and Lemma 6.6. \square

Definition 6.8. The *Euler class map* (for rank n oriented vector bundles) is the composition of maps of simplicial presheaves

$$e : BSL_n^+ \rightarrow BSL_n^+/BSL_{n-1}^+ \rightarrow P_{\leq n}(BSL_n^+/BSL_{n-1}^+).$$

By definition, it is trivial when restricted to BSL_{n-1}^+ .

Let $n \geq 2$. In view of Corollary 6.7, applying the functor $F \mapsto [X, F]_{Zar}$ to the sequence

$$BSL_{n-1}^+ \rightarrow BSL_n^+ \xrightarrow{e} P_{\leq n}(BSL_n^+/BSL_{n-1}^+)$$

yields the sequence

$$(6.4) \quad [X, BSL_{n-1}^+]_{Zar} \longrightarrow [X, BSL_n^+]_{Zar} \xrightarrow{\epsilon} H_{Zar}^n(X, \mathcal{K}_n^{MW}).$$

A sequence $U \rightarrow V \rightarrow W$ of sets with W pointed is called *exact* if every element of V which is sent to the base point in W comes from U .

Theorem 6.9. *Let $n \geq 2$ be an integer and let X be a noetherian separated scheme with infinite residue fields. Assume that the dimension of X is at most n . Then the sequence of sets (6.4) is exact.*

Proof. This follows from obstruction theory [Mor12, Corollary B.10] in view of Lemma 6.6. \square

One would like to replace BSL_r^+ with BSL_r for $r = n - 1, n$ in Theorem 6.9. This motivates the following.

Question 6.10. For which affine noetherian scheme X is the canonical map

$$[X, BGL_n]_{Zar} \rightarrow [X, BGL_n^+]_{Zar}$$

a bijection?

We will prove in Corollary 6.16 below that $[X, BGL_n]_{Zar}$ is a natural direct factor of $[X, BGL_n^+]_{Zar}$. This will suffice to prove that (6.4) is still exact when BSL_r^+ is replaced with BSL_r in case X is affine. The proof of Theorem 6.15 which leads to Corollary 6.16 requires a few lemmas needed to check the descent criterion in Theorem A.2.

Lemma 6.11. *Let R be a commutative ring. If $f, g \in R$ with $fR + gR = R$, then the following diagram is homotopy cartesian*

$$\begin{array}{ccc} B_{Zar}GL_n(R) & \longrightarrow & B_{Zar}GL_n(R_f) \\ \downarrow & & \downarrow \\ B_{Zar}GL_n(R_g) & \longrightarrow & B_{Zar}GL_n(R_{fg}). \end{array}$$

Proof. This follows from descent and can also be checked using Quillen's Theorem B [Qui73]. \square

For $n \in \mathbb{N}$, we write $\text{Vect}_n^R(R[T_1, \dots, T_n])$ for the full subcategory of the category $\text{Vect}_n(R[T_1, \dots, T_n])$ of those projective modules P which are extended from R , that is, which are isomorphic to $Q \otimes_R R[T_1, \dots, T_n]$ for some $Q \in \text{Vect}_n(R)$. Write $B_{Zar}^R GL_n(R[T_1, \dots, T_n])$ for the classifying space (that is, nerve) of the category $\text{Vect}_n^R(R[T_1, \dots, T_n])$.

Lemma 6.12. *Let R be a commutative ring. If $f, g \in R$ with $fR + gR = R$, then for all integers $q \geq 0$ the following diagram is homotopy cartesian*

$$\begin{array}{ccc} B_{Zar}^R GL_n(R[T_1, \dots, T_q]) & \longrightarrow & B_{Zar}^{R_f} GL_n(R_f[T_1, \dots, T_q]) \\ \downarrow & & \downarrow \\ B_{Zar}^{R_g} GL_n(R_g[T_1, \dots, T_q]) & \longrightarrow & B_{Zar}^{R_{fg}} GL_n(R_{fg}[T_1, \dots, T_q]). \end{array}$$

Proof. By Quillen's Patching Theorem [Qui76, Theorem 1'], a projective $R[T_1, \dots, T_q]$ -module P is extended from R if and only if P_f and P_g are extended from R_f and R_g . Hence, the lemma follows from Lemma 6.11 with $R[T_1, \dots, T_q]$ in place of R . \square

Write $B_{Zar}^R GL_n(\Delta R)$ for the diagonal of the simplicial space $q \mapsto B_{Zar}^R GL_n(\Delta_q R)$.

Corollary 6.13. *Let R be a commutative ring. If $f, g \in R$ with $fR + gR = R$, then the following diagram of simplicial sets is homotopy cartesian*

$$\begin{array}{ccc} B_{Zar}^R GL_n(\Delta R) & \longrightarrow & B_{Zar}^{R_f} GL_n(\Delta R_f) \\ \downarrow & & \downarrow \\ B_{Zar}^{R_g} GL_n(\Delta R_g) & \longrightarrow & B_{Zar}^{R_{fg}} GL_n(\Delta R_{fg}). \end{array}$$

Proof. For $q \in \mathbb{N}$, the diagram is homotopy cartesian for Δ_q in place of Δ , in view of Lemma 6.12. The Corollary now follows from the Bousfield-Friedlander Theorem [BF78, Theorem B.4] which we can apply since the simplicial set $q \mapsto \pi_0 B_{Zar}^R GL_n(\Delta_q R)$ of connected components is a constant simplicial set for any R . \square

Write $B_{Zar}^\bullet GL_n^\Delta$ the simplicial presheaf

$$X \mapsto B_{Zar}^R GL_n(\Delta R), \quad \text{where } R = \Gamma(X, O_X).$$

Inclusion of its degree zero space into the simplicial space induces a map of simplicial presheaves $B_{Zar} GL_n \rightarrow B_{Zar}^\bullet GL_n^\Delta$.

Theorem 6.14. *Let $X = \text{Spec } R$ where R is a noetherian ring. Then the natural maps of simplicial presheaves $BGL_n \rightarrow B_{Zar} GL_n \rightarrow B_{Zar}^\bullet GL_n^\Delta$ induce a bijection*

$$[X, BGL_n]_{Zar} \cong [X, B_{Zar}^\bullet GL_n^\Delta]_{Zar}.$$

Proof. This follows from Corollary 6.13 in view of Theorem A.2. \square

We can reformulate the theorem as follows. For a simplicial presheaf F defined on the category of schemes, we write $\text{Sing}^{\mathbb{A}^1} F$ for the simplicial presheaf $X \mapsto (q \mapsto F(X \times \text{Spec } \Delta_q \mathbb{Z}))$. The map of simplicial rings $\mathbb{Z} \rightarrow \Delta$ induces a natural map $F \rightarrow \text{Sing}^{\mathbb{A}^1} F$ of simplicial presheaves.

Theorem 6.15. *Let $X = \text{Spec } R$ where R is a noetherian ring. Then the natural map of simplicial presheaves $BGL_n \rightarrow \text{Sing}^{\mathbb{A}^1} BGL_n$ induces a bijection*

$$\Phi_n(X) = [X, BGL_n]_{Zar} \cong [X, \text{Sing}^{\mathbb{A}^1} BGL_n]_{Zar}.$$

Proof. This follows from Theorem 6.14 since the natural map of simplicial presheaves $\text{Sing}^{\mathbb{A}^1} BGL_n \rightarrow B_{Zar}^\bullet GL_n^\Delta$ is a weak equivalence at the local rings of X . \square

By definition of the presheaf of perfect groups \tilde{E}_n , the canonical map of simplicial presheaves $BGL_n \rightarrow \text{Sing}^{\mathbb{A}^1} BGL_n$ factors through BGL_n^+ . From Theorem 6.15 we therefore obtain the following.

Corollary 6.16. *Let $X = \text{Spec } R$ be an affine noetherian scheme. Then the string of maps of simplicial presheaves $BGL_n \rightarrow BGL_n^+ \rightarrow \text{Sing}^{\mathbb{A}^1} BGL_n$ induces the sequence of maps*

$$[X, BGL_n]_{Zar} \rightarrow [X, BGL_n^+]_{Zar} \rightarrow [X, \text{Sing}^{\mathbb{A}^1} BGL_n]_{Zar}$$

whose composition is a bijection.

Definition 6.17. Let X be a scheme with infinite residue fields and V an oriented rank n vector bundle on X . The *Euler class*

$$e(V) \in H_{Zar}^n(X, \mathcal{K}_n^{MW})$$

of V is the image of $[V] \in [X, BSL_n]_{Zar}$ under the canonical map

$$[X, BSL_n]_{Zar} \rightarrow [X, BSL_n^+]_{Zar} \xrightarrow{e} H_{Zar}^n(X, \mathcal{K}_n^{MW}).$$

By construction, we have $e(W \oplus O_X) = 0$ for any rank $n-1$ oriented vector bundle W .

Theorem 6.18. *Let R be a commutative noetherian ring of dimension $n \geq 2$. Assume that all its residue fields are infinite. Let P be an oriented rank n projective R -module. Then*

$$P \cong Q \oplus R \Leftrightarrow e(P) = 0 \in H_{Zar}^n(R, \mathcal{K}_n^{MW}).$$

Proof. We already know that $e(Q \oplus R) = 0$. So assume $e(P) = 0$. In view of Corollary 6.16, the maps of simplicial presheaves

$$BSL_r \rightarrow BSL_r^+ \rightarrow BGL_r^+ \rightarrow \text{Sing}^{\mathbb{A}^1} BGL_r$$

induce a commutative diagram

$$\begin{array}{ccccc} [X, BSL_{n-1}]_{Zar} & \longrightarrow & [X, BSL_{n-1}^+]_{Zar} & \longrightarrow & [X, BGL_{n-1}]_{Zar} \\ \downarrow & & \downarrow & & \downarrow \\ [X, BSL_n]_{Zar} & \longrightarrow & [X, BSL_n^+]_{Zar} & \longrightarrow & [X, BGL_n]_{Zar} \end{array}$$

where the horizontal composition is the map which forgets the orientation. The commutativity of this diagram together with Theorem 6.9 and the hypothesis $e(P) = 0$ implies the result. \square

Remark 6.19. Theorem 6.18 is a generalization of a theorem of Morel [Mor12, Theorem 8.14] who proved it for X smooth affine over an infinite perfect field. To compare the two versions, note that instead of our Milnor-Witt K -theory sheaf, Morel uses the unramified Milnor-Witt K -theory sheaf. But for a smooth X over an infinite field of characteristic not 2, the canonical map from our Milnor-Witt K -sheaf to Morel's Milnor-Witt K -sheaf is an isomorphism which follows from the exactness of the Gersten complex for Milnor-Witt K -theory of regular local rings containing an infinite field of characteristic not 2 [GSZ16]. Moreover, Morel uses Nisnevich cohomology instead of Zariski cohomology. Again because of the exactness of the Gersten complex for K^{MW} , the change of topology map is an isomorphism for X smooth over an infinite field of characteristic not 2:

$$H_{Zar}^*(X, \mathcal{K}^{MW}) \cong H_{Nis}^*(X, \mathcal{K}^{MW}).$$

Remark 6.20. Let L be a line bundle on $X = \text{Spec } R$. Theorem 6.18 has an evident generalization to rank n vector bundles P with orientation $w : \Lambda_R^n P \cong L$ in L . Equip $R^{n-1} \oplus L$ with the canonical orientation $\Lambda_R^n(R^{n-1} \oplus L) \cong \Lambda_R^{n-1} R^{n-1} \otimes \Lambda_R^1 L = L$, and denote by $SL_n^L(R)$ the group of orientation perserving R -linear automorphisms of $R^{n-1} \oplus L$. Then

$$\Phi_n^L(X) = [X, BSL_n^L]_{Zar}$$

is the set of isomorphism classes of rank n vector bundles on X with orientation in L . Define the sheaf $\mathcal{K}_n^{MW}(L)$ on X as

$$\mathcal{K}_n^{MW}(L) = \mathcal{H}_n(SL_n^L, SL_{n-1}^L).$$

Its stalks are, of course, the usual Milnor-Witt K -groups of the local rings of X . Replacing SL_n with SL_n^L everywhere, we obtain an Euler class map as in Definition 6.8 and an Euler class $e(P, L) \in H^n(R, \mathcal{K}_n^{MW}(L))$ for projective modules P with orientation in L as in Definition 6.17.

With the definitions in Remark 6.20 we have the following theorem whose proof is *mutatis mutandis* the same as in the case $L = R$ in Theorem 6.18.

Theorem 6.21. *Let R be a commutative noetherian ring of dimension $n \geq 2$. Assume that all its residue fields are infinite. Let L be a line bundle on R . Let P be a rank n projective R -module with orientation in L . Then*

$$P \cong Q \oplus R \Leftrightarrow e(P, L) = 0 \in H_{Zar}^n(R, \mathcal{K}_n^{MW}(L)).$$

For a field k , denote by $\mathcal{H}(k)$ the Morel-Voevodsky unstable \mathbb{A}^1 -homotopy category of smooth schemes over k [MV99]. Recall that $\Phi_n(X)$ denotes the set of isomorphism classes of rank n vector bundles on the scheme X . The arguments in the proof of Theorem 6.14 can be used to give a simple proof of a theorem of Morel [Mor12, Theorem 8.1 (3)]. Note that we do not need to exclude the case $n = 2$.

Theorem 6.22 (Morel). *Let k be an infinite perfect field. Then for any smooth affine k -scheme X , there is a natural bijection*

$$\Phi_n(X) \cong [X, BGL_n]_{\mathcal{H}(k)}.$$

Proof. Let R be a smooth k -algebra. For each $q \geq 0$, the simplicial presheaf

$$B_{Zar}GL_n\Delta_q = B\text{Vect}_n\Delta_q$$

has the affine B.G.-property for the Zariski and the Nisnevich topology (see [Mor12] for the definition), by descent, or an application of Quillen's theorem B.

By a result of Lindel [Lin82], for any smooth k -algebra R , extension by scalars induces a bijection $\Phi_n(R) \cong \Phi_n(R[T])$. In other words, the simplicial set of connected components $q \mapsto \Phi_n(\Delta_q R) = \pi_0 B_{Zar}GL_n(\Delta_q R)$ of the simplicial space $q \mapsto B_{Zar}GL_n\Delta_q R$ is constant. In view of the Bousfield-Friedlander Theorem [BF78, Theorem B.4.], it follows that the diagonal $\text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$ of the bisimplicial presheaf $q \mapsto B_{Zar}GL_n\Delta_q$ has the affine B.G.-property for the Zariski and the Nisnevich topology. By construction, the simplicial presheaf $\text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$ is \mathbb{A}^1 -invariant. We will show that the map of simplicial presheaves

$$(6.5) \quad \text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n \rightarrow L_{\mathbb{A}^1} \text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$$

is a weak equivalence on affine k -schemes, by an application of [Mor12, Theorem A.19]. We already know that the source of (6.5) is \mathbb{A}^1 -invariant, and satisfies the affine B.G.-property for the Zariski and the Nisnevich topology. Furthermore, $\text{Sing}^{\mathbb{A}^1} GL_n = \Omega_s^1 \text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$ has the affine B.G.-property for the Nisnevich topology because $\text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$ has. The π_0 sheaf of $\text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$ is trivial in the Zariski topology because over a local ring every rank n vector bundle is trivial. Finally, the π_1 sheaf of $\text{Sing}^{\mathbb{A}^1} B_{Zar}GL_n$ in the Zariski topology is the π_0 -sheaf of $\text{Sing}^{\mathbb{A}^1} GL_n$ which is the group of units (for integral schemes), hence

strongly \mathbb{A}^1 -invariant, since for a local ring R and $n \geq 1$, we have $SL_n R = E_n R$. In view of [Mor12, Theorem A.19], the map (6.5) is a weak equivalence on affine k -schemes.

Now, the map $BGL_n \rightarrow \text{Sing}^{\mathbb{A}^1} B_{Zar} GL_n$ is an \mathbb{A}^1 -weak equivalence, and, by Lindel's theorem, we have $\Phi_n(R) = \pi_0 B_{Zar} GL_n(\Delta R)$. This finishes the proof. \square

Remark 6.23. Similar arguments apply to the symplectic groups Sp_n in place of GL_n .

We finish the section with an analog of Theorem 6.18 for the Nisnevich topology. For a rank n oriented vector bundle V on a noetherian scheme X with infinite residue fields, we denote by $e_{Nis}(V) \in H_{Nis}^n(X, \mathcal{K}_n^{MW})$ the image of $e(V)$ under the change of topology map $H_{Zar}^n(X, \mathcal{K}_n^{MW}) \rightarrow H_{Nis}^n(X, \mathcal{K}_n^{MW})$.

Theorem 6.24. *Let R be a commutative noetherian ring of dimension $n \geq 2$. Assume that all its residue fields are infinite. Then there is an exact sequence of pointed sets*

$$\phi_{n-1}^+(R) \xrightarrow{\oplus R} \phi_n^+(R) \xrightarrow{e_{Nis}} H_{Nis}^n(X, \mathcal{K}_n^{MW}).$$

In particular, if P is an oriented rank n projective R -module, then

$$P \cong Q \oplus R \Leftrightarrow e_{Nis}(P) = 0 \in H_{Nis}^n(R, \mathcal{K}_n^{MW}).$$

Proof. The proof is the same as the proof of Theorem 6.18 replacing Theorem A.2 with the Nisnevich descent criterion of [AHW17, Theorem 3.3.4 (ii)]. \square

APPENDIX A. THE AFFINE B.G.-PROPERTY FOR THE ZARISKI TOPOLOGY

Definition A.1. Let X be a scheme and let $F : \text{Open}_X^{op} \rightarrow \text{sSets}$ be a simplicial presheaf on X . We say that F has the *affine B.G.-property for the Zariski topology* if $F(\emptyset)$ is contractible and for any affine $U = \text{Spec } R \in \text{Open}_X$ and $f, g \in R$ with $(f, g) = R$, the following square of simplicial sets is homotopy cartesian

$$(A.1) \quad \begin{array}{ccc} F(R) & \longrightarrow & F(R_f) \\ \downarrow & & \downarrow \\ F(R_g) & \longrightarrow & F(R_{fg}). \end{array}$$

The aim of this appendix is to give a proof of the following result due to Marc Hoyois [Hoy15]. Whereas Hoyois' proof uses ∞ -categories, we give a proof in the framework of model categories based on standard manipulations of homotopy limits.

Theorem A.2 (Hoyois). *Let X be a noetherian scheme and let $F : \text{Open}_X^{op} \rightarrow \text{sSets}$ be a simplicial presheaf on X which has the affine B.G.-property. Then for all affine $Y \in \text{Open}_X$, the following canonical map is a weak equivalence*

$$F(Y) \xrightarrow{\sim} F_{Zar}(Y),$$

where $F \rightarrow F_{Zar}$ is a fibrant replacement of F for the Zariski topology on X .

The proof will occupy the rest of this appendix. We start by reviewing basic properties of homotopy limits [BK72], [CS02].

Let $f : \mathcal{C} \rightarrow \mathcal{D}$ be a functor between small categories. For an object D of \mathcal{D} , the category $(f \downarrow D)$ has objects pairs (C, a) where C is an object of \mathcal{C} and $a : f(C) \rightarrow D$ is a map in \mathcal{D} . A map $(C, a) \rightarrow (C', a')$ in $(f \downarrow D)$ is a map

$C \rightarrow C'$ in \mathcal{C} which makes the induced triangle in \mathcal{D} commute. Composition is composition of maps in \mathcal{C} . There is a similar category $(D \downarrow f)$ whose objects are pairs $(C, a : D \rightarrow fC)$. When $f = id : \mathcal{C} \rightarrow \mathcal{C}$ is the identity functor, one writes $(\mathcal{C} \downarrow \mathcal{C})$ and $(C \downarrow C)$ for $(id \downarrow C)$ and $(C \downarrow id)$. For a small category \mathcal{C} , we denote by BC the classifying space of \mathcal{C} , that is, the nerve simplicial set of \mathcal{C} .

Let $F : \mathcal{C} \rightarrow \mathbf{sSets}$ be a functor from a small category \mathcal{C} to simplicial sets. Assume that F is object-wise fibrant, that is, FC is a fibrant simplicial set for all objects C of \mathcal{C} . Then the *homotopy limit* of F over \mathcal{C} is the simplicial set defined by the equalizer diagram

$$\mathrm{holim}_{\mathcal{C}} F \rightarrow \prod_{C \in \mathcal{C}} \mathrm{Hom}(B(\mathcal{C} \downarrow C), F(C)) \begin{array}{c} \xrightarrow{a} \\ \xrightarrow{b} \end{array} \prod_{\gamma : C \rightarrow C' \in \mathcal{C}} \mathrm{Hom}(B(\mathcal{C} \downarrow C), F(C'))$$

where a and b are induced by

$$\mathrm{Hom}(B(\mathcal{C} \downarrow C), F(C)) \xrightarrow{F\gamma} \mathrm{Hom}(B(\mathcal{C} \downarrow C), F(C'))$$

$$\mathrm{Hom}(B(\mathcal{C} \downarrow C'), F(C')) \xrightarrow{(\mathcal{C} \downarrow \gamma)} \mathrm{Hom}(B(\mathcal{C} \downarrow C), F(C')).$$

If F is not object-wise fibrant, we define the homotopy limit of F over \mathcal{C} as the homotopy limit of $\mathrm{Ex}^{\infty} F$ over \mathcal{C} as above where $F \rightarrow \mathrm{Ex}^{\infty} F$ is Kan's fibrant replacement functor in the category of simplicial sets. So, $F \rightarrow \mathrm{Ex}^{\infty} F$ is an object-wise weak equivalence, that is, $FC \rightarrow \mathrm{Ex}^{\infty} FC$ is a weak equivalence for all $C \in \mathcal{C}$, and $\mathrm{Ex}^{\infty} F$ is object-wise fibrant.

The homotopy limit has the following useful properties.

Functoriality. The homotopy limit $\mathrm{holim}_{\mathcal{C}} F$ is covariantly functorial in F and contravariantly functorial in \mathcal{C} . More precisely, define a category $[\mathrm{Cat}, \mathbf{sSets}]$ whose objects are pairs (\mathcal{C}, F) where \mathcal{C} is a small category and $F : \mathcal{C} \rightarrow \mathbf{sSets}$ is a functor. Given two objects (\mathcal{C}, F) and (\mathcal{D}, G) of $[\mathrm{Cat}, \mathbf{sSets}]$, a morphism $(\mathcal{C}, F) \rightarrow (\mathcal{D}, G)$ in $[\mathrm{Cat}, \mathbf{sSets}]$ is a pair (f, φ) where $f : \mathcal{C} \rightarrow \mathcal{D}$ is a functor and $\varphi : f^*G \rightarrow F$ a natural transformation. Composition is defined as $(g, \gamma) \circ (f, \varphi) = (gf, \varphi \circ f^*(\gamma))$. The homotopy limit defines a functor

$$(A.2) \quad \mathrm{holim} : [\mathrm{Cat}, \mathbf{sSets}]^{op} \rightarrow \mathbf{sSets}$$

which sends the map $(f, \varphi) : (\mathcal{C}, F) \rightarrow (\mathcal{D}, G)$ in $[\mathrm{Cat}, \mathbf{sSets}]$ to the map of simplicial sets

$$(f, \varphi)^* : \mathrm{holim}_{\mathcal{D}} G \rightarrow \mathrm{holim}_{\mathcal{C}} F$$

which is the composition of the two maps [BK72, XI §3.2]

$$\mathrm{holim}_{\mathcal{D}} G \xrightarrow{\mathrm{holim}(f)} \mathrm{holim}_{\mathcal{C}} f^*G \xrightarrow{\mathrm{holim}(\varphi)} \mathrm{holim}_{\mathcal{C}} F.$$

Homotopy Lemma. Let $F \rightarrow F'$ be a natural transformation of functors $F, F' : \mathcal{C} \rightarrow \mathbf{sSets}$ such that for all $C \in \mathcal{C}$ the map $F(C) \rightarrow F'(C)$ is a weak equivalence of simplicial sets, then the induced map on homotopy limits is a weak equivalence [BK72, XI §5.6]:

$$\mathrm{holim}_{\mathcal{C}} F \xrightarrow{\sim} \mathrm{holim}_{\mathcal{C}} F'.$$

Cofinality. A functor $f : \mathcal{C} \rightarrow \mathcal{D}$ between small categories is called *left cofinal* if for every $D \in \mathcal{D}$, the classifying space of the category $(f \downarrow D)$ is contractible.

Let $f : \mathcal{C} \rightarrow \mathcal{D}$ be a left cofinal functor. Then for every functor $F : \mathcal{D} \rightarrow \mathbf{sSets}$, the induced map on homotopy limits is a weak equivalence [BK72, XI §9.2]:

$$(f, 1)^* : \operatorname{holim}_{\mathcal{D}} F \xrightarrow{\sim} \operatorname{holim}_{\mathcal{C}} f^* F.$$

Fubini's theorem. A functor

$$(A.3) \quad \mathcal{C} \rightarrow [\operatorname{Cat}, \mathbf{sSets}]^{op} : C \mapsto (\mathcal{D}_C, F_C)$$

is given by the following data:

- a functor $\mathcal{D} : \mathcal{C}^{op} \rightarrow \operatorname{Cat} : C \mapsto \mathcal{D}_C$,
- for every object $C \in \mathcal{C}$ a functor $F_C : \mathcal{D}_C \rightarrow \mathbf{sSets}$, and
- for every map $\gamma : C_0 \rightarrow C_1$ a natural transformation $\delta_\gamma : \mathcal{D}_\gamma^* F_{C_0} \rightarrow F_{C_1}$ such that $\delta_1 = id$ and $\delta_{\gamma_1 \gamma_0} = \delta_{\gamma_1} \mathcal{D}_{\gamma_1}(\delta_{\gamma_0})$ for any two composable arrows γ_0, γ_1 in \mathcal{C} .

To give such data is equivalent to giving a functor

$$(A.4) \quad F : \mathcal{C} \not\! \! \! \mathcal{D} \rightarrow \mathbf{sSets}$$

where $\mathcal{C} \not\! \! \! \mathcal{D} = (\mathcal{C}^{op} \int \mathcal{D})^{op}$ is the opposite of the Grothendieck construction on the functor $\mathcal{D} : \mathcal{C}^{op} \rightarrow \operatorname{Cat}$. In detail, $\mathcal{C} \not\! \! \! \mathcal{D}$ is the category whose objects are pairs (C, x) where C is an object of \mathcal{C} and x is an object of \mathcal{D}_C . A map $(C_0, x_0) \rightarrow (C_1, x_1)$ in $\mathcal{C} \not\! \! \! \mathcal{D}$ is given by a pair (γ, a) where $\gamma : C_0 \rightarrow C_1$ is a map in \mathcal{C} and $a : x_0 \rightarrow \mathcal{D}_\gamma x_1$ is a map in \mathcal{D}_{C_0} . Composition is defined by $(\gamma_1, a_1) \circ (\gamma_0, a_0) = (\gamma_1 \gamma_0, \mathcal{D}_{\gamma_0}(a_1) \circ a_0)$. The functor (A.4) induced by the collection of data above sends (C, x) to $F_C(x)$ and a map (γ, a) to $\delta_\gamma(x_1) \circ F_{C_0}(a)$.

The composition of the functors (A.2) and (A.3) determine a functor

$$\mathcal{C} \rightarrow \mathbf{sSets} : C \mapsto \operatorname{holim}_{x \in \mathcal{D}_C} F_C(x)$$

which in turn defines a simplicial set

$$\operatorname{holim}_{C \in \mathcal{C}} \operatorname{holim}_{x \in \mathcal{D}_C} F_C(x).$$

On the other hand, the functor (A.4) also determines a simplicial set

$$\operatorname{holim}_{(C,x) \in \mathcal{C} \not\! \! \! \mathcal{D}} F_C(x).$$

The Fubini Theorem for homotopy limits asserts that these two simplicial sets are naturally weakly equivalent [CS02, III Theorem 26.8 and III 31.5]:

$$(A.5) \quad \operatorname{holim}_{C \in \mathcal{C}} \operatorname{holim}_{x \in \mathcal{D}_C} F_C(x) \simeq \operatorname{holim}_{(C,x) \in \mathcal{C} \not\! \! \! \mathcal{D}} F_C(x).$$

If $\mathcal{D} : \mathcal{C}^{op} \rightarrow \operatorname{Cat}$ is a constant functor, that is, $\mathcal{D}_\gamma = id$ for all maps γ in \mathcal{C} , then $\mathcal{C} \not\! \! \! \mathcal{D} = \mathcal{C} \times \mathcal{D}$ and Fubini's Theorem reduces to a weak equivalence [BK72, XI Example 4.3]

$$(A.6) \quad \operatorname{holim}_{C \in \mathcal{C}} \operatorname{holim}_{x \in \mathcal{D}} F_C(x) \simeq \operatorname{holim}_{(C,x) \in \mathcal{C} \times \mathcal{D}} F_C(x).$$

Homotopy pull-backs. A commutative square of simplicial sets

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ Z & \longrightarrow & W \end{array}$$

is homotopy cartesian if and only if the natural map, induced by the unique map from the index category to the final object in Cat ,

$$X \longrightarrow \text{holim}(Y \rightarrow W \leftarrow Z),$$

is a weak equivalence of simplicial sets [BK72, XI Example 4.1 (iv)].

Extended Functoriality. Let $(f_0, \varphi_0), (f_1, \varphi_1) : (\mathcal{C}, F) \rightarrow (\mathcal{D}, G)$ be morphisms in $[\text{Cat}, \text{sSets}]$. A natural transformation $\delta : (f_0, \varphi_0) \rightarrow (f_1, \varphi_1)$ in $[\text{Cat}, \text{sSets}]$ is a natural transformation of functors $\delta : f_0 \rightarrow f_1$ such that $\varphi_0 = \varphi_1 \circ G(\delta) : f_0^* G \rightarrow F$.

If there is a natural transformation $\delta : (f_0, \varphi_0) \rightarrow (f_1, \varphi_1)$ of maps in $[\text{Cat}, \text{sSets}]$ as above then the induced maps on homotopy limits

$$(f_0, \varphi_0)^*, (f_1, \varphi_1)^* : \text{holim}_{\mathcal{D}} G \rightarrow \text{holim}_{\mathcal{C}} F$$

are homotopic.

Proof. The Extended Functoriality is a consequence of Cofinality as follows. Denote by $p : \mathcal{C} \times [1] \rightarrow \mathcal{C}$ the projection. The two maps $(f_i, \varphi_i) : (\mathcal{C}, F) \rightarrow (\mathcal{D}, G)$ are the two compositions in a diagram in $[\text{Cat}, \text{sSets}]$

$$(\mathcal{C}, F) \begin{array}{c} \xrightarrow{(s_0, 1)} \\ \xrightarrow{(s_1, 1)} \end{array} (\mathcal{C} \times [1], p^* F) \xrightarrow{(f, \varphi)} (\mathcal{D}, G)$$

where $[1]$ is the poset $0 < 1$ and $s_i : \mathcal{C} \rightarrow \mathcal{C} \times [1] : C \mapsto (C, i)$ is the obvious inclusion, $i = 0, 1$. The functor $p : \mathcal{C} \times [1] \rightarrow \mathcal{C}$ is left cofinal since for every $C \in \mathcal{C}$ the composition

$$(\mathcal{C} \downarrow C) \xrightarrow{s_0} (p \downarrow C) \xrightarrow{p} (\mathcal{C} \downarrow C)$$

is the identity whereas the the composition

$$(p \downarrow C) \xrightarrow{p} (\mathcal{C} \downarrow C) \xrightarrow{s_0} (p \downarrow C)$$

admits a natural transformation to the identity. Since $(\mathcal{C} \downarrow C)$ has a final object, this category and hence $(p \downarrow C)$ are contractible. By Cofinality, the map

$$(p, 1)^* : \text{holim}_{\mathcal{C}} F \rightarrow \text{holim}_{\mathcal{C} \times [1]} p^* F$$

is a weak equivalence. Since $(s_i, 1)^*(p, 1)^* = 1$, the two maps $(s_i, 1)^*$ are homotopic, $i = 0, 1$. In particular, $(f_0, \varphi_0)^* = (s_0, 1)^*(f, \varphi)^*$ is homotopic to $(f_1, \varphi_1)^* = (s_1, 1)^*(f, \varphi)^*$. \square

Most functors we want to take a homotopy limit of factor through the category Open_X^{op} of open subsets of a space X . Since the category Open_X^{op} is a poset, this simplifies the treatment, and we introduce the following category Cat_X . Its objects are pairs (\mathcal{C}, U) where \mathcal{C} is a small category and $U : \mathcal{C} \rightarrow \text{Open}_X^{op}$ is a functor. A map $f : (\mathcal{C}, U) \rightarrow (\mathcal{D}, V)$ in Cat_X can be thought of as a ‘‘refinement’’. It is a functor $f : \mathcal{C} \rightarrow \mathcal{D}$ such that $U(C) \subset V(f(C))$ for all $C \in \mathcal{C}$. Composition in Cat_X is composition of functors. A natural transformation $\delta : f \rightarrow g$ of maps $f, g : (\mathcal{C}, U) \rightarrow (\mathcal{D}, V)$ in Cat_X is by definition a natural transformation $\delta : f \rightarrow g$ of functors $f, g : \mathcal{C} \rightarrow \mathcal{D}$.

If $F : \text{Open}_X^{op} \rightarrow \text{sSets}$ is a simplicial presheaf on X , then $\text{holim} \circ F$ defines a functor $\text{Cat}_X^{op} \rightarrow \text{sSets}$ by $(\text{holim} \circ F)(\mathcal{C}, U) = \text{holim}_{\mathcal{C}} F U$. A map $f : (\mathcal{C}, U) \rightarrow (\mathcal{D}, V)$ in Cat_X induces a map $(f, \text{can}) : (\mathcal{C}, F U) \rightarrow (\mathcal{D}, F V)$ in $[\text{Cat}, \text{sSets}]$ where $\text{can} : f^* F V \rightarrow F U$ is the restriction map. Thus, F defines a functor

$$F : \text{Cat}_X \rightarrow [\text{Cat}, \text{sSets}] : (\mathcal{C}, U) \mapsto (\mathcal{C}, F U)$$

which sends natural transformations to natural transformations, and $\text{holim} \circ F$ is the composition

$$(A.7) \quad \text{holim} \circ F : \text{Cat}_X^{op} \xrightarrow{F} [\text{Cat}, \text{sSets}]^{op} \xrightarrow{\text{holim}} \text{sSets}.$$

Call two maps $f, g : (\mathcal{C}, U) \rightarrow (\mathcal{D}, V)$ in Cat_X homotopic if there is a zigzag $f = f_0 \rightarrow f_1 \leftarrow f_2 \rightarrow \cdots \leftarrow f_n = g$ of natural transformations of maps $(\mathcal{C}, U) \rightarrow (\mathcal{C}, V)$ in Cat_X . By the Extended functoriality for homotopy limits, the functor (A.7) sends homotopic maps to homotopic maps.

The category Cat_X has a final object, namely $(*, X)$ where $*$ denotes the one-object-one-morphism category, that is, the final object in Cat . In particular, for any (\mathcal{C}, U) in Cat_X , there is a natural map of simplicial sets

$$F(X) \rightarrow \text{holim}_{\mathcal{C}} F(U).$$

If I is a set, we denote by $\mathcal{P}_0(I)$ the category of non-empty subsets $S \subset I$ where we have a unique arrow $S \rightarrow S'$ if $S \subset S'$, otherwise there is no arrow. An open cover $\mathcal{U} = \{U_i \rightarrow U\}_{i \in I}$ of some open $U \subset X$ defines a functor $U : \mathcal{P}_0(I) \rightarrow \text{Open}_X^{op} : S \mapsto U_S$ where $U_S = \bigcap_{s \in S} U_s$.

Definition A.3. Let X be a noetherian scheme, and $\mathcal{U} = \{U_i \rightarrow U \mid i \in I\}$ an open cover of some open subset $U \subset X$. We say that a simplicial presheaf $F : \text{Open}_X^{op} \rightarrow \text{sSets}$ has *descent for \mathcal{U}* if the following canonical map is a weak equivalence

$$F(U) \xrightarrow{\sim} \text{holim}_{\emptyset \neq S \subset I} F(U_S).$$

The following is a version of [Voe10, Lemma 5.6].

Lemma A.4 (Refinement Lemma). *Let $\mathcal{U} = \{U_i \rightarrow X\}_{i \in I}$ and $\mathcal{V} = \{V_j \rightarrow X\}_{j \in J}$ be open covers of X , and assume that \mathcal{V} is a refinement of \mathcal{U} , that is, there is a map $f : J \rightarrow I$ such that $V_j \subset U_{f(j)}$ for all $j \in J$. Let F be a simplicial presheaf on X . If F has descent for \mathcal{V} and for $\mathcal{V} \cap U_S = \{V_j \cap U_S \rightarrow U_S\}_{j \in J}$ for all $S \subset I$, then F has descent for \mathcal{U} .*

Proof. Consider the diagram in Cat_X

$$\begin{array}{ccc} (*, X) & \longleftarrow & (\mathcal{P}_0(I), U) \\ \uparrow & \nearrow f & \uparrow p_I \\ (\mathcal{P}_0(J), V) & \longleftarrow_{p_J} & (\mathcal{P}_0(I) \times \mathcal{P}_0(J), U \cap V). \end{array}$$

The square and the upper left triangle commute in Cat_X . We check that the lower right triangle commutes up to homotopy. To this end, consider the map in Cat_X

$$g : (\mathcal{P}_0(I) \times \mathcal{P}_0(J), U \cap V) \rightarrow (\mathcal{P}_0(I), U) : (S, T) \mapsto S \cup f(T)$$

which is well-defined as $U_S \cap V_T \subset U_{S \cup f(T)}$ in view of the equality $U_{S \cup f(T)} = U_S \cap U_{f(T)}$ and the inclusion $V_T \subset U_{f(T)}$. Now, the (unique) natural transformations

$p_I \rightarrow g$ and $f \circ p_J \rightarrow g$ show that the lower right triangle commutes up to homotopy in Cat_X . Applying the functor $\text{holim} \circ F$ yields a diagram of simplicial sets

$$(A.8) \quad \begin{array}{ccc} F(X) & \longrightarrow & \text{holim}_{\emptyset \neq S \subset I} F(U_S) \\ \downarrow & \searrow & \downarrow \\ \text{holim}_{\emptyset \neq T \subset J} F(V_T) & \longrightarrow & \text{holim}_{\substack{\emptyset \neq T \subset J \\ \emptyset \neq S \subset I}} F(U_S \cap V_T) \end{array}$$

in which the outer square and the upper triangle commute and the lower triangle commutes up to homotopy. The left vertical map is a weak equivalence, by assumption. By the Fubini Theorem for homotopy limits (A.6), the right vertical map can be identified with the map

$$\text{holim}_{\emptyset \neq S \subset I} F(U_S) \rightarrow \text{holim}_{\emptyset \neq S \subset I} \text{holim}_{\emptyset \neq T \subset J} F(U_S \cap V_T)$$

induced by the maps

$$F(U_S) \rightarrow \text{holim}_{\emptyset \neq T \subset J} F(U_S \cap V_T)$$

which are weak equivalence, by assumption. Hence both vertical maps in diagram (A.8) are weak equivalences. It follows that the diagonal map is a weak equivalence, and hence, so are the horizontal maps. \square

Corollary A.5. *Let F be a simplicial presheaf on X and $\{U_i \rightarrow U\}_{i \in I}$ an open cover of some open $U \subset X$. If for some $i \in I$, the map $U_i \rightarrow U$ is the identity, then F has descent for the cover $\{U_i \rightarrow U\}_{i \in I}$.*

Proof. By hypothesis, the cover $\{1 : U \rightarrow U\}$ refines $\{U_i \rightarrow U\}$. Since F has descent for any cover of the form $\{1 : V \rightarrow V\}$, the Refinement Lemma A.4 implies the result. \square

Corollary A.6. *Let F be a simplicial presheaf on X . Let \mathcal{U} and \mathcal{V} be open covers of some open $U \subset X$. If \mathcal{V} is obtained from \mathcal{U} by repeating some open sets, then F has descent for \mathcal{U} if and only if it has descent for \mathcal{V} .*

Proof. By assumption, \mathcal{U} refines \mathcal{V} and \mathcal{V} refines \mathcal{U} . The result follows from the Refinement Lemma A.4 whose hypothesis we check using Corollary A.5. \square

Lemma A.7 (Covering Lemma). *Let F be a simplicial presheaf on X and $V \subset X$ some open subset. Let $\{V_i \rightarrow V\}_{i \in I}$ and $\{U_{i,j} \rightarrow V_i\}_{j \in J}$ be open covers for $i \in I$. For a non-empty $S \subset I$ and function $\sigma : S \rightarrow J$ write $U_\sigma = \bigcap_{i \in S} U_{i, \sigma(i)}$. Assume that F has descent for $\{V_i \rightarrow V\}_{i \in I}$ and for $\{U_\sigma \rightarrow V_S\}_{\sigma : S \rightarrow J}$, $\emptyset \neq S \subset I$. Then F has descent for $\{U_{i,j} \rightarrow V\}_{(i,j) \in I \times J}$.*

Proof. For two sets S, J write J^S for the set of functions $S \rightarrow J$. As before, for a non-empty $T \subset J^S$, write U_T for $\bigcap_{\sigma \in T} U_\sigma$. By assumption, we have weak equivalences of simplicial sets

$$F(V) \xrightarrow{\sim} \text{holim}_{S \in \mathcal{P}_0(I)} F(V_S) \xrightarrow{\sim} \text{holim}_{S \in \mathcal{P}_0(I)} \text{holim}_{T \in \mathcal{P}_0(J^S)} F(U_T).$$

By the Fubini Theorem for homotopy limits (A.5), the right hand term is $\text{holim}_{\mathcal{C}} f^* F U$ for the functor (map of posets)

$$f : \mathcal{C} = \mathcal{P}_0(I) \# \mathcal{P}_0(J^?) = \{(S, T) \mid S \in \mathcal{P}_0(I), T \in \mathcal{P}_0(J^S)\} \longrightarrow \mathcal{P}_0(I \times J)$$

defined by

$$f(S, T) = \{(i, j) \in I \times J \mid i \in S, j \in \{\sigma(i) \mid \sigma \in T\}\}$$

where $FU = F \circ U$ is the usual functor with

$$U : \mathcal{P}_0(I \times J) \rightarrow \text{Open}_X^{\text{op}} : R \mapsto U_R = \bigcap_{(i,j) \in R} U_{i,j}.$$

By Cofinality, we are done once we show that the functor f is left cofinal. Thus, for $R \in \mathcal{P}_0(J^S)$, we have to check that $(f \downarrow R)$ is contractible. But the category $(f \downarrow R)$, considered as a full subcategory of \mathcal{C} , has a final object, namely (S_R, T_R) where

$$\begin{aligned} S_R &= \{i \in I \mid \exists j \in J \mid (i, j) \in R\}, \\ R_i &= \{j \in J \mid (i, j) \in R\}, \\ T_R &= \{\sigma : S_R \rightarrow J \mid \sigma(i) \in R_i\}. \end{aligned}$$

Therefore, $(f \downarrow R)$ is contractible, and we are done. \square

Corollary A.8. *Let $Y \subset X$ be an open subset of a space X . If a simplicial presheaf F on X has descent for the open covers $\{V_j \rightarrow V\}_{j \in J}$, $\{V \rightarrow Y, W \rightarrow Y\}$ and $\{V_j \cap W \rightarrow V \cap W\}_{j \in J}$, then F has descent for $\{V_j \rightarrow Y, W \rightarrow Y\}_{j \in J}$.*

Proof. In view of the hypothesis and Corollary A.6 we can apply the Covering Lemma A.7 to $I = \{0, 1\}$, $V_0 = V$, $V_1 = W$, $U_{0,j} = V_j$, $U_{1,j} = W$. Therefore, F has descent for $\{U_{i,j} \rightarrow Y\}_{i \in I, j \in J}$ which, after omitting repetitions, is $\{V_j \rightarrow Y, W \rightarrow Y\}_{j \in J}$. By Corollary A.6, we are done. \square

Let X be a noetherian scheme and $U \subset X$ an open subscheme. We will call a finite cover $\{U_i \rightarrow U\}_{i \in I}$ of U *elementary* if there exists a total order on I such that for all $i \in I$, there are $f, g \in \Gamma(U_{\leq i}, \mathcal{O}_X)$ such that (f, g) generates the unit ideal in $\Gamma(U_{\leq i}, \mathcal{O}_X)$ and such that $U_i = (U_{\leq i})_f$ and $U_{< i} = (U_{\leq i})_g$ where $U_{\leq i} = \bigcup_{j \leq i} U_j$ and $U_{< i} = \bigcup_{j < i} U_j$. Note that elementary covers are closed under taking base change. Note also that if U is affine then U_i , $U_{< i}$ and $U_{\leq i}$ are also affine.

Lemma A.9. *Let R be a noetherian ring. Then any open cover of $\text{Spec } R$ can be refined by a finite elementary open cover.*

Proof. Since R is noetherian, any cover of $X = \text{Spec } R$ can be refined by a finite cover. So, it suffices to prove the claim for finite covers $\{U_i \rightarrow X\}_{i=1, \dots, n}$. We will prove by induction on $n \in \mathbb{N}_{\geq 1}$ that any cover consisting of n open subsets can be refined by an elementary open cover. If $n = 1$ then $U_1 = X$ and the cover is already elementary as we can choose $f = 1$ and $g = 0$. Assume now that $n \geq 2$. Let I and J be the vanishing ideals of $X - U_n$ and $X - U_{< n}$. Since U_n and $U_{< n}$ cover X , we have $I + J = R$, and we can choose $f \in I$, $g \in J$ with $f + g = 1$. Then $X_f \subset U_n$ and $X_g \subset U_{< n}$, and X_f and X_g cover X . By induction hypothesis, the cover $\{(U_i)_g \rightarrow (U_{< n})_g = X_g\}_{i=1, \dots, n-1}$ can be refined by an elementary cover $\{V_i \rightarrow X_g\}_{i \in I}$. Then $\{V_i \rightarrow X, X_f \rightarrow X\}_{i \in I}$ is an elementary cover of X which refines $\{U_i \rightarrow X\}_{i=1, \dots, n}$. \square

Lemma A.10. *Let X be a noetherian scheme and F a simplicial presheaf on X which has the affine B.G.-property. Then for every open affine $U \subset X$, the simplicial presheaf F has descent for all elementary open covers of U .*

Proof. We will prove the claim by induction on the cardinality of an elementary open cover of an affine open subset of X . If $n \leq 2$, the claim follows from the definition of the affine B.G.-property. Now assume $n \geq 3$. Let $\{U_i \rightarrow U\}_{i=1, \dots, n}$ be an elementary cover of an open affine $U \subset X$. By induction hypothesis and the definition of elementary cover, the simplicial presheaf F has descent for the covers $\{U_i \rightarrow U_{<n}\}_{i=1, \dots, n-1}$, $\{U_i \cap U_n \rightarrow U_{<n} \cap U_n\}_{i=1, \dots, n-1}$ and $\{U_n \rightarrow U, U_{<n} \rightarrow U\}$. By Corollary A.8, the simplicial presheaf F has descent for $\{U_i \rightarrow U\}_{i=1, \dots, n}$. \square

Lemma A.11. *Let X be a noetherian scheme and F a simplicial presheaf on X which has the affine B.G.-property. Then for every open affine $U \subset X$, the simplicial presheaf F has descent for all open affine covers of U .*

Proof. Note that an elementary open cover of an affine scheme is an affine cover. Now the claim follows from Lemma A.10 and the Refinement Lemma A.4 which we can apply since elementary covers are closed under base change, and every intersection of affine open subsets in U is affine. \square

Denote by Sch a small full subcategory of the category of schemes closed under taking open subschemes and fibre products. For instance, Sch could be the category of open subsets of a given scheme, the category of finite type S -scheme, or smooth S -scheme for a noetherian scheme S . Let $\text{Aff} \subset \text{Sch}$ be the full subcategory of affine schemes. For a simplicial presheaf F on Sch , its homotopy right Kan extension from Aff to Sch is the simplicial presheaf \hat{F} on Sch defined by

$$(A.9) \quad \hat{F}(X) = \text{holim}_{U \in (\text{Aff} \downarrow X)} F(U).$$

The canonical map $((\text{Aff} \downarrow X), F) \rightarrow (*, F(X))$ in $[\text{Cat}, \text{sSets}]$ induces a map of simplicial presheaves

$$F \rightarrow \hat{F}.$$

When $U \in \text{Sch}$ is affine then this map induces a weak equivalence of simplicial sets $F(U) \xrightarrow{\sim} \hat{F}(U)$ since then $(\text{Aff} \downarrow U)$ has a final object.

Lemma A.12. *Let $X \in \text{Sch}$ be a scheme, let L_i be line bundles on X , and let $f_i \in \Gamma(X, L_i)$ be global sections of L_i , $i = 1, \dots, n$. Assume that $X = \bigcup_{i \in I} X_{f_i}$. If F is a simplicial presheaf on Sch which has the affine B.G.-property, then its homotopy right Kan extension \hat{F} has descent for the open cover $\{X_{f_i} \rightarrow X\}_{i \in I}$.*

Proof. Let $y : Y \rightarrow X$ be an affine map of schemes. Consider the functor

$$f : (\text{Aff} \downarrow X) \rightarrow (\text{Aff} \downarrow Y) : (V \rightarrow X) \mapsto y^*V = (V \times_X Y \rightarrow Y).$$

The induced functor on opposite categories f^{op} is left cofinal because for every $w : W \rightarrow Y$ in $(\text{Aff} \downarrow Y)$, the category $(w \downarrow f^{op})^{op} = (w \downarrow f)$ has an initial object given by $yw : W \rightarrow X$ and $(1, yw) : W \rightarrow W \times_X Y$. For $\emptyset \neq S \subset I$ and $Y = U_S = \bigcap_{i \in S} X_{f_i} \rightarrow X$ the open inclusion, Cofinality for homotopy limits then yields a weak equivalence of simplicial sets

$$\text{holim}_{W \in (\text{Aff} \downarrow U_S)} F(W) \xrightarrow{\sim} \text{holim}_{V \in (\text{Aff} \downarrow X)} F(V \times_X U_S).$$

Taking homotopy limit over $\mathcal{P}_0(I)$, we obtain from the Homotopy Lemma the weak equivalence of simplicial sets

$$\text{holim}_{S \in \mathcal{P}_0(I)} \text{holim}_{W \in (\text{Aff} \downarrow U_S)} F(W) \xrightarrow{\sim} \text{holim}_{S \in \mathcal{P}_0(I)} \text{holim}_{V \in (\text{Aff} \downarrow X)} F(V \times_X U_S).$$

The left hand side is

$$\text{holim}_{S \in \mathcal{P}_0(I)} \hat{F}(U_S)$$

and the right hand side is

$$\operatorname{holim}_{V \in (\operatorname{Aff} \downarrow X)} \operatorname{holim}_{S \in \mathcal{P}_0(I)} F(V \times_X U_S) = \operatorname{holim}_{V \in (\operatorname{Aff} \downarrow X)} F(V) = \hat{F}(X)$$

since F has descent for open covers of V , by Lemma A.11. Thus, we have a sequence of maps in which the second map is a weak equivalence of simplicial sets

$$\hat{F}(X) \longrightarrow \operatorname{holim}_{S \in \mathcal{P}_0(I)} \hat{F}(U_S) \xrightarrow{\sim} \hat{F}(X).$$

We are done once we show that the composition is homotopic to the identity. The existence of the homotopy follows from the Extended Functoriality for homotopy limits since the following diagram in $[\operatorname{Cat}, \operatorname{sSets}]$ commutes up to natural transformation

$$\begin{array}{ccc} (\mathcal{P}_0(I) \times (\operatorname{Aff} \downarrow X), f^*F) & \xrightarrow{f} & (\mathcal{P}_0(I) \oint (\operatorname{Aff} \downarrow U_?), F) \\ & \searrow & \downarrow \\ & & ((\operatorname{Aff} \downarrow X), F) \end{array}$$

where the horizontal functor is $(S, V \rightarrow X) \mapsto (S, V \times_X U_S \rightarrow U_S)$, the diagonal functor is $(S, V \rightarrow X) \mapsto (V \rightarrow X)$ and the vertical functor is $(S, W \rightarrow U_S) \mapsto (W \rightarrow X)$ using the inclusion $U_S \subset X$. The functor F sends $(S, W \rightarrow U_S)$ and $(W \rightarrow X)$ to $F(W)$. The natural transformation at $(S, V \rightarrow X)$ is the projection map $V \times_X U_S \rightarrow V$. \square

Recall that a quasi-compact scheme X admits an ample family of line bundles if the open subsets X_f form a basis for the Zariski topology on X where $f \in \Gamma(X, L)$ and L runs through all line bundles on X . For instance, any quasi-affine scheme has an ample family of line bundles.

Theorem A.13. *Let Sch be a small category of noetherian schemes closed under open immersions and fibre products. Let F be a simplicial presheaf on Sch which has the affine B.G.-property. Let \hat{F} be the homotopy right Kan extension of F from the full subcategory Aff of affine schemes to Sch ; see (A.9). Let $X \in \operatorname{Sch}$ be a noetherian scheme with an ample family of line bundles. Then \hat{F} has descent for all open covers of X .*

Proof. Since X has an ample family of line bundles, every open cover \mathcal{U} of X can be refined by a cover as in A.12. Since those covers are closed under base change, Lemma A.12 together with the Refinement Lemma A.4 implies that \hat{F} has descent for \mathcal{U} . \square

Proof of Theorem A.2. For an open inclusion $j : Y \subset X$, the natural map of simplicial presheaves $(j^*F)_{\operatorname{Zar}} \rightarrow j^*(F_{\operatorname{Zar}})$ is a map of fibrant objects and a weak equivalence for the Zariski topology, hence it is an object-wise weak equivalence. Therefore, we can replace F with j^*F and assume that $X = Y$ is affine. Then X and all its open subsets have an ample family of line bundles. Let $U, V \subset X$ be open subsets. By Theorem A.13 with $\operatorname{Sch} = \operatorname{Open}_X$ and $U \cup V$ in place of X , the simplicial presheaf \hat{F} has descent for the cover $\{U \rightarrow U \cup V, V \rightarrow U \cup V\}$ of $U \cup V$. That is, \hat{F} sends the square (6.1) to a homotopy cartesian square of simplicial sets. By [BG73, Theorem 4], the map $\hat{F} \rightarrow \hat{F}_{\operatorname{Zar}}$ from \hat{F} to its Zariski fibrant replacement is an object-wise weak equivalence. Since $F \rightarrow \hat{F}$ is an equivalence on affine schemes, the composition $F \rightarrow \hat{F}_{\operatorname{Zar}}$ is an equivalence on affine schemes and a Zariski weak equivalences to a fibrant simplicial presheaf. \square

REFERENCES

- [AHW17] Aravind Asok, Marc Hoyois, and Matthias Wendt. Affine representability results in \mathbb{A}^1 -homotopy theory, I: Vector bundles. *Duke Math. J.*, 166(10):1923–1953, 2017.
- [Bas64] H. Bass. K -theory and stable algebra. *Inst. Hautes Études Sci. Publ. Math.*, (22):5–60, 1964.
- [Bas73] Hyman Bass. Some problems in “classical” algebraic K -theory. In *Algebraic K-theory, II: “Classical” algebraic K-theory and connections with arithmetic (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 3–73. Lecture Notes in Math., Vol. 342. Springer, Berlin, 1973.
- [BF78] A. K. Bousfield and E. M. Friedlander. Homotopy theory of Γ -spaces, spectra, and bisimplicial sets. In *Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II*, volume 658 of *Lecture Notes in Math.*, pages 80–130. Springer, Berlin, 1978.
- [BG73] Kenneth S. Brown and Stephen M. Gersten. Algebraic K -theory as generalized sheaf cohomology. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 266–292. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [BK72] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972.
- [BM99] Jean Barge and Fabien Morel. Cohomologie des groupes linéaires, K -théorie de Milnor et groupes de Witt. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(3):191–196, 1999.
- [BM00] Jean Barge and Fabien Morel. Groupe de Chow des cycles orientés et classe d’Euler des fibrés vectoriels. *C. R. Acad. Sci. Paris Sér. I Math.*, 330(4):287–290, 2000.
- [Bro82] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [CS02] Wojciech Chachólski and Jérôme Scherer. Homotopy theory of diagrams. *Mem. Amer. Math. Soc.*, 155(736):x+90, 2002.
- [Fas08] Jean Fasel. Groupes de Chow-Witt. *Mém. Soc. Math. Fr. (N.S.)*, (113):viii+197, 2008.
- [GSZ16] Stefan Gille, Stephen Scully, and Changlong Zhong. Milnor-Witt K -groups of local rings. *Adv. Math.*, 286:729–753, 2016.
- [Hoy15] Marc Hoyois. *Personal communication*, 2015.
- [HT10] Kevin Hutchinson and Liqun Tao. Homology stability for the special linear group of a field and Milnor-Witt K -theory. *Doc. Math.*, (Extra volume: Andrei A. Suslin sixtieth birthday):267–315, 2010.
- [HT13] Kevin Hutchinson and Liqun Tao. Milnor-Witt K -theory and tensor powers of the augmentation ideal of $\mathbb{Z}[F^\times]$. *Algebra Colloq.*, 20(3):515–522, 2013.
- [Hut90] Kevin Hutchinson. A new approach to Matsumoto’s theorem. *K-Theory*, 4(2):181–200, 1990.
- [Ker09] Moritz Kerz. The Gersten conjecture for Milnor K -theory. *Invent. Math.*, 175(1):1–33, 2009.
- [Lin82] Hartmut Lindel. On the Bass-Quillen conjecture concerning projective modules over polynomial rings. *Invent. Math.*, 65(2):319–323, 1981/82.
- [Mat69] Hideya Matsumoto. Sur les sous-groupes arithmétiques des groupes semi-simples déployés. *Ann. Sci. École Norm. Sup. (4)*, 2:1–62, 1969.
- [May67] J. Peter May. *Simplicial objects in algebraic topology*. Van Nostrand Mathematical Studies, No. 11. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967.
- [Maz05] A. Mazzoleni. A new proof of a theorem of Suslin. *K-Theory*, 35(3-4):199–211 (2006), 2005.
- [Mil70] John Milnor. Algebraic K -theory and quadratic forms. *Invent. Math.*, 9:318–344, 1969/1970.
- [Moo68] Calvin C. Moore. Group extensions of p -adic and adelic linear groups. *Inst. Hautes Études Sci. Publ. Math.*, (35):157–222, 1968.
- [Mor04] Fabien Morel. Sur les puissances de l’idéal fondamental de l’anneau de Witt. *Comment. Math. Helv.*, 79(4):689–703, 2004.
- [Mor12] Fabien Morel. \mathbb{A}^1 -algebraic topology over a field, volume 2052 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2012.

- [Mur94] M. Pavaman Murthy. Zero cycles and projective modules. *Ann. of Math. (2)*, 140(2):405–434, 1994.
- [MV99] Fabien Morel and Vladimir Voevodsky. \mathbf{A}^1 -homotopy theory of schemes. *Inst. Hautes Études Sci. Publ. Math.*, (90):45–143 (2001), 1999.
- [NS89] Yu. P. Nesterenko and A. A. Suslin. Homology of the general linear group over a local ring, and Milnor’s K -theory. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(1):121–146, 1989.
- [OVV07] D. Orlov, A. Vishik, and V. Voevodsky. An exact sequence for $K_*^M/2$ with applications to quadratic forms. *Ann. of Math. (2)*, 165(1):1–13, 2007.
- [Qui73] Daniel Quillen. Higher algebraic K -theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [Qui76] Daniel Quillen. Projective modules over polynomial rings. *Invent. Math.*, 36:167–171, 1976.
- [Sah89] Chih-Han Sah. Homology of classical Lie groups made discrete. III. *J. Pure Appl. Algebra*, 56(3):269–312, 1989.
- [SS03] Stefan Schwede and Brooke Shipley. Equivalences of monoidal model categories. *Algebr. Geom. Topol.*, 3:287–334 (electronic), 2003.
- [Sus82] A. A. Suslin. Stability in algebraic K -theory. In *Algebraic K-theory, Part I (Oberwolfach, 1980)*, volume 966 of *Lecture Notes in Math.*, pages 304–333. Springer, Berlin, 1982.
- [Vas69] L. N. Vaseršteĭn. On the stabilization of the general linear group over a ring. *Math. USSR-Sb.*, 8:383–400, 1969.
- [Vas71] L. N. Vaseršteĭn. The stable range of rings and the dimension of topological spaces. *Funkcional. Anal. i Priložen.*, 5(2):17–27, 1971.
- [vdK76] Wilberd van der Kallen. Injective stability for K_2 . In *Algebraic K-theory (Proc. Conf., Northwestern Univ., Evanston, Ill., 1976)*, pages 77–154. Lecture Notes in Math., Vol. 551. Springer, Berlin, 1976.
- [vdK77] Wilberd van der Kallen. The K_2 of rings with many units. *Ann. Sci. École Norm. Sup. (4)*, 10(4):473–515, 1977.
- [vdK80] Wilberd van der Kallen. Homology stability for linear groups. *Invent. Math.*, 60(3):269–295, 1980.
- [Voe10] Vladimir Voevodsky. Homotopy theory of simplicial sheaves in completely decomposable topologies. *J. Pure Appl. Algebra*, 214(8):1384–1398, 2010.

MARCO SCHLICHTING, MATHEMATICS INSTITUTE, ZEEMAN BUILDING, UNIVERSITY OF WARWICK,
 COVENTRY CV4 7AL, UK
E-mail address: m.schlichting@warwick.ac.uk