

Acta Biologica Hungarica 68(2), pp. 137–149 (2017)
DOI: 10.1556/018.68.2017.2.2

ANTI-MUTAGENIC POTENTIAL OF ALGAL EXTRACTS ON CHROMOSOMAL ABERRATIONS IN *ALLIUM CEPA* L.

HODA ANWER MANSOUR,* HALA MAHFOUZ and NESMA MAHER

Botany Department, Faculty of Science, Ain Shams University, Abassia, Cairo, Egypt

(Received: December 6, 2016; accepted: March 16, 2017)

In the present study, sodium azide (SA) toxicity and the anti-mutagenic effects of different algal extracts at 0.1% and 0.2% concentrations were studied on the mitotic index (MI), chromosomal and nuclear aberrations using *Allium cepa* L. root assay. Moreover, phytochemical screening of photosynthetic pigments, antioxidants compounds, total antioxidant, DPPH scavenging activity, polysaccharides, and phenolic contents were done for two red seaweeds (*Laurencia obtusa* (Hudson) Lamouroux and *Polysiphonia morrowii* Harvey) and for one brown seaweed (*Dictyopteris delicatula* Lamouroux). Treatment with 300 µg/ml sodium azide (SA) induced the highest number of aberrations in *A. cepa* root. A highly significant decrease in the MI appeared after treatment with SA, whereas its value increased following different algal extracts treatments. The highest anti-mutagenic inhibition activity of *Dictyopteris delicatula* added at 0.2% concentration was 72.96%, 69.84%, 56.89% and 43.59% with the algal polyphenol, polysaccharide, aqueous and methanol extract treatments, respectively. The different algal extracts minimized the genotoxicity and exhibited anti-mutagenic potential against SA in a dose-dependent manner. Phytochemical studies showed that *Dictyopteris delicatula* contained the highest total phenol, chlorophyll-*a* and carotenoid quantity. Moreover it exhibited the highest total antioxidant and DPPH scavenging activities. Total polysaccharides and the weight percentage of sulphated polysaccharides were relatively higher in *Polysiphonia morrowii* followed by *Laurencia obtusa*. Hydroquinone and bromophenol were detected only in the studied brown and red seaweeds, respectively. *Polysiphonia morrowii* and *Laurencia obtusa* contained the highest quantity of galactose, rhamnose and xylose, while *Dictyopteris delicatula* contained fucose and mannitol as main monosaccharide units. In conclusion, the studied seaweeds may be considered as rich sources of natural antioxidants. Meanwhile the investigated different algal extracts can minimize the genotoxicity in a dose-dependent manner and exhibit anti-mutagenic potential against the mutagenic substance sodium azide.

Keywords: Algal extracts – antioxidant compounds and activity – anti-mutagenic effects – sodium azide – genotoxicity – *Allium cepa* L.

INTRODUCTION

Seaweeds are considered as one of the natural renewable resources. However, they produce a wide variety of chemically active secondary metabolites such as fatty acids, sterols, phenolic compounds, terpenes, enzymes, polysaccharides, alkaloids, flavo-

*Corresponding author; e-mail address: rodynarwan@yahoo.com

noids, ascorbic acids, glutathione and carotenoids. Recent reports revealed that marine algal extracts, that are rich of natural antioxidant compounds with anti-mutagenic, anti-carcinogenic, anti-genotoxic properties, prevent the deleterious consequences of oxidative stress and improve antioxidant status [10, 23, 46]. This may occur through the direct reaction of these natural algal compounds with reactive oxygen species (ROS) by inactivating, removing them and eventually preventing DNA damages or any carcinogenous changes [42]. In this sense, various bioactive compounds obtained from algae showed different physiological effects on humans and other living organisms that may exert a broad range of biological activities against cancer, tuberculosis, inflammations, and other diseases caused by fungi, bacteria or viruses [1, 42, 54].

Higher plants are often used in many experimental (test) studies since they are not expensive, can be correlated with other animal test systems, and are easily available [18, 31]. The cytological analysis with respect to mitotic behavior considered to be one of the most reliable way to estimate the potency of any mutagens such as, for example sodium azide. The main aim of this work using *Allium cepa* root assay was designed to investigate the anti-mutagenic potential of some marine macro-algal extracts against sodium azide (SA). Also, it was intended to provide a complete picture about some phytochemical constituents of the studied algal species.

MATERIALS AND METHODS

Algal sampling and identification

Three macro-algal species were collected from the coastal zone of El-Temsah Lake (Ismailia Governorate, Egypt). The collected algal samples were washed with tap and distilled water to remove the debris and were air dried. Each algal sample was divided into three parts; the first was preserved in freezer for fresh weight analysis, while the second one was shaded air-dried and grounded into fine powder. The third part was preserved in 4% formalin for identification. Algal identification was based on morphological features determined by microscopical examination, using an Olympus DP SOFT microscope equipped with a Canon Powershot G12 digital camera. Algal samples were identified according to Aleem [3]. Two marine macro-algal samples belonged to Rhodophyta (red algae) and were identified as *Laurencia obtusa* (Hudson) Lamouroux and *Polysiphonia morrowii* Harvey, while the third one belonged to Pheophyta (brown algae) and was identified as *Dictyopteris delicatula* Lamouroux.

Preparation of seaweed extracts

Aqueous extract was prepared by mixing 100 g of dried seaweed powder with one liter of distilled water and homogenized with glass homogenizer. After centrifugation

at 1200×g for 20 min, supernatants were collected and filtered with Whatman No. 1 filter paper followed by 0.22 µm pore sized filter (Millipore, filter type GV) to obtain the water soluble extract. The remaining pellets were further mixed with 20 ml of distilled water, homogenized and centrifuged as described above. Finally, all aqueous extracts were stored at -4 °C for further experiments.

Methanol, phenol and polysaccharide extracts of seaweed powder (100 g) were prepared according to Machu et al. [34], Malik and Singh [36] and Naguib [41]. Total phenols, total polysaccharides, weight % of sulfated polysaccharides, monosaccharides and phenolic acids fractionations (using HPLC analysis) were determined according to A.O.A.C. [5], Blakeney and Mutton [8] and Naguib [41].

Analyses of photosynthetic pigments, antioxidant compounds, DPPH scavenging and total antioxidant activities of algae

Photosynthetic pigments particularly chlorophyll-*a*, chlorophyll-*b* and carotenoids were extracted in 80% acetone and determined spectrophotometrically as recommended by Metzner et al. [38]. Antioxidant compounds (as total phenol, ascorbic acid and glutathione) were estimated according to Namvar et al. [42]. Total antioxidant and DPPH scavenging activities were determined according to the method of Naga and Yukimoto [40].

Anti-mutagenic potential of the extracts

Definite concentrations (0.1% and 0.2%) of the previously prepared algal extracts were used for the *A. cepa* L. root (var. Giza 6, kindly supplied by the Agricultural Research Center (ARC), Giza, Egypt) treatment to study their anti-mutagenic effects.

Onion bulbs were grown at room temperature (28±0.5 °C). When the roots reached length of 2 to 3 cm, they were transferred to the test solutions. For anti-mutagenic analysis, 25 ml of the algal extracts (0.1% and 0.2%) were applied on the *A. cepa* root tips for 3 hours which had been previously treated for 3 hours with 300 µg/ml of sodium azid (a mutagenic substance). SA and distilled water served as positive and negative control, respectively.

Microscopic analysis

The treated roots were fixed in Carnoy's solution (methanol/glacial acetic acid) for 24 hours. After fixation, roots were kept in 70% alcohol in a refrigerator. Slide preparation and staining with Leucobasic Fuchsin were done according to Darlington and La Cour [14]. The preparations were examined under light microscope, 90 fields were completely analyzed for each concentration. The percentage of mitotic index as

well as the frequencies of different mitotic abnormalities were determined. Reduction percentage of abnormalities induced by mutagenic substance (R) was calculated using the formula of Barcelos et al. [7] as shown below:

$$\% \text{ of R} = (A - B) / (A - C) \times 100$$

where (A) is the proportion of aberrations induced by a known mutagen, (B) is the proportion of aberrations induced by a test sample (different algal extracts), (C) is the proportion of aberrations induced by a negative control.

Statistical analysis

Data were subjected to *t*-test and to the analysis of variance (ANOVA) with statistical significance at $p < 0.05$ being tested using the Duncan's Test and Pearson correlation.

RESULTS AND DISCUSSION

Although the anti-mutagenic potential of algal extracts has been extensively studied and well documented [34, 45, 46], yet there is no report on the anti-mutagenic effects of algal extracts in plant test systems. Exposure *A. cepa* root tips to 300 µg/ml of SA for 3 h induced genotoxicity reflected by the high frequency of different types of chromosomal and nuclear abnormalities (Table 1, Fig. 1). Mitotic index (MI) is an indicator of cell proliferation and can be used to evaluate the level of cytotoxicity of an agent [16]. Moreover, data in Table 1 and Fig. 1-I) show a high decrease in the mitotic index after treatment with SA (1.46%) as compared to the negative control (7.58%). Administration of algal extracts to the *A. cepa* root tips exerted a protective potential against the mito-inhibitory effect of SA. However, MI was significantly increased (5.27%) by the treatment with (0.2%) polysaccharide extract of *Polysiphonia morrowii*. Sulfated polysaccharides of algae have a wide range of biological activities including free-radical scavengers, anti-mutagenic, anti-carcinogenic, anti-genotoxic and antioxidants that can prevent the oxidative damages in living organisms [23, 28, 48].

The most common types of abnormalities observed were stickiness, bridges C-M, disturbed meta-anaphase, lagging chromosomes and micronuclei. Considerable frequency of micronuclei was observed in the cells of mitotic division at interphase especially after treatment with SA [17, 19, 37]. The formation of micronuclei is regarded as an induction of mutagenicity and cancer risk [18, 27]. The present results indicate that the maximum percentages of nuclear and total chromosomal abnormalities reached 9.09% and 60.00%, respectively, following SA treatment (300 µg/ml) as compared with the negative control (0.00% and 0.86). Moreover, the total percentage of abnormalities in the root tip cells of *A. cepa* was significantly decreased by the different treatments of algal extracts (Table 1, Fig. 1-II). Effect of polyphenol

Table 1
Effects of different algal extracts on mitotic index, total abnormalities and total inhibitions

Treatments	% of conc.	Mitotic index	% of total Ab.	Inhibition of Ab. %	% of different types of abnormalities													
					Metaphase			Ana-telophase			Interphase							
					Sti.	Lag.	Dis.	C-M	Brid.	Dis.	Lag.	Sti.	Micro.					
Negative control (H ₂ O)	–	7.58±0.65	0.86±0.10	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Positive control (SA)	300 µg/ml	1.46±0.32	60.00±0.12	–	27.37	6.32	11.58	5.26	19.23	3.85	3.85	38.46	9.09					
Polysaccharides extracts																		
<i>Laurencia obtusa</i>	0.1	3.99±0.61 ^j	31.10±0.12 ⁱ	49.11	16.94	2.42	6.45	8.06	19.78	13.19	9.89	46.15	1.37					
	0.2	3.76±0.25 ^l	28.13±0.15 ^l	53.92	19.05	2.86	9.52	2.86	12.12	4.04	8.08	32.32	1.37					
<i>Polysiphonia morrowii</i>	0.1	3.78±0.81 ^k	26.39±0.13 ^m	56.99	56.99	21.74	6.09	8.70	9.62	1.92	1.92	5.22	1.24					
	0.2	5.27±0.52 ^a	20.83±0.05 ^t	65.34	65.34	16.38	9.48	7.76	10.81	1.91	4.46	2.59	1.54					
<i>Dictyopteris delicatula</i>	0.1	4.37±0.35 ^h	19.25±0.03 ^u	68.18	68.18	10.17	3.39	6.76	7.62	3.81	3.81	8.47	1.39					
	0.2	5.02±0.68 ^c	17.93±0.02 ^w	69.84	69.84	8.78	4.73	4.05	7.09	3.94	5.51	–	1.63					
Polyphenol extracts																		
<i>Laurencia obtusa</i>	0.1	4.81±0.67 ^f	25.78±0.15 ^p	59.10	59.10	12.93	3.45	6.90	14.89	6.38	4.96	8.62	0.29					
	0.2	5.01±0.39 ^d	26.21±0.09 ^o	58.55	58.55	18.55	8.06	5.64	9.23	3.07	6.15	4.84	0.40					
<i>Polysiphonia morrowii</i>	0.1	4.34±0.24 ^f	24.72±0.17 ^q	61.17	61.17	14.29	3.57	8.04	11.113	1.71	2.56	8.04	0.26					
	0.2	4.54±0.26 ^e	18.16±0.14 ^v	72.21	72.21	10.66	5.74	6.56	6.56	1.64	1.64	9.02	0.23					
<i>Dictyopteris delicatula</i>	0.1	4.91±0.63 ^e	22.97±0.25 ^r	64.07	64.07	14.07	2.96	5.93	12.42	5.23	5.88	2.96	0.22					
	0.2	5.17±0.22 ^b	17.32±0.26 ^x	72.96	72.96	8.82	5.15	3.68	9.52	2.38	3.17	2.21	0.32					

Table 1 (cont.)

Treatments	% of conc.	Mitotic index	% of total Ab.	Inhibition of Ab. %	% of different types of abnormalities									
					Metaphase			Ana-telophase			Interphase			
					Sti.	Lag.	Dis.	C-M	Brid.	Dis.	Lag.	Sti.	Micro.	
Methanol extracts														
<i>Laurencia obtusa</i>	0.1	1.68±0.54 ^x	42.61±0.32 ^a	29.98	29.98	35.19	7.41	3.70	24.39	12.20	7.32	3.70	1.72	
	0.2	1.95±0.14 ^y	35.38±0.24 ^e	34.21	34.21	33.33	4.44	2.22	37.50	5.00	5.00	4.44	1.65	
<i>Polysiphonia morrowii</i>	0.1	2.05±0.64 ^u	38.40±0.15 ^b	36.29	36.29	39.99	13.33	3.33	29.82	5.26	3.51	3.33	1.94	
	0.2	2.87±0.25 ⁿ	36.41±0.85 ^c	39.78	39.78	23.25	9.30	2.33	37.50	12.50	6.25	1.16	1.81	
<i>Dictyopteris delicatula</i>	0.1	2.17±0.65 ^s	35.33±0.23 ^d	41.39	41.39	18.84	4.35	8.70	55.55	5.56	8.33	2.90	1.60	
	0.2	2.99±0.68 ^m	34.29±0.32 ^f	43.59	43.59	26.60	5.32	6.38	32.76	6.90	6.90	2.13	1.42	
Aqueous extracts														
<i>Laurencia obtusa</i>	0.1	2.08±0.87 ^t	34.13±0.36 ^g	44.07	44.07	29.99	5.71	2.86	26.78	8.93	5.36	—	—	
	0.2	2.17±0.65 ^s	32.00±0.023 ^h	47.40	47.40	28.57	4.76	3.17	24.49	16.33	4.08	4.76	—	
<i>Polysiphonia morrowii</i>	0.1	2.83±0.25 ^o	21.62±0.14 ^s	46.05	46.05	15.96	8.51	2.13	50.00	11.91	9.52	1.06	1.81	
	0.2	2.51±0.54 ^p	28.21±0.19 ^k	53.03	53.03	27.59	3.45	3.45	22.58	3.23	3.23	1.72	1.75	
<i>Dictyopteris delicatula</i>	0.1	2.19±0.21 ^r	28.29±0.32 ^j	53.21	53.21	23.08	3.08	4.62	22.00	4.00	6.00	—	1.57	
	0.2	2.41±0.52 ^q	26.37±0.36 ⁿ	56.89	56.89	19.32	3.41	2.27	28.57	7.14	4.76	1.14	—	

% of total abnormalities and total % of inhibition in *Allium cepa* root tips [SA = Sodium azide, Ab. = abnormalities, Sti. = Stickiness, Dis. = Disturbed, Lag. = Lagging, C-M = C metaphase, Brid. = Bridge and Micro. = Micronuclei].
Values significant at $p < 0.05$ by ANOVA.




Fig. 1. (I). Changes in the mitotic index (MI) and (II). Frequency of total abnormalities stages after treating *Allium cepa* root tips (pretreated with SA) with the different concentrations of macro-algal extracts for 3 hours. For (I & II), A = negative control+positive control+algal polysaccharide extracts; B = negative control+positive control+algal polyphenol extracts; C = negative control+positive control+algal methanol extracts and D = negative control+positive control+algal aqueous extracts

(17.32%) and polysaccharide (17.93%) extracts of *Dictyopteris delicatula* obtained at 0.2% alga concentration showed the lowest percentage of total chromosomal aberrations. Additionally, *Dictyopteris delicatula* at the same concentration showed the highest activity in total abnormalities inhibition (72.96%, 69.84%, 56.89% and 43.59%) with the polyphenol, polysaccharide, aqueous and methanol treatments, respectively. The lowest inhibitory effect (29.98%) was induced by the (0.1%) methanol treatment of *Laurencia obtusa*. Phenolic compounds were found to be one of the most effective antioxidants and they were reported to have very strong anti-mutagenic properties in brown and red algae [20, 24, 28].

Table 2
Photosynthetic pigments, antioxidant compounds, DPPH scavenging and total antioxidant activities of algae

Marine macro-algae	Pigments ($\mu\text{g}\cdot\text{g}^{-1}$ fw)			Total phenols ($\text{mg}\cdot\text{g}^{-1}$ dw)	Ascorbic acid ($\mu\text{M}\cdot\text{g}^{-1}$ fw)	Glutathione ($\text{mM}\cdot\text{g}^{-1}$ fw)	DPPH ($\mu\text{g}\cdot\text{g}^{-1}$ dw)	Total antioxidant activity ($\mu\text{g}\cdot\text{g}^{-1}$ dw)
	Chl. a	Chl. b	Carotenoids					
<i>Laurencia obtusa</i>	2.9±0.6 ^b	3.28±0.1 ^c	1.72±0.03 ^b	23.2±5.83 ^a	2.28±0.06 ^a	20.6±0.23 ^b	1.91±0.08 ^a	82.8±7.80 ^c
<i>Polysiphonia morrowii</i>	2.4±0.51 ^a	2.7±0.1 ^b	2.02±0.03 ^a	27.3±8.90 ^b	2.53±0.08 ^b	26.6±0.35 ^c	3.33±0.41 ^b	107.2±2.21 ^b
<i>Dictyopteris delicatula</i>	9.17±0.21 ^c	1.34±0.01 ^a	6.62±0.5 ^c	39.4±11.6 ^c	2.96±0.07 ^c	11.6±0.18 ^a	5.11±0.39 ^c	156±2.65 ^a

Values significant at $p < 0.05$ by ANOVA.

Phytochemical screening data shown in Table 2 reveal that chlorophyll-*a* and carotenoid contents reached their highest values (9.17 and 6.62 $\mu\text{g} \cdot \text{g}^{-1}$ fw, respectively) in *Dictyopteris delicatula*. Chlorophyll-*b* had its maximum concentration (3.28 $\mu\text{g} \cdot \text{g}^{-1}$ fw) in *Laurencia obtusa*. According to Christaki et al. [11] and Kumar et al. [26], brown algae are commonly characterized by the highest carotenoid contents. The studied *Polysiphonia morrowii* and *Dictyopteris delicatula* species showed a relatively high level of glutathione (26.6 $\mu\text{M} \cdot \text{g}^{-1}$ fw), and ascorbic acid (2.96 $\mu\text{M} \cdot \text{g}^{-1}$ fw), respectively, compared to that measured in some higher plants. Our results are in accordance with the findings of Aguilera et al. [2] and Kumar et al. [26]. Moreover, *Dictyopteris delicatula* exhibited the highest total antioxidant and DPPH scavenging capacities (156 and 5.11 $\mu\text{g} \cdot \text{g}^{-1}$ dw respectively), followed by *Polysiphonia morrowii* (107.2 and 3.33 $\mu\text{g} \cdot \text{g}^{-1}$ dw) and *Laurencia obtusa* (82.8 and 1.91 $\mu\text{g} \cdot \text{g}^{-1}$ dw, respectively). *Dictyopteris delicatula* was found to possess immense antioxidant potential in the form of total antioxidant, and reduction power activities [9, 33, 35].

Data in Table 2 show significant differences ($p < 0.05$) in the total phenols contents among the tested *Dictyopteris* (39.4 $\text{mg} \cdot \text{g}^{-1}$ dw), *Laurencia* and *Polysiphonia* (23.2 and 27.2 $\text{mg} \cdot \text{g}^{-1}$ dw, respectively). Phenolic compounds are potential antioxidants and free-radical scavengers. Brown algae contain, in general, higher amounts of phenolic compounds than red and green algae [48]. Higher antioxidant contents in *Dictyopteris* may be attributed to its high ascorbic acid, carotenoids and the total phenolic contents [11, 48]. However, there are a number of studies proving a linear relationship between the total antioxidant capacity and total phenolic contents in algae [34].

Table 3
Area % of the phenolic acids fractions (mg/100 g dw) of algae

Phenolic acids	Red seaweeds		Brown seaweed
	<i>Laurencia obtusa</i>	<i>Polysiphonia morrowii</i>	<i>Dictyopteris delicatula</i>
	(%) phenolic acids fractions		
Cinnamic acid	0.12	0.089	0.93
Isopropyl 3,5-dimethoxy-4-hydroxycinnamate	0.05	N.D.	0.3
Gallic acid	0.07	0.23	1.57
Caffeic acid	1.2	1.5	1.09
Coumaric acid	0.27	0.17	1.82
Ferulic acid	0.9	0.51	0.26
Hydroquinone	N.D.	N.D.	0.97
Salicylic acid (methyl-4-hydroxy benzoate)	N.D.	0.01	0.72
o-Bromophenol	0.24	0.78	N.D.

(N.D. = not detected).

The dominant components of phenolic acids fraction (Table 3) in *Dictyopteris* were coumaric acid (1.82%) followed by gallic (1.57%), caffeic acid (1.09%), hydroquinone (0.97%) cinnamic acid (0.93%) and salicylic acid (0.72%). The studied red algae were dominated by caffeic, ferulic, *o*-bromophenol and coumaric acid. These were in agreement with the findings of Hsu [22] and Onofrejovaa et al. [44]. Hayat et al. [21] reported that cinnamic acid and some of its esters occur in brown alga *Spatoglossum variabile* (family: Dictyoaceae). Moreover, our present data show that hydroquinone can be detected only in the studied brown alga species while bromophenol was characteristic for the investigated red seaweeds. This finding is confirmed by the data of Leandrini de Oliveira et al. [28] and Li et al. [30].

The results shown in Table 4 reveal that the studied red weed species (*Polysiphonia morrowii* and *Laurencia obtusa*) contain the most monosaccharides sugar galactose, rhamnose and xylose [12, 53], whereas *Dictyopteris delicatula* showed to contain highest area (%) from fucose followed by mannitol as main monosaccharide units. These are in agreement with the findings of Dai-Hung and Se-Kwon [13], and Li et al. [29]. Total polysaccharides and the weight % of sulfated polysaccharides dominated in *Polysiphonia morrowii* (71.1 mg/g DW and 31.07%), followed by *Laurencia obtusa* (37.8 mg/g DW and 25.7%) and *Dictyopteris delicatula* (28.8 mg/g DW and 18.1%). However, several investigations have reported that sulfated polysaccharides are commonly occurring at high concentrations in red marine algae [52, 55].

Table 4
Area % of the monosaccharides fractions (mg/100 g dw), weight % of sulfated polysaccharides and total polysaccharides of algae

Name of sugars	Red seaweeds		Brown seaweed
	<i>Laurencia obtusa</i>	<i>Polysiphonia morrowii</i>	<i>Dictyopteris delicatula</i>
(%) of polysaccharides fractions			
Glucose	22.75	24.75	8.64
Fructose	1.36	2.09	7.61
Arabinose	0.44	0.69	0.16
Rhamnose	50.06	58.2	9.03
Xylose	30.3	39.53	23.50
Mannose	0.2	0.06	5.03
Fucose	0.013	0.07	41.02
Galactose	52.07	81.15	0.41
Mannitol	0.07	0.03	39.30
w% of SO ₄	25.7	31.07	18.1
Total polysaccharides (mg·g ⁻¹ dw)	37.8±2.71 ^b	71.1±0.92 ^a	28.8±0.89 ^c

Values significant at p<0.05 by ANOVA.

The estimated biogenic algal compounds such as phenolic acids, polysaccharides, ascorbic acid, glutathione and a significant radical scavenging activities are most likely responsible for the anti-mutagenic effect of the studied algal extracts [15, 29, 47].

CONCLUSIONS

The present study demonstrates the possible pharmaceutical importance of algae with their chemo-preventive potential in the field of cogency. The investigated different algal extracts of *Laurencia obtusa* (Hudson) Lamouroux, *Polysiphonia morrowii* Harvey and *Dictyopteris delicatula* Lamouroux have the potency to suppress or modulate the mutagenic effect induced by SA in a dose-dependent manner. However, further experiments using different test-systems are required to establish adequate procedures for the biological use of this algal extracts.

REFERENCES

1. Abdul, Q. A., Choi, R. J., Jung, H. A., Choi, J. S. (2016) Health benefit of fucosterol from marine algae: a review. *J. Sci. Food Agric.* 96, 1856–1866.
2. Aguilera, J., Dummermuth, A., Karsten, U., Schriek, R., Wiencke, C. (2002) Enzymatic defenses against photo-oxidative stress induced by ultraviolet radiation in Arctic marine macroalgae. *Polar Biol.* 25, 432–441.
3. Aleem, A. A. (1993) The Marine Algae of Alexandria, University of Alexandria, Alexandria, Egypt.
4. Amer, S. M., Ali, E. M. (1986) Cytological effects of pesticides. XVII. Effect of the insecticide dichlorvos on root mitosis of *Vicia faba*. *Cytologia* 51, 21–25.
5. A.O.A.C. (1995) Official Methods of Analysis of the Association of Official Agricultural Chemists. Pub A.O.A.C. Washington D.C., U.S.A.
6. Azuine, M. A., Bhide, S. V. (1992) Chemo-preventive effect of turmeric against stomach and skin tumors induced by chemical carcinogens in Swiss mice. *Nutr. Cancer* 17, 77–83.
7. Barcelos, G. R. M., Shimabukuro, F., Mori, M. P., Maciel, M. A. M., de Syllos Cólus, I. M. (2007) Evaluation of mutagenicity and anti-mutagenicity of cashew stem bark methanolic extract *in vitro*. *J. Ethnopharmacol.* 114, 268–273.
8. Blakeney, A. B., Mutton, L. L. (1980) A simple calorimetric method for the determination of sugars in fruits and vegetables. *J. Sci. Food Agric.* 31, 889–897.
9. Burritt, D. J., Larkindale, J., Hurd, C. L. (2002) Antioxidant metabolism in the intertidal red seaweed *Stictosiphonia arbuscula* following desiccation. *Planta* 215, 829–838.
10. Chojnacka, K., Saeid, A., Witkowska, Z., Tuhy, L. (2012) Biologically active compounds in seaweed extracts – the prospects for the application. *The Open Conf. Proceed. J.* 3, 20–28.
11. Christaki, E., Bonos, E., Giannenas, I., Florou-Paneri, P. (2013) Functional properties of carotenoids originating from algae. *J. Sci. Food and Agri.* 93, 5–11.
12. Costa, L. S., Fidelis, G. P., Cordeiro, S. L., Oliveira, R. M., Sabry, D. A. (2010) Biological activities of sulfated polysaccharides from tropical seaweeds. *Biomed. Pharmacother.* 64, 21–28.
13. Dai-Hung, N., Se-Kwon, K. (2013) Sulfated polysaccharides as bioactive agents from marine algae. *Inter. J. Biol. Macromolec.* 62, 70–75.
14. Darlington, C. D., La Cour, L. F. (1976) *The Handling of Chromosomes*, 6th edn. Wiley, New York.
15. El-Din, N. S., El-Sherif, Z. M. (2012) Nutritional value of some algae from the north-western Mediterranean coast of Egypt. *J. Appl. Phycol.* 24, 613–626.

16. Elghamery, A. A., Elkholy, M. A., El-Yousse, R. A. (2003) Evaluation of cytological effects of Zn²⁺ in relation to germination and root growth of *Nigella sativa* L. and *Triticum aestivum* L. *Mutation Res.* 537, 2941.
17. Fernandes, T. C. C., Mazzero, D. E. C., Marin-Morales, M. A. (2007) Mechanism of micronuclei formation in polyploidized cells of *Allium cepa* exposed to trifluralin herbicide. *Pesticide Biochem. Physiol.* 88, 252–259.
18. Fiskesjö, G. (1988) The *Allium test* – an alternative in environmental studies: the relative toxicity of metal ions. *Mutat. Res.* 197, 243–260.
19. Gadano, A., Gurni, A., López, P., Ferraro, G., Carballo, M. (2002) *In vitro* genotoxic evaluation of the medicinal plant *Chenopodium ambrosioides* L. *J. Ethnopharmacol.* 81, 11–16.
20. Gupta, S., Abu-Ghannam, N. (2011) Bioactive potential and possible health effects of edible brown seaweeds. *Trends. Food Sci. Technol.* 22, 315–326.
21. Hayat, S., Atta-Rahman, C. M. I., Khan, K. M., Abbaskhan, A. (2002) Two new cinnamic acid esters from marine brown alga *Spatoglossum variabile*. *Chem. Pharm. Bull.* 50, 1297–1299.
22. Hsu, C. Y. (2008) Antioxidant activity of *Pyrrosia petiolosa*. *Fitoterap.* 79, 64–66.
23. Hwang, P., Wu, C., Gau, S., Chien, S., Hwang, D. (2010) Antioxidant and immune-stimulating activities of hot water extract from seaweed *Sargassum hemiphyllum*. *J. Marine Sci. Technol.* 18, 41–46.
24. Kayalvizhi, K., Subramanian, N. V., Boopathy, S., Kathiresan, K. (2014) Antioxidant properties of brown seaweeds *Turbinaria ornata* (Turner) J. Agardh, 1848 and *Padina tetrastratica* (Hauck). *J. Biotechnol. Sci.* 2, 29–37.
25. Kim, S. H., Choi, D. S., Athukorala, Y., Jeon, Y. J., Senevirathne, M., Rha, C. K. (2007) Antioxidant activity of sulfated polysaccharides isolated from *Sargassum fulvellum*. *J. Food Sci. Nutr.* 12, 65–73.
26. Kumar, N. J. I., Kumar, R. N., Bora, A., Amb, M. K., Chakraborty, S. (2009) An evaluation of the pigment composition of eighteen marine macro-algae collected from Okha Coast, Gulf of Kutch, India. *Our Nat.* 7, 48–55.
27. Lamsal, K., Ghimire, B. K., Sharma, P., Ghimirar, A. K., Kim, S. W., Yu, C. Y., Chung, M., Lee, Y. S., Kim, J. S., Shakya, S. R. (2010) Genotoxicity evaluation of the insecticide ethion in root of *Allium cepa* L. *Afr. J. Biotechnol.* 9, 4204–4210.
28. Leandrini de Oliveira, A. L., Da Silva, D. B., Lopes, N. P., Deboni, H. M. (2012) Chemical constituents from red algae *Bostrychia radicans* (Rhodomelaceae): New amides and phenolic compounds. *Quim. Nova* 35, 2186–2188.
29. Li, B., Lu, F., Wei, X., Zhao, R. (2008) Fucoidan: Structure and bioactivity. *Molecules* 13, 1671–1695.
30. Li, K., Li, X. M., Nai-Yun, J., Bin-Gui Wang, B. G. (2008) Bromophenols from the marine red alga *Polysiphonia urceolata* with DPPH radical scavenging activity. *J. Nat. Prod.* 71, 28–30.
31. Liman, R., Akyıl, D., Eren, Y., Konuk, M. (2010) Testing of the mutagenicity and genotoxicity of metolcarb by using both *Ames/Salmonella* and *Allium* test. *Chemosphere* 80, 1056–1061.
32. Lu, Xi, Z., Chun, E. C., Ting, T., Guo, J. W., Hong, G., Xu, L., Yun, Z., Yuan, W., Cheng, C., Pei, M. H. (2011) Anti-cancer effects of polysaccharide and phycocyanin from *Porphyra yezoensis*. *J. Mar. Sci. Technol.* 19, 377–382.
33. Luo, H. Y., Wang, B., Yu, C. G., Qu, Y. L., Su, C. L. (2010) Evaluation of antioxidant activities of five selected brown seaweeds from China. *J. Med. Plants Res.* 4, 2557–2565.
34. Machu, L., Misurcova, L., Ambrozova, J. V., Orsavova, J., Mlcek, J., Sochor, J., Jurikova, T. (2015) Phenolic content and antioxidant capacity in algal food products. *Molecules* 20, 1118–1133.
35. Magalhaes, K. D., Costa, L. S., Fidelis, G. P., Oliveira, R. M., Nobre, L. T. D. B., Santos, N. D., Camara, R. B. G., Albuquerque, I. R. L., Cordeiro, S. L., Sabry, D. A. et al. (2011) Anticoagulant, antioxidant and antitumor activities of heterofucans from the seaweed *Dictyopteria delicatula*. *Int. J. Mol. Sci.* 12, 3352–3365.
36. Malik, C. P., Singh, M. B. (1980) Plant enzymology and histo-enzymology. Kalyani Publishers. New Delhi.
37. Maruhen, A., Datsch, S., Diego, L. R., Thayná, A. S., Gabriela, M. V., Carlye, N. C., Michele, K., Luciana, P., Grégio, A. (2016) Mutagenicity of two herbicides widely used on soybean crops by the *Allium cepa* test. *Cytotechnol* 68, 1215–1222.

38. Metzner, H., Rau, H., Senger, H. (1965) Untersuchungen zur Synchronisierbarkeit einzelner Pigment. Mangol Mutanten Von Chloella. *Plant* 65, 186.
39. Mukherjee, S. P., Choudhuri, M. A. (1983) Implications of water stress-induced changes in the levels of endogenous ascorbic acid and hydrogen peroxide in *Vigna* seedlings. *Physiol. Plant* 58, 166–170.
40. Nagai, T., Yukimoto, T. (2003) Preparation and functional properties of beverages made from sea algae. *Food Chem.* 81, 327–332.
41. Naguib, M. I. (1963) Colorimetric estimation of plant polysaccharides. *Z. Zucker.*, 16, 15–18.
42. Namvar, F., Mohamed, S., Fard, S. G. (2012) Polyphenol-rich seaweed (*Eucheuma cottonii*) extract suppresses breast tumour via hormone modulation and apoptosis induction. *Food Chem.* 130, 376–382.
43. Nwosu, F., Morris, J., Lund, V. A., Stewart, D., Ross, H. A., McDougall, G. J. (2011) Anti-proliferative and potential anti-diabetic effects of phenolic-rich extracts from edible marine algae. *Food Chem.* 126, 1006–1012.
44. Onofrejovaa, L., Vašičkováa, J., Klejdusa, B., Stratil, P., Mišurcova, L., Kračmar, S. C., Kopecky, B. J., Vaceka, J. (2010) Bioactive phenols in algae: The application of pressurized-liquid and solid-phase extraction techniques. *J. Pharm. Biomed. Anal.* 51, 464–470.
45. Osama, A. H., Abu Zinadah, W., Khalil, K. B., El Ashmaouil, M. H., Faiza, A., Abou Alsoud, E. M. (2013) Evaluation of the anti-genotoxicity and growth performance impacts of green algae on *Mugil cephalus*. *Life Sci. J.* 10, 1543–1554.
46. Osuna-Ruiz, I., López-Saiz, C. M., Burgos-Hernández, A., Velázquez, C., Nieves-Soto, M., Hurtado-Oliva, M. A. (2016) Antioxidant, anti-mutagenic and anti-proliferative activities in selected seaweed species from Sinaloa, Mexico. *Pharm. Biol.* 9, 1–15.
47. Sachindra, N. M., Sato, E., Maeda, H., Hosokawa, M., Niwano, Y., Kohno, M., Miyashita, K. (2007) Radical scavenging and singlet oxygen quenching activity of marine carotenoid fucoxanthin and its metabolites. *J. Agri. Food Chem.* 55, 8516–8522.
48. Schoenwaelder, M. E. A. (2008) The biology of phenolic containing vesicles. *Algae* 23, 163–175.
49. Shahin, S. A., El-Amoodi, K. H. H. (1991) Induction of numerical chromosomal aberrations during DNA synthesis using fungicides nimrod and rubigan-4 in root tips of *Vicia faba* L. *Mutat Res.* 261, 169–176.
50. Sobral, O., Marin-Morales, M. A., Ribeiro, R. (2013) Could contaminant induced mutations lead to a genetic diversity over estimation? *Ecotoxicol.* doi:10.1007/s10646-013-1079-4.
51. Souza, M. C. R., Marques, C. T., Dore, C. M. G., Silva, F. R. F., Rocha, H. A. O., Leite, E. L. (2007) Antioxidant activities of sulfated polysaccharides from brown and red seaweeds. *J. Appl. Phycol.* 19, 153–160.
52. Synytsya, A., Kim, W. J., Kim, S. M., Pohl, R., Kvasnicka, F. (2010) Structure and antitumor activity of fucoidan isolated from sporophyll of Korean brown seaweed *Undaria pinnatifida*. *Carbohydr. Polym.* 81, 41–48.
53. Usov, A. I. (2011) Polysaccharides of the red algae. *Adv. Carbohydr. Chem. Biochem.* 65, 115–217.
54. Yvonne, V. Y., Natalie, A. (2006) Antioxidant and anti-proliferative activities of extracts from a variety of edible seaweeds. *Food Chem. Toxicol.* 44, 1144–1150.
55. Zakaria, N. A., Ibrahim, D., Sulaiman, S. F., Supardy, A. (2011) Assessment of antioxidant activity, total phenolic content and in-vitro toxicity of Malaysian red seaweed, *Acanthophora spicifera*. *J. Chem. Pharm. Res.* 3, 182–191.