

# PUBLISHED VERSION


Gudrun Schober, Kylie Lange, Robert E. Steinert, Amy T. Hutchison, Natalie D. Luscombe-Marsh, Maria F. Landrock, Michael Horowitz, Radhika V. Seimon and Christine Feinle-Bisset  
**Contributions of upper gut hormones and motility to the energy intake-suppressant effects of intraduodenal nutrients in healthy, lean men - a pooled-data analysis**  
Physiological Reports, 2016; 4(17):e12943-1-e12943-14

© 2016 The Authors. Physiological Reports published by Wiley Periodicals, Inc. on behalf of the American Physiological Society and The Physiological Society. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Originally published at:  
<http://doi.org/10.14814/phy2.12943>

## PERMISSIONS

<http://creativecommons.org/licenses/by/4.0/>



The image shows the Creative Commons Attribution 4.0 International License (CC BY 4.0) logo and a summary of its terms. The logo consists of the 'CC' symbol and a person icon. Below the logo, the text reads 'Attribution 4.0 International (CC BY 4.0)'. A disclaimer states: 'This is a human-readable summary of (and not a substitute for) the license. Disclaimer.' The main heading is 'You are free to:', followed by two bullet points: 'Share — copy and redistribute the material in any medium or format' and 'Adapt — remix, transform, and build upon the material for any purpose, even commercially.' A circular seal on the right says 'Free Cultural Works APPROVED FOR'. Below this, it states: 'The licensor cannot revoke these freedoms as long as you follow the license terms.' A horizontal line separates this from the next section, 'Under the following terms:'. This section includes a person icon and the text: 'Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.' Below that, it says: 'No additional restrictions — You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.'

14 August 2017

<http://hdl.handle.net/2440/106383>

## ORIGINAL RESEARCH

## Contributions of upper gut hormones and motility to the energy intake-suppressant effects of intraduodenal nutrients in healthy, lean men – a pooled-data analysis

Gudrun Schober<sup>1</sup>, Kylie Lange<sup>1,2</sup>, Robert E. Steinert<sup>1,2</sup>, Amy T. Hutchison<sup>1,2,3</sup>, Natalie D. Luscombe-Marsh<sup>2,4</sup>, Maria F. Landrock<sup>1</sup>, Michael Horowitz<sup>1,2</sup>, Radhika V. Seimon<sup>5</sup> & Christine Feinle-Bisset<sup>1,2</sup>

1 University of Adelaide Discipline of Medicine, Adelaide, Australia

2 NHMRC Centre of Excellence in Translating Nutritional Science to Good Health, University of Adelaide, Adelaide, Australia

3 South Australian Health and Medical Research Institute, Adelaide, Australia

4 CSIRO Animal, Food and Health Sciences, Adelaide, Australia

5 Boden Institute of Obesity, Nutrition, Exercise & Eating Disorders, University of Sydney, Sydney, Australia

### Keywords

Appetite perceptions, determinants of energy intake, cholecystokinin, glucagon-like peptide-1, peptide YY, pyloric pressures.

### Correspondence

Christine Feinle-Bisset, University of Adelaide Discipline of Medicine, Royal Adelaide Hospital, Adelaide, SA 5000, Australia.  
Tel: +61-8-8222-5247  
Fax: +61-8-8223-3870  
E-mail: christine.feinle@adelaide.edu.au

### Funding Information

NHMRC Senior Research Fellowships (grant 627002, 2010–15; grant 1103020, 2016–20; to CF-B), Royal Adelaide Hospital Research Committee Mary Overton Early Career Fellowship (2012–14; to RES), Royal Adelaide Hospital Research Committee project grants 2013 and 2014 (to RES & CF-B), Diabetes Australia Research Trust grant 2014 (to CF-B), Faculty of Health Sciences, The University of Adelaide Postgraduate Scholarship (2010–13; to ATH), NHMRC New Investigator Project grant (grant 627118, 2010–14; to NDL-M); Stiftung zur Förderung der Ernährungsforschung in der Schweiz (Swiss Foundation for Nutrition Research; grant 459, 2015; to GS), and NHMRC Early Career Fellowship (grant 1072771, 2014–17; to RVS).

Received: 12 July 2016; Accepted: 8 August 2016

doi: 10.14814/phy2.12943

*Physiol Rep*, 4 (17), 2016, e12943, doi: 10.14814/phy2.12943

### Abstract

We have previously identified pyloric pressures and plasma cholecystokinin (CCK) concentrations as independent determinants of energy intake following administration of intraduodenal lipid and intravenous CCK. We evaluated in healthy men whether these parameters also determine energy intake in response to intraduodenal protein, and whether, across the nutrients, any predominant gastrointestinal (GI) factors exist, or many factors make small contributions. Data from nine published studies, in which antropyloroduodenal pressures, GI hormones, and GI /appetite perceptions were measured during intraduodenal lipid or protein infusions, were pooled. In all studies energy intake was quantified immediately after the infusions. Specific variables for inclusion in a mixed-effects multivariable model for determination of independent predictors of energy intake were chosen following assessment for collinearity, and within-subject correlations between energy intake and these variables were determined using bivariate analyses adjusted for repeated measures. In models based on all studies, or lipid studies, there were significant effects for amplitude of antral pressure waves, premeal glucagon-like peptide-1 (GLP-1) and time-to-peak GLP-1 concentrations, GLP-1 AUC and bloating scores ( $P < 0.05$ ), and trends for basal pyloric pressure (BPP), amplitude of duodenal pressure waves, peak CCK concentrations, and hunger and nausea scores ( $0.05 < P \leq 0.094$ ), to be independent determinants of subsequent energy intake. In the model including the protein studies, only BPP was identified as an independent determinant of energy intake ( $P < 0.05$ ). No single parameter was identified across all models, and effects of the variables identified were relatively small. Taken together, while GI mechanisms contribute to the regulation of acute energy intake by lipid and protein, their contribution to the latter is much less. Moreover, the effects are likely to reflect small, cumulative contributions from a range of interrelated factors.

## Introduction

In healthy humans, energy intake and expenditure are, in most cases, balanced precisely over long periods of time, so that body weight is stable. This energy homeostasis is controlled by complex interactions between central and peripheral feedback signals, including neurohumoral responses to ingested food (Woods et al. 2000; Cummings and Overduin 2007). The arrival of nutrients in the small intestine modulates a number of gastrointestinal (GI) functions, including gastroduodenal motility (Heddl et al. 1988a; Cook et al. 1997; Pilichiewicz et al. 2007b), associated with a slowing of gastric emptying (Heddl et al. 1989), and the stimulation of putative eating-inhibitory gut hormones, including cholecystokinin (CCK) (Feltrin et al. 2007; Pilichiewicz et al. 2007b), glucagon-like peptide-1 (GLP-1) (Feinle et al. 2002; Pilichiewicz et al. 2007a; Ryan et al. 2013), and peptide YY (PYY) (Feltrin et al. 2007; Pilichiewicz et al. 2007b), as well as the suppression of ghrelin (Parker et al. 2005; Feltrin et al. 2006; Cukier et al. 2008), the only known orexigenic gut hormone (Tschop et al. 2000; Cummings et al. 2001; Wren et al. 2001). The stimulation of phasic and tonic pyloric pressures is critical for the slowing of gastric emptying by nutrients (Heddl et al. 1989). Among the macronutrients, lipid appears to have the most potent effects to modulate GI motor and hormone functions (Andrews et al. 1998; Pilichiewicz et al. 2007a,b; Seimon et al. 2009a; Ryan et al. 2013). Our pooled data analysis of eight studies from our laboratory, in which antropyloroduodenal pressures, GI hormones, and GI/appetite perceptions were measured during intraduodenal lipid, or intravenous CCK, infusions, indicated that the magnitude of the stimulation of pyloric pressures and plasma CCK concentrations is independent determinants of subsequent energy intake in healthy men (Seimon et al. 2010), consistent with the concept that both pyloric pressures and CCK are important, in the acute regulation of energy intake.

Protein is generally regarded as the most satiating macronutrient (Latner and Schwartz 1999; Weigle et al. 2005; Batterham et al. 2006; Westerterp-Plantenga et al. 2009), although recent studies from our laboratory in lean subjects have shown that orally ingested high-protein or high-fat meals (Brennan et al. 2012), and intraduodenal infusions of pure fat or protein (Ryan et al. 2013), have comparable effects to reduce energy intake. However, in these studies, despite equipotent effects on energy intake, the effects of intraduodenal protein to stimulate pyloric pressures and plasma CCK and GLP-1 concentrations were much less than those of intraduodenal lipid, whereas protein had more potent effects to stimulate plasma insulin and glucagon (Ryan et al. 2013). These observations,

accordingly, suggest that the energy intake-suppressant effects of protein may not be mediated by these GI mechanisms to the same extent as for lipid, which warrants further evaluation. Over the last years, we have performed a number of studies relating to the effects of protein and amino acids on GI function and energy intake (Ryan et al. 2012, 2013; Steinert et al. 2014, 2015). This analysis was stimulated by the recognition that, in our previous pooled analysis (Seimon et al. 2010), (1) effects of protein were not evaluated; and (2) we included studies utilizing intravenous hormone infusions, particularly exogenous CCK, which is known to potently stimulate pyloric pressures and increase plasma CCK concentrations to supra-physiological levels, and, thus, may have led to an overestimation of the role of these two outcomes. Because of the potent energy intake-suppressant, but more modest GI, effects of protein, we hypothesized, based on our recent study (Ryan et al. 2013), that, in contrast to lipid, GI mechanisms would not be identified as independent determinants of energy intake in the varying protein interventions/loads used.

## Methods

### Subjects

A total of 117 healthy, normal-weight men, aged  $26 \pm 3$  years, and with a BMI of  $22.6 \pm 0.8$  kg/m<sup>2</sup>, were included in this pooled data analysis (Feltrin et al. 2004, 2006, 2008; Little et al. 2005; Pilichiewicz et al. 2007b; Seimon et al. 2009a; Ryan et al. 2012, 2013; Steinert et al. 2014, 2015) (Table 1). All participants were unrestrained eaters with a score of  $\leq 12$  on the eating-restraint component of the three-factor eating questionnaire (Stunkard and Messick 1985). Subjects who smoked, consumed  $>20$  g of alcohol/day, had a history of GI symptoms, or took medication known to affect appetite, eating, or GI function were excluded. Each subject provided informed, written consent before their inclusion, and the Royal Adelaide Research Ethics Committee approved the study protocols, and studies were performed in accordance with the Declaration of Helsinki.

### Study design

Data from nine published studies (Feltrin et al. 2004, 2006, 2008; Little et al. 2005; Pilichiewicz et al. 2007b; Seimon et al. 2009a; Ryan et al. 2012, 2013; Steinert et al. 2014, 2015), performed in our laboratory between 2003–13, were pooled for this analysis. All studies used identical methods and techniques and evaluated the outcome measures of interest (Table 2). While, as in our previous pooled analysis (Seimon et al. 2010), data were analyzed

**Table 1.** Subject and protocol details for each study included in the data analysis.

Study no.	Authors (References)	Subject (n)	Age (year)	BMI (kg/m <sup>2</sup> )	Study intervention
1	Feltrin et al. (2004, 2006) <sup>1</sup>	7 <sup>2</sup>	24 ± 4	22.0 ± 1.6	ID saline (control), 0.375 kcal/min decanoic or dodecanoic acid for 90 min. Buffet meal at 90 min.
2	Little et al. (2005)	13	23 ± 2	23.6 ± 0.5	ID saline, 0.1, 0.2, or 0.4 kcal/min dodecanoic acid for 90 min. Buffet meal at 90 min.
3	Pilichiewicz et al. (2007b)	16	31 ± 3	23.8 ± 0.5	ID saline, 0.25, 1.5, or 4 kcal/min of 10% Intralipid for 50 min. Buffet meal at 50 min.
4	Feltrin et al. (2008)	13	26 ± 2	22.9 ± 0.6	ID saline, 0.4 kcal/min of dodecanoic, or oleic acid for 60 min. Buffet meal at 60 min.
5	Seimon et al. (2009b)	10	25 ± 3	22.8 ± 0.4	ID saline, 2.8 kcal/min of fat emulsions with droplet sizes of 0.26, 30, or 170 μm for 120 min. Buffet meal at 120 min.
6	Ryan et al. (2012)	16	27 ± 3	22.1 ± 0.6	ID saline, 0.5, 1.5, or 3 kcal/min of 18.1% hydrolyzed whey protein for 60 min. Buffet meal at 60 min.
7	Ryan et al. (2013)	20	27 ± 3	22.4 ± 0.4	ID saline, 3 kcal/min of 18.1% hydrolyzed whey protein or 20% Intralipid for 90 min. Buffet meal at 90 min.
8	Steinert et al. (2014)	10	27 ± 9	22.5 ± 2.1	ID saline, 0.075 or 0.15 kcal/min of L-tryptophan for 90 min. Buffet meal at 90 min.
9	Steinert et al. (2015)	12	25 ± 2	21.9 ± 0.4	ID saline, 0.15 or 0.45 kcal/min of L-leucine for 90 min. Buffet meal at 90 min.

ID, intraduodenal. Data are means ± SEM.

<sup>1</sup>Parts of the hormone data were analyzed and published separately, which resulted in two publications.

<sup>2</sup>Plasma was available in only seven of the eight subjects included in the analyses.

using the same statistical tests that would be appropriate for an individual participant meta-analysis, it would be inappropriate to refer to our analysis as such because the studies included were not identified through a systematic review (Simmonds et al. 2005).

### Study protocols

Each study assessed the effects of ID infusions of nutrients (e.g., triglyceride or fatty acids, protein or amino acids) and appropriate control solutions on antropyloroduodenal pressures, GI hormone release, GI/appetite perceptions, and energy intake. Information regarding the study interventions in each study is provided in Table 1. Detailed study protocols have been described in the original publications (Feltrin et al. 2004, 2006, 2008; Little et al. 2005; Pilichiewicz et al. 2007b; Seimon et al. 2009a; Ryan et al. 2012, 2013; Steinert et al. 2014, 2015). Briefly, in all studies, participants attended the laboratory in the Discipline of Medicine, Royal Adelaide Hospital, at 0830 h after an overnight fast, for multiple study visits in randomized order. On arrival, a small-diameter manometric catheter (outer diameter: 3.5 mm; total length: 100 cm; Dentsleeve International Ltd, Mississauga, Ontario, Canada) was inserted through an anesthetized nostril

into the stomach and allowed to pass into the duodenum by peristalsis (Heddle et al. 1989). The catheter consisted of 16 side holes, spaced at 1.5 cm intervals, to measure pressures in the antrum (channels 1–6), pylorus (channels 7–9; 4.5-cm sleeve sensor), and duodenum (channels 10–16), and an additional side hole located in the duodenum for ID infusions. An intravenous cannula was placed into a right forearm vein for regular blood sampling for subsequent measurements of plasma GI hormone, and/or insulin, concentrations. Once the catheter was placed correctly, as described (Heddle et al. 1988a), fasting motility was observed until the occurrence of phase III of the interdigestive migrating motor complex. Immediately after the end of phase III activity, in phase I, a baseline blood sample was taken, and the subject completed a visual analog scale (VAS) questionnaire to assess GI perceptions (Parker et al. 2004). ID infusion of nutrient or control solutions commenced, and antropyloroduodenal pressures were recorded throughout the infusion period. Further blood samples were collected, and VAS questionnaires completed, at regular intervals. At the end of each ID infusion, the catheter and cannula were removed, and the subject was presented with a standardized, cold, buffet-style meal (Nair et al. 2009). The meal consisted of white and whole-meal breads, sliced, cold meats (ham,

**Table 2.** Variables measured in each study.

Study no.	Authors (References)	APD pressures					Hormones					Perceptions					
		APWs	IPPWs	BPP	DPWs	Ghr	CCK	PYY	GLP-1	Insulin	Glu <sup>2</sup>	Hunger	Desire to eat	Prosp. cons.	Fullness	Nausea	Bloating
1	Feltrin et al. (2004)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
2	Feltrin et al. (2006) <sup>1</sup>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
3	Little et al. (2005)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
4	Plichewicz et al. (2007b)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
5	Feltrin et al. (2008)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
6	Seimon et al. (2009b)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
7	Seimon et al. (2012)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
8	Ryan et al. (2013)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
9	Steinert et al. (2014)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
9	Steinert et al. (2015)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

APD, antrotyloroduodenal; APWs, antral pressure waves; IPPWs, isolated pyloric pressure waves; BPP, basal pyloric pressure; DPWs, duodenal pressure waves; Ghr, Ghrelin; CCK, cholecystokinin; GLP-1, glucagon-like peptide 1; PYY, peptide YY; Glu, Glucagon; Prosp. cons., prospective consumption; X, variable measured.

<sup>1</sup>Parts of the hormone data were analyzed and published separately, which resulted in two publications.

<sup>2</sup>Not included in the subsequent statistical analyses because glucagon was only evaluated in a small number of studies.

chicken), cheese, lettuce, tomato, cucumber, mayonnaise, butter, apple, banana, yoghurt, chocolate custard, fruit salad, iced coffee, orange juice, and water. Subjects were allowed to eat as much as they wished for up to 30 min, until they felt comfortably full.

## Measurements

### APD pressures

APD pressures were digitized using a computer-based system running commercially available software and analyzed for (1) the number and amplitude of antral and duodenal pressure waves; (2) basal pyloric pressure (BPP; “tone”); and (3) the number and amplitude of isolated pyloric pressure waves (IPPWs) (Hedde et al. 1988b). Antral and duodenal pressure waves were expressed as total numbers and mean amplitudes (mmHg). For IPPWs, number of premeal IPPWs (i.e., during the 15-min period immediately before the buffet meal), the peak number, and the time (min)-to-peak number during ID infusions, total number, and areas under the curve (AUCs, calculated using the trapezoidal rule) of total number (min) and amplitude (mmHg·min) were quantified. BPP was expressed as peak pressures (mmHg), time (min)-to-peak pressure, and AUC (mmHg·min) over the infusion period.

### Hormone concentrations

Venous blood samples were collected, processed, and stored for later analysis of total ghrelin, CCK, total PYY, total GLP-1, and insulin. Plasma total ghrelin (pmol/L) was measured by radioimmunoassay, without peptide extraction (Phoenix Pharmaceuticals, Mountain View, CA). No cross-reactivities with any relevant molecule have been reported. Intra- and interassay coefficients of variation (CVs) were 5.0% and 15.0%, respectively. The detection limit was 13 pmol/L. Plasma CCK-8 (pmol/L) was measured by radioimmunoassay after ethanol extraction using an adaptation of a previously published method (Santangelo et al. 1998). Intra- and interassay CVs were 8.3% and 12.6%, respectively. The detection limit was 1 pmol/L. Plasma total GLP-1 (pmol/L) was measured by radioimmunoassay (Millipore, Billerica, MA). The antibody used did not cross-react with glucagon, gastric inhibitory polypeptide, or other gut or pancreatic peptides. Intra- and interassay CVs were 4.8% and 6.8%, respectively. The detection limit was 3 pmol/L. Plasma total PYY (pmol/L) was measured by radioimmunoassay (Linco Research, St. Charles, MO). Intra- and interassay CVs were 5.3% and 7.0%, respectively. The detection limit was 2.5 pmol/L. Plasma insulin (mU/L) was

measured by an ELISA assay (Mercodia, Uppsala, Sweden). Intra- and interassay CVs were 2.9% and 8.8%, respectively. The detection limit was 1 mU/L.

For CCK, premeal and peak concentrations (pmol/L), the time (min)-to-peak concentrations, and the AUC (pmol·min/L) over the infusion period were calculated. For ghrelin, PYY, GLP-1, and insulin, premeal concentrations (pmol/L or mU/L, as appropriate) and AUC (pmol·min/L or mU·min/L, as appropriate) were calculated. Peak concentrations were not calculated for ghrelin, PYY, GLP-1, and insulin because concentrations were shown to continue to rise, or decline (for ghrelin), throughout the infusion period.

### GI/appetite perceptions and energy intake

Hunger, desire to eat, prospective consumption, fullness, nausea, and bloating were rated using validated 100-mm VAS scales (Parker *et al.* 2004) and expressed as AUCs over the infusion period (mm·min). Energy intake (kJ) was determined from the food consumed (g) at the buffet meal, by weighing the different food items before and after consumption, and then calculating their energy content using commercially available software (Foodworks; Xyris Software, Highgate Hill, QLD, Australia) (Feltrin *et al.* 2004; Ryan *et al.* 2013).

### Statistical analysis

Statistical analyses were performed using SPSS 22 software (2013, IBM Corp, Armonk, NY).

Within-subject correlations, adjusted for repeated measures, were performed to assess the strength of bivariate relations between energy intake and each hormone, motility, and GI/appetite perception variable (Bland and Altman 1995). Variables were then entered simultaneously into a multivariable maximum-likelihood linear mixed-effects model, adjusted for repeated visits per subjects and the clustering of subjects within studies, to determine the independent effects of each variable on energy intake (Raudenbush and Bryk 2002). This is equivalent to the one-step analysis approach in a meta-analysis of individual participant data (Riley *et al.* 2010).

Analyses were conducted for all studies combined (models 1a, 1b), for the lipid-based studies (models 2a, 2b), and for the protein-based studies (model 3). Because GLP-1 and PYY were not measured in all studies, the multivariable analyses were conducted as two separate models for 'all studies' and for the 'lipid studies'. Models 1a and 2a considered all variables measured across all studies plus PYY. Models 1b and 2b considered all variables measured across all studies plus GLP-1. Variables were included in the multivariable models after screening

for multicollinearity and excluding related parameters from the same underlying motility, hormone, or GI/appetite perception variable, as indicated by the variance inflation factors and condition indices. Where related variables were collinear, parameters were selected for inclusion based on how they best reflected the observed physiological response over the study periods. These selections were made without referring to the model results to avoid post hoc variable selection. Further exclusion of variables was conducted for the lipid and protein subgroups, to reduce the models to a number of parameters appropriate for the reduced sample sizes. For this purpose, F-statistics were used to exclude the variables that least contributed to the model, analogous to backward variable selection. Thus, the following variables were excluded in the process: IPPW time-to-peak number, IPPW number AUC, BPP time-to-peak pressure, CCK premeal, and CCK time to peak. Based on the same principles, further variables were excluded to reach a final model with an appropriate number of parameters, including, from the lipid subgroup, mean amplitude of antral pressure waves, IPPW amplitude AUC, BPP AUC, mean amplitude of duodenal pressure waves, AUCs for CCK, PYY, GLP-1, prospective consumption and fullness, and from the protein subgroup, additionally IPPW peak number, number of duodenal pressure waves, and desire to eat AUC. At each step, the models were examined for consistency and robustness of the parameter estimates.

Data are expressed as means  $\pm$  standard error of the mean (SEM) or standard deviation (SD), as indicated, and differences were considered significant when  $P < 0.05$ .

## Results

### Bivariate correlation analyses

Within-subject correlations between energy intake and each of the measured variables are presented in Table 3. Collinearity was present between a number of variables; thus, related variables from the same underlying motility, hormone, or GI/appetite perception variable were excluded from the multivariate model to guarantee a robust estimation of the regression effects.

### Models 1a and 1b (all studies)

Within the variables characterizing IPPWs, peak number, total number, and AUC of the number were strongly associated with each other (all  $r \geq 0.83$ ). Of these variables, AUC of the number was excluded from all multivariate models because it least characterized the IPPW response. Among the CCK variables, peak concentration

**Table 3.** Within-subject correlations between energy intake and GI motor, hormone, and GI/appetite perception variables.

Variable	All studies			Lipid studies			Protein studies			
	n <sup>1</sup>	Mean ± SD	r	n <sup>1</sup>	Mean ± SD	r	n <sup>1</sup>	Mean ± SD	r	P
<b>Antral pressure waves</b>										
Number, total	400	59 ± 86	0.14	256	57 ± 98	0.13	164	77 ± 80	0.21	0.028
Amplitude, mean (mmHg)	400	62 ± 80	0.23	256	52 ± 87	0.22	164	104 ± 105	0.26	0.006
<b>Isolated pyloric pressure waves</b>										
Number premeal/15 min	384	9 ± 12	-0.17	240	9 ± 12	-0.09	164	8 ± 11	-0.32	0.001
Peak number/15 min	384	20 ± 14	-0.24	240	21 ± 14	-0.22	164	17 ± 12	-0.34	<0.001
Time-to-peak number (min)	359	29 ± 22	0.05	216	28 ± 22	0.18	163	31 ± 23	-0.16	0.104
Total number	400	53 ± 60	-0.19	256	56 ± 67	-0.14	164	45 ± 43	-0.38	<0.001
AUC number (min)	384	727 ± 784	-0.21	240	815 ± 886	-0.18	164	540 ± 530	-0.38	<0.001
AUC amplitude (mmHg·min)	384	1699 ± 1105	-0.10	240	1863 ± 1126	-0.01	164	1404 ± 1021	-0.24	0.010
<b>Basal pyloric pressures</b>										
Peak pressure (mmHg)	393	4.0 ± 5.0	-0.14	249	4.4 ± 5.4	-0.13	164	3.0 ± 4.1	-0.28	0.003
Time-to-peak pressure (min)	387	31 ± 24	0.03	243	29 ± 24	0.08	164	37 ± 25	0.00	0.971
AUC (mmHg·min)	393	62 ± 206	-0.11	249	77 ± 232	-0.12	164	38 ± 148	-0.16	0.088
<b>Duodenal pressure waves</b>										
Number, total	397	450 ± 336	0.21	253	451 ± 376	0.28	164	480 ± 270	0.14	0.153
Amplitude, mean (mmHg)	400	56 ± 47	0.11	256	44 ± 43	0.14	164	87 ± 53	0.12	0.213
<b>Ghrelin</b>										
Premeal (pmol/L)	181	266 ± 213	0.34	47	199 ± 155	0.62	147	275 ± 221	0.25	0.012
AUC (pmol·min/L)	181	18,866 ± 13,491	0.34	47	19,034 ± 13,164	0.59	147	18,120 ± 13,242	0.22	0.028
<b>Cholecystokinin</b>										
Premeal (pmol/L)	382	4.6 ± 3.4	-0.35	232	5.1 ± 3.9	-0.37	170	3.6 ± 2.1	-0.45	<0.001
Peak concentration (pmol/L)	382	6.0 ± 4.5	-0.25	232	6.9 ± 5.3	-0.26	170	4.4 ± 2.3	-0.43	<0.001
Time-to-peak conc (min)	352	29 ± 25	-0.19	202	27 ± 25	-0.18	168	32 ± 24	-0.24	0.012
AUC (pmol·min/L)	359	338 ± 265	-0.24	209	392 ± 325	-0.27	170	254 ± 111	-0.45	<0.001
<b>Peptide YY</b>										
Premeal (pmol/L)	309	42.9 ± 27.1	-0.25	166	47.7 ± 33.6	-0.30	156	36.4 ± 14.8	-0.30	0.002
AUC (pmol·min/L)	309	5260 ± 7830	-0.14	166	7544 ± 10,123	-0.20	156	2580 ± 838	-0.23	0.016
<b>Glucagon-like peptide-1</b>										
Premeal (pmol/L)	263	31.0 ± 19.8	-0.26	113	29.0 ± 25.7	-0.21	170	31.5 ± 13.5	-0.50	<0.001
Time-to-peak conc (min)	241	50.9 ± 24.1	-0.20	102	51.5 ± 24.0	-0.31	156	49.9 ± 24.2	-0.15	0.148
AUC (pmol·min/L)	263	2118 ± 1104	-0.17	113	2152 ± 1502	-0.22	170	2099 ± 677	-0.35	<0.001
<b>Insulin</b>										
Premeal (mU/L)	190	8.0 ± 10.8	-0.40	40	3.5 ± 2.0	-0.40	170	8.5 ± 11.3	-0.44	<0.001
AUC (mU·min/L)	190	520 ± 600	-0.42	40	351 ± 173	-0.38	170	525 ± 632	-0.47	<0.001
<b>GI/appetite perceptions (mm·min)</b>										
AUC Hunger	390	-398 ± 1405	0.11	240	-682 ± 1545	0.14	170	51 ± 1046	0.02	0.855
AUC Desire to eat	390	-522 ± 1393	0.11	240	-857 ± 1538	0.18	170	28 ± 947	-0.04	0.702

(Continued)

Table 3. Continued.

Variable	All studies			Lipid studies			Protein studies			
	<i>n</i> <sup>1</sup>	Mean ± SD	<i>r</i>	<i>n</i> <sup>1</sup>	Mean ± SD	<i>r</i>	<i>n</i> <sup>1</sup>	Mean ± SD	<i>r</i>	<i>P</i>
AUC Prospective consumption	390	-597 ± 1302	0.16	240	-947 ± 1427	0.25	0.001	-63 ± 846	-0.02	0.835
AUC Fullness	390	1027 ± 1338	-0.10	240	1420 ± 1470	-0.11	0.164	437 ± 817	-0.08	0.381
AUC Nausea	390	224 ± 907	-0.23	240	277 ± 1084	-0.31	<0.001	142 ± 495	-0.17	0.065
AUC Bloating	390	558 ± 1205	-0.27	240	781 ± 1403	-0.37	<0.001	236 ± 727	-0.06	0.517

AUC, area under the curve; GI, gastrointestinal

<sup>1</sup>Variations in the number of study visits (*n*) for the various variables due to missing data

and AUC were strongly correlated ( $r = 0.87$ ), and peak concentration was included because it best described the hormone profile.

### Models 2a and 2b (lipid studies)

Of the variables characterizing IPPWs in these studies, peak number, total number, and AUC of the number were strongly associated with each other (all  $r \geq 0.82$ ); thus, AUC of the number was excluded because it least characterized the IPPW response. For the BPP variables, peak pressure and AUC were strongly correlated ( $r = 0.85$ ); thus, peak pressure was included because it showed a slightly stronger correlation with energy intake. Among the CCK variables, peak concentration and AUC were strongly correlated ( $r = 0.88$ ), and peak concentration was included because it best characterized the CCK response. Within the GI/appetite perception variables, AUCs for hunger, desire to eat, and prospective consumption were correlated moderately with each other (all  $r \leq 0.77$ ); thus, AUC for prospective consumption was excluded.

### Model 3 (protein studies)

All variables characterizing IPPWs were very strongly correlated ( $r \geq 0.89$ ); thus, total number was included in the multivariate model because it showed the highest F-statistics. Of the BPP variables, peak pressure and AUC were strongly correlated ( $r = 0.70$ ), and peak pressure was selected for inclusion because it showed a stronger correlation with energy intake. Within the CCK variables, peak concentration and AUC were strongly correlated ( $r = 0.91$ ); thus, peak concentration was included because it best characterized the CCK response. AUCs for hunger, desire to eat, and prospective consumption were modestly associated with each other (all  $r \leq 0.66$ ); thus, hunger was included based on the highest F-statistics. GLP-1 and PYY premeal concentrations were strongly correlated with the corresponding AUCs ( $r \geq 0.88$ ); premeal concentrations were included because they showed stronger correlations with energy intake.

None of the other variables showed any multicollinearity, therefore, they were entered into the multivariable models automatically. For the protein and lipid-based studies, additional variables were excluded on the basis of clinical utility and the smallest F-statistics to reduce the models to a number of parameters appropriate for the reduced sample sizes.

### Multivariable mixed-effects models

Model 1a (all variables measured across all studies plus PYY) identified antral amplitude ( $P < 0.05$ ), a trend for



peak BPP ( $P = 0.082$ ), as well as bloating ( $P < 0.05$ ) as predictors of energy intake (Table 4); increases in antral amplitude or peak BPP by 1 mmHg, or in bloating by 1 mm·min, while controlling for all the other variables, were associated with an increase of 2.4 kJ, or decreases of ~37 kJ or ~0.2 kJ in energy intake, respectively. Model 1b (all variables measured across all studies plus GLP-1) identified a trend for peak CCK concentration ( $P = 0.074$ ), premeal GLP-1 concentration ( $P < 0.01$ ), and GLP-1 AUC ( $P < 0.01$ ), and a trend for hunger ( $P = 0.09$ ) as determinants of energy intake. Increases in CCK or GLP-1 by 1 pmol/L, in GLP-1 AUC by 1 pmol·min/L or in hunger AUC by 1 mm·min were associated with reductions of ~66, ~27, and ~0.6 kJ, or an increase of ~0.3 kJ, in energy intake, respectively.

Model 2a (lipid-based studies including PYY) identified trends for number of premeal IPPWs ( $P = 0.057$ ), peak BPP ( $P = 0.067$ ), and duodenal amplitude ( $P = 0.072$ ), as well as bloating ( $P < 0.001$ ), as predictors of energy intake, so that increases by  $n = 1$ , 1 mmHg, or 1 mm·min were associated with decreases of 15 kJ or 32 kJ, an increase of 5.4 kJ, or a decrease of ~0.2 kJ, in energy intake, respectively (Table 4). In model 2b (lipid-based studies including GLP-1), increases in peak BPP and plasma CCK peak concentration were associated with trends for an increase ( $P = 0.055$ ), or a decrease ( $P = 0.066$ ), in energy intake, respectively; an increase in peak BPP by 1 mmHg was associated with an increase in energy intake of ~69 kJ, whereas an increase in plasma CCK peak concentration by 1 pmol/L reduced energy intake by ~80 kJ. The time-to-peak concentration of plasma GLP-1 was also associated with energy intake ( $P < 0.05$ ); an increase by 1 min reduced energy intake by ~11 kJ. Finally, there were trends for hunger ( $P = 0.061$ ) and nausea ( $P = 0.094$ ) to be associated with energy intake; increases in the hunger or nausea AUC by 1 mm·min were associated with an increase, or decrease, in energy intake of 0.4 kJ, or 0.2 kJ, respectively.

Model 3 (all variables measured in the protein-based studies) only identified peak BPP as an independent determinant of energy intake ( $P < 0.05$ ; Table 4); thus, an increase by 1 mmHg was associated with a decrease in energy intake of ~48 kJ.

## Discussion

This study has evaluated, using a pooled-data analysis, the GI motor and hormone determinants of acute energy intake in response to intraduodenal administration of nutrients, particularly protein and lipid. The major findings are that the models, based either on all studies or the lipid studies, identified a range of GI factors, including amplitude of antral pressure waves, premeal GLP-1 and

time-to-peak GLP-1 concentrations, GLP-1 AUC and bloating scores, and trends for BPP, amplitude of duodenal pressure waves, peak CCK concentrations, and hunger and nausea scores, as independent determinants of energy intake. In contrast, in the model that included the protein studies only BPP was identified as an independent determinant of energy intake. Moreover, the effects of variables identified as independent determinants were small. That said, it is important to recognize that the reported effects relate to a change of 1 unit for each variable, for example, an increase in plasma CCK by 1 pmol/L was associated with a reduction in energy intake of ~66–80 kJ across models, and lipid-stimulated plasma CCK may increase by 3–5 pmol/L. Taken together, our observations indicate that (i) the contributions of GI factors to energy intake regulation are nutrient specific and (ii) while GI mechanisms are important in the regulation of acute energy intake (Cummings and Overduin 2007; Steinert et al. 2013), this effect most likely reflects the outcome of relatively small, cumulative contributions from a range of interrelated factors. A clear implication is that ‘GI’ strategies for the management of obesity are unlikely to prove effective if only one mechanism is targeted.

The studies included in this analysis utilized intraduodenal nutrient infusions, which allow the delivery of nutrients into the proximal small intestine (the major location of nutrient sensors, from which signals to initiate feedback control of upper GI motility, slowing of gastric emptying, and gut hormone release, are initiated) to be standardized. The rationale for this approach is that it excludes the confounding effects of the substantial interindividual variation in gastric emptying and ‘bypasses’ gastric distension which modulates energy intake. In addition, the delivery of nutrients prior to a meal “mimics” the preload concept, which aims to administer nutrients that stimulate specific GI functions to influence outcomes (e.g., energy intake or blood glucose) in response to that meal (Gentilcore et al. 2006). We infused long-chain triglyceride emulsions, whey protein, or their digestive products, fatty acids, and amino acids, as these have consistently been shown, in studies from our laboratory and those of others, to suppress subsequent energy intake in humans when given acutely (Matzinger et al. 2000; Feltrin et al. 2004; Ryan et al. 2013).

Our main hypothesis was that with protein, unlike lipid, GI factors were not the major driving force in determining subsequent acute energy intake. Indeed, our analysis identified a range of factors as contributors to energy intake regulation in response to intraduodenal lipid, whereas only a single factor, BPP, was identified for protein, suggesting that other factors are more important in the case of protein. Although these could not be determined in this study, important roles for, for example,

**Table 4.** Results of mixed-effects multivariable models for determination of independent predictors of energy intake.

Variable	All studies						Lipid studies						Protein studies					
	Model 1a (incl PYY)			Model 1b (incl GLP-1)			Model 2a (incl PYY)			Model 2b (incl GLP-1)			Model 3 (incl PYY & GLP-1)					
	$\beta$	<i>t</i>	<i>P</i>	<i>b</i> $\pm$ SEM	$\beta$	<i>t</i>	<i>P</i>	<i>b</i> $\pm$ SEM	$\beta$	<i>t</i>	<i>P</i>	<i>b</i> $\pm$ SEM	$\beta$	<i>t</i>	<i>P</i>			
<b>Antral pressure waves</b>																		
Number, total	-1.4 $\pm$ 1.2	-0.06	-1.14	0.256	-0.4 $\pm$ 1.0	-0.03	-0.44	0.657	-1.7 $\pm$ 1.5	-0.09	-1.12	0.265	-	-	-	-		
Amplitude, mean (mmHg)	2.4 $\pm$ 1.0	0.13	2.44	0.015	1.6 $\pm$ 1.2	0.09	1.38	0.170	-	-	-	-	-	-	-	-		
<b>Isolated pyloric pressure waves</b>																		
Number premeal/15 min	4.9 $\pm$ 7.2	0.04	0.68	0.494	-0.04 $\pm$ 12.3	0.00	0.00	0.997	-15 $\pm$ 8.0	0.13	1.92	0.057	-	-	-	-		
Peak number/15 min	-12.2 $\pm$ 8.4	-0.11	-1.44	0.150	-9.9 $\pm$ 13.4	-0.08	-0.74	0.463	-15.2 $\pm$ 9.3	-0.14	-1.63	0.105	-8.1 $\pm$ 19.3	-0.07	-0.42	0.675		
Time-to-peak number (min)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Total number	0.04 $\pm$ 2.4	0.00	0.02	0.987	0.2 $\pm$ 4.3	0.01	0.06	0.956	1.2 $\pm$ 2.3	0.06	0.53	0.598	1.9 $\pm$ 4.3	0.07	0.44	0.664		
AUC number (min)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
AUC amplitude (mmHg min)	-0.02 $\pm$ 0.07	-0.02	-0.32	0.752	-0.01 $\pm$ 0.09	-0.01	-0.12	0.906	-	-	-	-	-	-	-	-		
<b>Basal pyloric pressures</b>																		
Peak pressure (mmHg)	-37 $\pm$ 21	-0.11	-1.75	0.082	-28 $\pm$ 29	-0.07	-0.95	0.346	-32 $\pm$ 17.0	-0.10	-1.85	0.067	69 $\pm$ 35	0.17	1.96	0.055		
Time-to-peak pressure (min)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
AUC (mmHg min)	0.28 $\pm$ 0.51	0.03	0.55	0.584	0.77 $\pm$ 0.71	0.08	1.09	0.277	-	-	-	-	-	-	-	-		
<b>Duodenal pressure waves</b>																		
Number, total	-0.07 $\pm$ 0.29	-0.01	-0.25	0.806	0.05 $\pm$ 0.3	0.01	0.14	0.885	0.1 $\pm$ 0.34	0.02	0.29	0.770	0.3 $\pm$ 0.3	0.08	0.99	0.329		
Amplitude, mean (mmHg)	-1.7 $\pm$ 2.5	-0.05	-0.68	0.503	-0.3 $\pm$ 2.8	-0.01	-0.11	0.911	5.4 $\pm$ 3.0	0.16	1.81	0.072	-	-	-	-		
<b>Ghrelin</b>																		
Premeal (pmol/L)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
AUC (pmol min/L)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
<b>Cholecystokinin</b>																		
Premeal (pmol/L)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Peak concentration (pmol/L)	-1.8 $\pm$ 19.2	-0.01	-0.09	0.927	-66 $\pm$ 37	-0.13	-1.80	0.074	-3 $\pm$ 18.0	-0.01	-0.18	0.860	-80 $\pm$ 43	-0.20	-1.87	0.066		
Time-to-peak conc (min)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
AUC (pmol min/L)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

(Continued)

Table 4. Continued.

Variable	All studies																				
	Model 1a (incl PYY) (n = 288)				Model 1b (incl GLP-1) (n = 263)				Model 2a (incl PYY) (n = 142)				Model 2b (incl GLP-1) (n = 82)				Model 3 (incl PYY & GLP-1) (n = 170)				
	b ± SEM	β	t	P	b ± SEM	β	t	P	b ± SEM	β	t	P	b ± SEM	β	t	P	b ± SEM	β	t	P	
Peptide YY																					
Premeal (pmol/L)	-3.0 ± 3.7	-0.05	-0.81	0.417	-	-	-	-	-3.2 ± 4.1	-0.07	-0.78	0.436	-	-	-	-	-2.2 ± 9.4	-0.02	-0.23	0.819	-
AUC (pmol·min/L)	0.02 ± 0.02	0.09	0.98	0.330	-	-	-	-	0.01 ± 0.02	0.09	0.68	0.498	-	-	-	-	-	-	-	-	-
Glucagon-like peptide-1																					
Premeal (pmol/L)	-	-	-	-	-27.1 ± 9.9	-0.33	-2.75	0.007	-	-	-	-	4.4 ± 5.8	0.07	0.76	0.451	-8.7 ± 13.6	-0.07	-0.64	0.525	-
Time-to-peak conc (min)	-	-	-	-	-2.0 ± 3.4	-0.03	-0.58	0.560	-	-	-	-	-11 ± 4.0	-0.16	-2.46	0.018	4.9 ± 4.4	0.07	1.12	0.268	-
AUC (pmol·min/L)	-	-	-	-	0.57 ± 0.19	0.38	2.97	0.003	-	-	-	-	-	-	-	-	-	-	-	-	-
Insulin																					
Premeal (mU/L)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-17 ± 11	-0.13	-1.55	0.125	-
AUC (mU·min/L)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
GI/appetite perceptions (mm·min)																					
AUC Hunger	0.01 ± 0.08	0.01	0.15	0.881	0.25 ± 0.15	0.19	1.68	0.094	-0.01 ± 0.08	-0.01	-0.13	0.894	0.4 ± 0.2	0.41	1.91	0.061	0.1 ± 0.1	0.06	0.89	0.378	-
AUC Desire	-0.07 ± 0.11	-0.07	-0.70	0.483	-0.11 ± 0.16	-0.09	-0.68	0.499	-0.04 ± 0.09	-0.04	-0.43	0.670	-0.3 ± 0.2	-0.29	-1.30	0.198	-	-	-	-	-
to eat																					
AUC Prospective consumption	0.09 ± 0.11	0.08	0.87	0.385	0.004 ± 0.15	0.00	0.03	0.978	-	-	-	-	-	-	-	-	-	-	-	-	-
AUC Fullness	-0.03 ± 0.07	-0.03	-0.51	0.612	-0.07 ± 0.11	-0.05	-0.62	0.537	-	-	-	-	-	-	-	-	-	-	-	-	-
AUC Nausea	-0.11 ± 0.08	-0.07	-1.39	0.166	-0.19 ± 0.12	-0.12	-1.63	0.105	-0.12 ± 0.07	-0.10	-1.56	0.122	-0.2 ± 0.1	-0.20	-1.71	0.094	-0.1 ± 0.2	-0.04	-0.64	0.522	-
AUC Bloating	-0.17 ± 0.07	-0.14	-2.49	0.014	0.03 ± 0.13	0.02	0.24	0.811	-0.23 ± 0.06	-0.25	-3.67	0.000	-0.1 ± 0.2	-0.13	-0.81	0.421	-0.1 ± 0.2	-0.03	-0.31	0.756	-

n, number of visits; AUC, area under the curve; GI, gastrointestinal; GLP-1, glucagon-like peptide-1; PYY, peptide Y.

<sup>1</sup>Only included in "Protein studies" model due to insufficient data from lipid studies.

diet-induced thermogenesis (Westerterp-Plantenga et al. 1999), intestinal gluconeogenesis (Mithieux 2009), and direct effects of elevated plasma amino acid concentrations within the brain (Tome 2004), have been identified by others.

The increased recognition of the fundamental role for the upper GI tract in regulating energy intake, but also blood glucose, has prompted extensive research into potential therapeutic agents that target these mechanisms. Among the gut peptides, CCK, PYY, and GLP-1 have received much interest, due to their satiating effects when infused intravenously in healthy or obese humans, however, as has been shown in the case of PYY(3-36), the satiating effects could only be achieved at doses that elevated plasma PYY concentrations to supraphysiological levels and also induced nausea (Degen et al. 2005). Investigations of the role of endogenous gut hormones have been more limited, due to the relative lack of specific receptor antagonists for use in humans. Studies using the CCK-A receptor antagonist, lorexglumide, have shown that blockade of CCK-A receptors has only a modest (Beglinger et al. 2001), or no (Lieverse et al. 1995), effect to increase energy intake, suggesting that while endogenous CCK may play a role, its individual contribution is small. GLP-1 agonists that stimulate insulin in a glucose-dependent manner, but also slow gastric emptying, and, thus, small intestinal glucose delivery and absorption, are now widely used in the management of type 2 diabetes (Nauck et al. 1997). Their use is also associated with a modest reduction in body weight in obese patients with and without type 2 diabetes, and the GLP-1 agonist, liraglutide, in a dose of 3.0 mg, has recently been approved for management of obesity in humans. However, these drugs are expensive and associated with adverse effects, particularly nausea and diarrhea (Horowitz et al. 2008), and their longer-term safety remains uncertain. Their limited effect to decrease body weight may reflect the targeting of only one mechanism.

The notion that there is not a “dominant” GI factor, but that numerous factors interact, to suppress energy intake, is not surprising, considering that under physiological conditions a range of hormones are released, and other functions, including motility, are stimulated. For example, we have shown that the potent suppressive effects of specific nutrients, when administered intraduodenally, particularly the fatty acid, lauric acid, and also the amino acid, L-tryptophan, on energy intake are associated with marked effects on the release of a number of gut hormones and modulation of upper gut motility (Felttrin et al. 2004; Steinert et al. 2014). Further evidence for the role of interrelated GI functions in the regulation of energy intake comes from our previous pooled data analysis (Seimon et al. 2010), in which the magnitude of

stimulation of both pyloric pressures and plasma CCK was identified as major independent determinants of subsequent energy intake. In contrast to this previous analysis, we were, however, unable to identify any specific parameter as a major determinant of subsequent energy intake. Instead, our data suggest that a number of factors make small contributions to determine energy intake. There was also an inconsistency; in model 2b (lipid-based studies that measured GLP-1), an increase in peak BPP by 1 mmHg was associated with a small increase, rather than a decrease, in energy intake, which is difficult to reconcile with our other findings. However, the statistical analysis used considers each factor while keeping all other variables constant, occasionally resulting in nonintuitive outcomes.

Our observations are clinically relevant, given that they provide evidence as to why any treatments that only target one pathway, for example, administration of a particular gut hormone, or its analog, are not, in the absence of aversive effects, very effective in reducing energy intake and, in the longer-term, body weight. A number of studies have investigated combination approaches (Gutzwiller et al. 2004; Neary et al. 2005; Steinert et al. 2010) and found that oral or intravenous administration of combinations of gut hormones suppressed hunger or energy intake more than individual gut hormones (reviewed in (Steinert et al. 2016)). For example, in a recent study, combined intravenous infusion of GLP-1 and PYY(3-36) reduced energy intake compared with placebo more than the sum of infusion of either hormone alone (Schmidt et al. 2014). A similar approach has been used for combinations of centrally acting drugs, for example, phentermine and topiramate, although their side-effects are considerable (Bray et al. 2016).

Our study has a number of limitations, which should be taken into account in interpreting our data. Some gut hormones that may modulate eating were either not measured (e.g., glucagon (Geary et al. 1992), pancreatic polypeptide (Jesudason et al. 2007)), or measured only in a small number of studies (e.g., ghrelin, insulin), and it remains uncertain how their inclusion in the analyses may have affected outcomes. These and other GI, or extra-GI, mechanisms, not taken into account in our studies, including plasma amino acid concentrations, warrant further investigation in prospective studies. As in our previous study (Seimon et al. 2010), PYY was not identified as an independent determinant of energy intake, but PYY was only assessed in a subset of studies, hence, the analyses may have not been sufficiently powered. That said, GLP-1 data were also available only in some studies, yet, GLP-1 concentrations were identified as an independent determinant of subsequent energy intake. The studies were performed in healthy, young men, hence, we

cannot be certain that the findings can be extended to women, overweight, obese, or elderly individuals, although this is intuitively unlikely. As gastric distension is involved in the acute regulation of energy intake, and enhances the effect of intraduodenal nutrients on GI perceptions (Feinle et al. 1997), the magnitude of effects identified in our analyses may be greater in the presence of gastric distension. Some selection of parameters to be included into the models was required. Therefore, it is possible that significant effects may have been confounded with other parameters of the same outcome (e.g., for peak vs. AUC vs. mean data of a variable). A statistical comparison of the lipid and protein cohorts to formally test for a difference in the independent determinants was not possible due to the available sample size. The comparison of results between these two subgroups is therefore qualitative only. Finally, principle components analysis, as has been used recently in a study to evaluate gastrointestinal and psychological traits associated with obesity (Acosta et al. 2015), was not deemed an appropriate analysis method in the case of our data, as our data were collected over multiple visits in each subject and there was clustering of subjects within separate studies.

In conclusion, our analysis indicates that a number of GI factors, including GI pressures, plasma CCK, and GLP-1 concentrations and some GI/appetite perception variables, are determinants of energy intake at a subsequent meal in response to intraduodenal lipid, but to a much lesser extent, protein. However, individual effects were small, therefore, the overall effect on energy intake, particularly by lipid, most likely reflects small, cumulative contributions from a combination of these, interrelated, factors. The findings have implications for the design of effective approaches for the management of obesity that target the GI tract, that is, these are unlikely to be effective if only one mechanism is targeted.

## Acknowledgments

RES' current address: DSM Nutritional Products, R&D Human Nutrition and Health, Kaiseraugst, Switzerland

## Conflict of Interest

None of the authors have any conflicts of interests to declare.

## References

Acosta, A., M. Camilleri, A. Shin, M. I. Vazquez-Roque, J. Iturrino, D. Burton, et al. 2015. Quantitative gastrointestinal and psychological traits associated with obesity and response to weight-loss therapy. *Gastroenterology* 148:537–546 e534.

- Andrews, J. M., S. Doran, G. S. Hebbard, G. Rassias, W. M. Sun, and M. Horowitz. 1998. Effect of glucose supplementation on appetite and the pyloric motor response to intraduodenal glucose and lipid. *Am. J. Physiol.* 274: G645–G652.
- Batterham, R. L., H. Heffron, S. Kapoor, J. E. Chivers, K. Chandarana, H. Herzog, et al. 2006. Critical role for peptide YY in protein-mediated satiation and body-weight regulation. *Cell Metab.* 4:223–233.
- Beglinger, C., L. Degen, D. Matzinger, M. D'Amato, and J. Drewe. 2001. Loxiglumide, a CCK-A receptor antagonist, stimulates calorie intake and hunger feelings in humans. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 280:R1149–R1154.
- Bland, J. M., and D. G. Altman. 1995. Calculating correlation coefficients with repeated observations: part 1—correlation within subjects. *BMJ* 310:446.
- Bray, G. A., G. Fruhbeck, D. H. Ryan, and J. P. Wilding. 2016. Management of obesity. *Lancet* 387:1947–56.
- Brennan, I. M., N. D. Luscombe-Marsh, R. V. Seimon, B. Otto, M. Horowitz, J. M. Wishart, et al. 2012. Effects of fat, protein, and carbohydrate and protein load on appetite, plasma cholecystokinin, peptide YY, and ghrelin, and energy intake in lean and obese men. *Am. J. Physiol. Gastrointest. Liver Physiol.* 303:G129–G140.
- Cook, C. G., J. M. Andrews, K. L. Jones, G. A. Wittert, I. M. Chapman, J. E. Morley, et al. 1997. Effects of small intestinal nutrient infusion on appetite and pyloric motility are modified by age. *Am. J. Physiol.* 273:R755–R761.
- Cukier, K., A. N. Pilichiewicz, R. Chaikomin, I. M. Brennan, J. M. Wishart, C. K. Rayner, et al. 2008. Effect of small intestinal glucose load on plasma ghrelin in healthy men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 295:R459–R462.
- Cummings, D. E., and J. Overduin. 2007. Gastrointestinal regulation of food intake. *J. Clin. Invest.* 117:13–23.
- Cummings, D. E., J. Q. Purnell, R. S. Frayo, K. Schmidova, B. E. Wisse, and D. S. Weigle. 2001. A preprandial rise in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes* 50:1714–1719.
- Degen, L., S. Oesch, M. Casanova, S. Graf, S. Ketterer, J. Drewe, et al. 2005. Effect of peptide YY3-36 on food intake in humans. *Gastroenterology* 129:1430–1436.
- Feinle, C., D. Grundy, and N. W. Read. 1997. Effects of duodenal nutrients on sensory and motor responses of the human stomach to distension. *Am. J. Physiol. Gastrointest. Liver Physiol.* 273:G721–G726.
- Feinle, C., I. M. Chapman, J. Wishart, and M. Horowitz. 2002. Plasma glucagon-like peptide-1 (GLP-1) responses to duodenal fat and glucose infusions in lean and obese men. *Peptides* 23:1491–1495.
- Feltrin, K. L., T. J. Little, J. H. Meyer, M. Horowitz, A. J. Smout, J. Wishart, et al. 2004. Effects of intraduodenal fatty

- acids on appetite, antropyloroduodenal motility, and plasma CCK and GLP-1 in humans vary with their chain length. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 287:R524–R533.
- Feltrin, K. L., M. Patterson, M. A. Ghatei, S. R. Bloom, J. H. Meyer, M. Horowitz, et al. 2006. Effect of fatty acid chain length on suppression of ghrelin and stimulation of PYY, GLP-2 and PP secretion in healthy men. *Peptides* 27:1638–1643.
- Feltrin, K. L., T. J. Little, J. H. Meyer, M. Horowitz, T. Rades, J. Wishart, et al. 2007. Effects of lauric acid on upper gut motility, plasma cholecystokinin and peptide YY, and energy intake are load, but not concentration, dependent in humans. *J. Physiol.* 581:767–777.
- Feltrin, K. L., T. J. Little, J. H. Meyer, M. Horowitz, T. Rades, J. Wishart, et al. 2008. Comparative effects of intraduodenal infusions of lauric and oleic acids on antropyloroduodenal motility, plasma cholecystokinin and peptide YY, appetite, and energy intake in healthy men. *Am. J. Clin. Nutr.* 87:1181–1187.
- Geary, N., H. R. Kissileff, F. X. Pi-Sunyer, and V. Hinton. 1992. Individual, but not simultaneous, glucagon and cholecystokinin infusions inhibit feeding in men. *Am. J. Physiol.* 262:R975–R980.
- Gentilcore, D., R. Chaikomin, K. L. Jones, A. Russo, C. Feinle-Bisset, J. M. Wishart, et al. 2006. Effects of fat on gastric emptying of and the glycemic, insulin, and incretin responses to a carbohydrate meal in type 2 diabetes. *J. Clin. Endocrinol. Metab.* 91:2062–2067.
- Gutzwiller, J. P., L. Degen, D. Matzinger, S. Prestin, and C. Beglinger. 2004. Interaction between GLP-1 and CCK-33 in inhibiting food intake and appetite in men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 287:R562–R567.
- Heddl, R., J. Dent, N. W. Read, L. A. Houghton, J. Toouli, M. Horowitz, et al. 1988a. Antropyloroduodenal motor responses to intraduodenal lipid infusion in healthy volunteers. *Am. J. Physiol.* 254:G671–G679.
- Heddl, R., J. Dent, J. Toouli, and N. W. Read. 1988b. Topography and measurement of pyloric pressure waves and tone in humans. *Am. J. Physiol.* 255:G490–G497.
- Heddl, R., P. J. Collins, J. Dent, M. Horowitz, N. W. Read, B. Chatterton, et al. 1989. Motor mechanisms associated with slowing of the gastric emptying of a solid meal by an intraduodenal lipid infusion. *J. Gastroenterol. Hepatol.* 4:437–447.
- Horowitz, M., T. Vilsboll, M. Zdravkovic, M. Hammer, and S. Madsbad. 2008. Patient-reported rating of gastrointestinal adverse effects during treatment of type 2 diabetes with the once-daily human GLP-1 analogue, liraglutide. *Diabetes Obes. Metab.* 10:593–596.
- Jesudason, D. R., M. P. Monteiro, B. M. McGowan, N. M. Neary, A. J. Park, E. Philippou, et al. 2007. Low-dose pancreatic polypeptide inhibits food intake in man. *Br. J. Nutr.* 97:426–429.
- Latner, J. D., and M. Schwartz. 1999. The effects of a high-carbohydrate, high-protein or balanced lunch upon later food intake and hunger ratings. *Appetite* 33:119–128.
- Lieverse, R. J., A. A. Masclee, J. B. Jansen, L. C. Rovati, and C. B. Lamers. 1995. Satiety effects of the type A CCK receptor antagonist loxiglumide in lean and obese women. *Biol. Psychiatry* 37:331–335.
- Little, T. J., K. L. Feltrin, M. Horowitz, A. J. Smout, T. Rades, J. H. Meyer, et al. 2005. Dose-related effects of lauric acid on antropyloroduodenal motility, gastrointestinal hormone release, appetite, and energy intake in healthy men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 289:R1090–R1098.
- Matzinger, D., L. Degen, J. Drewe, J. Meuli, R. Duebendorfer, N. Ruckstuhl, et al. 2000. The role of long chain fatty acids in regulating food intake and cholecystokinin release in humans. *Gut* 46:688–693.
- Mithieux, G. 2009. A novel function of intestinal gluconeogenesis: central signaling in glucose and energy homeostasis. *Nutrition* 25:881–884.
- Nair, N. S., I. M. Brennan, T. J. Little, D. Gentilcore, T. Hausken, K. L. Jones, et al. 2009. Reproducibility of energy intake, gastric emptying, blood glucose, plasma insulin and cholecystokinin responses in healthy young males. *Br. J. Nutr.* 101:1094–1102.
- Nauck, M. A., J. J. Holst, B. Willms, and W. Schmiegel. 1997. Glucagon-like peptide 1 (GLP-1) as a new therapeutic approach for type 2-diabetes. *Exp. Clin. Endocrinol. Diabetes* 105:187–195.
- Parker, B. A., K. Sturm, C. G. MacIntosh, C. Feinle, M. Horowitz, and I. M. Chapman. 2004. Relation between food intake and visual analogue scale ratings of appetite and other sensations in healthy older and young subjects. *Eur. J. Clin. Nutr.* 58:212–218.
- Parker, B. A., S. Doran, J. Wishart, M. Horowitz, and I. M. Chapman. 2005. Effects of small intestinal and gastric glucose administration on the suppression of plasma ghrelin concentrations in healthy older men and women. *Clin. Endocrinol. (Oxf)* 62:539–546.
- Pilichiewicz, A. N., R. Chaikomin, I. M. Brennan, J. M. Wishart, C. K. Rayner, K. L. Jones, et al. 2007a. Load-dependent effects of duodenal glucose on glycemia, gastrointestinal hormones, antropyloroduodenal motility, and energy intake in healthy men. *Am. J. Physiol. Endocrinol. Metab.* 293:E743–E753.
- Pilichiewicz, A. N., P. Papadopoulos, I. M. Brennan, T. J. Little, J. H. Meyer, J. M. Wishart, et al. 2007b. Load-dependent effects of duodenal lipid on antropyloroduodenal motility, plasma CCK and PYY, and energy intake in healthy men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 293:R2170–R2178.
- Raudenbush, S. W. B., and A. S. Bryk. 2002. Hierarchical linear models: applications and data analysis methods. SAGE Publications Inc, London.

- Riley, R. D., P. C. Lambert, and G. Abo-Zaid. 2010. Meta-analysis of individual participant data: rationale, conduct, and reporting. *BMJ* 340:c221.
- Ryan, A. T., C. Feinle-Bisset, A. Kallas, J. M. Wishart, P. M. Clifton, M. Horowitz, et al. 2012. Intraduodenal protein modulates antropyloroduodenal motility, hormone release, glycemia, appetite, and energy intake in lean men. *Am. J. Clin. Nutr.* 96:474–482.
- Ryan, A. T., N. D. Luscombe-Marsh, A. A. Saies, T. J. Little, S. Standfield, M. Horowitz, et al. 2013. Effects of intraduodenal lipid and protein on gut motility and hormone release, glycemia, appetite, and energy intake in lean men. *Am. J. Clin. Nutr.* 98:300–311.
- Santangelo, A., M. Peracchi, D. Conte, M. Fraquelli, and M. Porrini. 1998. Physical state of meal affects gastric emptying, cholecystokinin release and satiety. *Br. J. Nutr.* 80:521–527.
- Schmidt, J. B., N. T. Gregersen, S. D. Pedersen, J. L. Arentoft, C. Ritz, and T. W. Schwartz. 2014. Effects of PYY3-36 and GLP-1 on energy intake, energy expenditure, and appetite in overweight men. *Am. J. Physiol. Endocrinol. Metab.* 306: E1248–E1256.
- Seimon, R. V., K. L. Feltrin, J. H. Meyer, I. M. Brennan, J. M. Wishart, M. Horowitz, et al. 2009a. Effects of varying combinations of intraduodenal lipid and carbohydrate on antropyloroduodenal motility, hormone release, and appetite in healthy males. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 296:R912–R920.
- Seimon, R. V., T. Wooster, B. Otto, M. Golding, L. Day, T. J. Little, et al. 2009b. The droplet size of intraduodenal fat emulsions influences antropyloroduodenal motility, hormone release, and appetite in healthy males. *Am. J. Clin. Nutr.* 89:1729–1736.
- Seimon, R. V., K. Lange, T. J. Little, I. M. Brennan, A. N. Pilichiewicz, K. L. Feltrin, et al. 2010. Pooled-data analysis identifies pyloric pressures and plasma cholecystokinin concentrations as major determinants of acute energy intake in healthy, lean men. *Am. J. Clin. Nutr.* 92:61–68.
- Simmonds, M. C., J. P. Higgins, L. A. Stewart, J. F. Tierney, M. J. Clarke, and S. G. Thompson. 2005. Meta-analysis of individual patient data from randomized trials: a review of methods used in practice. *Clin. Trials* 2:209–217.
- Steinert, R. E., B. Poller, M. C. Castelli, J. Drewe, and C. Beglinger. 2010. Oral administration of glucagon-like peptide 1 or peptide YY 3-36 affects food intake in healthy male subjects. *Am. J. Clin. Nutr.* 92:810–817.
- Steinert, R. E., C. Feinle-Bisset, N. Geary, and C. Begliner. 2013. Digestive physiology of the pig symposium: secretion of gastrointestinal hormones and eating control. *J. Anim. Sci.* 91:1963–1973.
- Steinert, R. E., N. D. Luscombe-Marsh, T. J. Little, S. Standfield, B. Otto, M. Horowitz, et al. 2014. Effects of intraduodenal infusion of L-tryptophan on ad-libitum eating, antropyloroduodenal motility, glycemia, insulinemia, and gut peptide secretion in healthy men. *J. Clin. Endocrinol. Metab.* 99:3275–3284.
- Steinert, R. E., M. F. Landrock, S. S. Ullrich, S. Standfield, B. Otto, M. Horowitz, et al. 2015. Effects of intraduodenal infusion of the branched-chain amino acid, leucine, on ad libitum eating, gut motor and hormone functions, and glycemia in healthy men. *Am. J. Clin. Nutr.* 102:820–827.
- Steinert, R. E., C. Begliner, and W. Langhans. 2016. Intestinal GLP-1 and satiation: from man to rodents and back. *Int. J. Obes.* 40:198–205.
- Stunkard, A. J., and S. Messick. 1985. The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *J. Psychosom. Res.* 29:71–83.
- Tome, D. 2004. Protein, amino acids and the control of food intake. *Br. J. Nutr.* 92(Suppl 1):S27–S30.
- Tschop, M., D. L. Smiley, and M. L. Heiman. 2000. Ghrelin induces adiposity in rodents. *Nature* 407:908–913.
- Weigle, D. S., P. A. Breen, C. C. Matthys, H. S. Callahan, K. E. Meeuws, V. R. Burden, et al. 2005. A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. *Am. J. Clin. Nutr.* 82:41–48.
- Westerterp-Plantenga, M. S., V. Rolland, S. A. Wilson, and K. R. Westerterp. 1999. Satiety related to 24 h diet-induced thermogenesis during high protein/carbohydrate vs high fat diets measured in a respiration chamber. *Eur. J. Clin. Nutr.* 53:495–502.
- Westerterp-Plantenga, M. S., A. Nieuwenhuizen, D. Tome, S. Soenen, and K. R. Westerterp. 2009. Dietary protein, weight loss, and weight maintenance. *Annu. Rev. Nutr.* 29:21–41.
- Woods, S. C., M. W. Schwartz, D. G. Baskin, and R. J. Seeley. 2000. Food intake and the regulation of body weight. *Annu. Rev. Psychol.* 51:255–277.
- Wren, A. M., L. J. Seal, M. A. Cohen, A. E. Brynes, G. S. Frost, K. G. Murphy, et al. 2001. Ghrelin enhances appetite and increases food intake in humans. *J. Clin. Endocrinol. Metab.* 86:5992.