

PUBLISHED VERSION

Tiger Zhou, Emmanuelle Souzeau, Shiwani Sharma, Owen M. Siggs, Ivan Goldberg, Paul R. Healey, Stuart Graham, Alex W. Hewitt, David A. Mackey, Robert J. Casson, John Landers, Richard Mills, Jonathan Ellis, Paul Leo, Matthew A. Brown, Stuart MacGregor, Kathryn P. Burdon and Jamie E. Craig
Rare variants in optic disc area gene CARD10 enriched in primary open-angle glaucoma
Molecular Genetics & Genomic Medicine, 2016; 4(6):624-633


© 2016 The Authors. Molecular Genetics & Genomic Medicine published by Wiley Periodicals, Inc. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Originally published at:

<http://doi.org/10.1002/mgg3.248>

PERMISSIONS

<http://creativecommons.org/licenses/by/4.0/>



The image is a graphic for the Creative Commons Attribution 4.0 International License. It features a blue header with the CC and BY icons and the text "Attribution 4.0 International (CC BY 4.0)". Below the header, there is a disclaimer: "This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#)." The main content is divided into two sections: "You are free to:" and "Under the following terms:". The "You are free to:" section lists "Share" (copy and redistribute) and "Adapt" (remix, transform, and build upon) with a "Free Cultural Works APPROVED FOR" seal. The "Under the following terms:" section lists "Attribution" (give credit, link to license, indicate changes) and "No additional restrictions" (no legal terms or technological measures). A person icon is used for the Attribution term.

Attribution 4.0 International (CC BY 4.0)

This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#).

You are free to:

- Share** — copy and redistribute the material in any medium or format
- Adapt** — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the [license terms](#).

Under the following terms:

- Attribution** — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
- No additional restrictions** — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.

23 August 2017

<http://hdl.handle.net/2440/106830>

ORIGINAL ARTICLE

Rare variants in optic disc area gene *CARD10* enriched in primary open-angle glaucoma

Tiger Zhou¹, Emmanuelle Souzeau¹, Shiwani Sharma¹, Owen M. Siggs¹, Ivan Goldberg^{2,3}, Paul R. Healey², Stuart Graham², Alex W. Hewitt⁴, David A. Mackey⁵, Robert J. Casson⁶, John Landers¹, Richard Mills¹, Jonathan Ellis⁷, Paul Leo⁷, Matthew A. Brown⁷, Stuart MacGregor⁸, Kathryn P. Burdon^{1,4} & Jamie E. Craig¹

¹Department of Ophthalmology, Flinders University, Adelaide, SA, Australia

²Discipline of Ophthalmology, University of Sydney, Sydney, NSW, Australia

³Glaucoma Unit, Sydney Eye Hospital, Sydney, NSW, Australia

⁴Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS, Australia

⁵Centre for Ophthalmology and Visual Science, Lions Eye Institute, University of Western Australia, Perth, WA, Australia

⁶Discipline of Ophthalmology & Visual Sciences, University of Adelaide, Adelaide, SA, Australia

⁷Diamantina Institute, Translational Research Institute, Princess Alexandra Hospital, University of Queensland, Woolloongabba, QLD, Australia

⁸Statistical Genetics, QIMR Berghofer Medical Research Institute, Royal Brisbane Hospital, Brisbane, QLD, Australia

Keywords

CARD10, genome-wide association study, rare variants, whole exome sequencing

Correspondence

Tiger Zhou, Flinders Medical Centre, Flinders Drive, Bedford Park, SA 5042, Australia.

Tel: +61 0882046987;

Fax: +61 0882770899;

E-mail: tiger.zhou@health.sa.gov.au

Funding Information

Tiger Zhou is supported by a Lions Medical Research Foundation Scholarship in Medicine.

Stuart MacGregor is supported by an Australian Research Council Future Fellowship. Kathryn P. Burdon is a National Health and Medical Research Council (NHMRC) Senior Research Fellow. Jamie E. Craig is a National Health and Medical Research Council (NHMRC) Practitioner Fellow.

Received: 6 June 2016; Revised: 8 August

2016; Accepted: 26 August 2016

Molecular Genetics & Genomic Medicine
2016; 4(6): 624–633

doi: 10.1002/mgg3.248

Abstract**Background**

Genome-wide association studies (GWAS) have identified association of common alleles with primary open-angle glaucoma (POAG) and its quantitative endophenotypes near numerous genes. This study aims to determine whether rare pathogenic variants in these disease-associated genes contribute to POAG.

Methods

Participants fulfilled strict inclusion criteria of advanced POAG at a young age of diagnosis. Myocilin mutation carriers were excluded using direct sequencing. Whole exome sequencing was performed on 187 glaucoma cases and 103 local screened nonglaucoma controls then joint-called with exomes of 993 previously sequenced Australian controls. GWAS-associated genes were assessed for enrichment of rare predicted pathogenic variants in POAG. Significantly enriched genes were compared against Exome Aggregation Consortium (ExAC) public control.

Results

Eighty-six GWAS disease or trait-associated glaucoma genes were captured and sequenced. *CARD10* showed enrichment after Bonferroni correction for rare variants in glaucoma cases (OR = 13.2, $P = 6.94 \times 10^{-5}$) with mutations identified in 4.28% of our POAG cohort compared to 0.27% in controls. *CARD10* was significantly associated with optic disc parameters in previous GWAS. The whole GWAS gene set showed no enrichment in POAG overall (OR = 1.12, $P = 0.51$).

Conclusion

We report here an enrichment of rare predicted pathogenic coding variants within a GWAS-associated locus in POAG (*CARD10*). These findings indicate that both common and rare pathogenic coding variants in *CARD10* may contribute to POAG pathogenesis.

Introduction

Globally, glaucoma is the most common cause of irreversible and preventable blindness and the second most common cause of all blindness after cataract, accounting for 8% of blindness worldwide in 2010 (Pascolini and Mariotti 2012). The defining feature of glaucoma is an optic neuropathy characterized by optic disc cupping with corresponding visual field loss (Foster et al. 2002). With early identification and intervention, mainly with intraocular pressure (IOP)-lowering therapy, vision loss is usually preventable (Leske et al. 2003). The disease incorporates a spectrum of subtypes ranging from congenital to age-related, normal or raised IOP, open or closed iridocorneal angle and primary or secondary etiology (Kwon et al. 2009). Primary open-angle glaucoma (POAG) is the most common disease subset and the high IOP phenotype is predominant in Caucasian and African ethnicities (Tham et al. 2014). While elevated IOP is the single greatest risk factor for POAG, other associated endophenotypes include optic disc morphology and central corneal thickness (CCT) (Miglior et al. 2007; Jiang et al. 2012; Hollands et al. 2013).

POAG exhibits high heritability, with family history conferring up to a tenfold increase in disease risk, implying a genetic etiology (Wolfs et al. 1998). Several genes with high penetrance mutations have been discovered through family-based linkage studies (*MYOC* - *601652, *OPTN* - *602432, *TBK1* - *604834, *CYP11B1* - *601771), although collectively they account for <5% of total disease burden in POAG (Fingert 2011). The common disease, common variant hypothesis has been tested in POAG. Genome-wide association studies (GWAS) suggest at least part of the disease burden is attributable to common variants near or within a number of genes including *ABCA1* (+600046) (Chen et al. 2014; Gharahkhani et al. 2014; Hysi et al. 2014), *AFAP1* (*608252) (Gharahkhani et al. 2014), *ATOH7* (*609875) (Ramdas et al. 2011), *CAV1-CAV2* (*601047; *601048) (Thorleifsson et al. 2010; Wiggs et al. 2011; Hysi et al. 2014), *CDKN2B-AS1* (*613149) (Burdon et al. 2011; Nakano et al. 2012; Osman et al. 2012), *GAS7* (*603127) (Hysi et al. 2014), *GMDS* (*602884) (Gharahkhani et al. 2014), *PMM2* (*601785) (Chen et al. 2014), *SIX1-SIX6* (*601205; *606326) (Ramdas et al. 2011; Osman et al. 2012; Wiggs et al. 2012), and *TMCO1* (*614123) (Burdon et al. 2011; van Koolwijk et al. 2012; Hysi et al. 2014). Some POAG risk loci have shown significant associations with quantitative traits related to the disease such as IOP, optic disc morphology, and central corneal thickness (Lu et al. 2013; Hysi et al. 2014; Springelkamp et al. 2014). However the magnitude of odds ratios for all genome-wide associated disease polymorphisms are less than two, even

for the strongest loci including *TMCO1* (Burdon et al. 2011) and *CDKN2B-AS1* (Burdon et al. 2011; Wiggs et al. 2012), with a significant proportion of normal controls also carrying the risk alleles at associated polymorphisms (Mackey and Hewitt 2014). Furthermore, the majority of the GWAS signals are located in the noncoding regions of the genome. There is as yet no clear understanding of the mechanism of association of these noncoding polymorphisms with disease etiology (Manolio et al. 2009).

The exact contribution of rare variants to glaucoma is not known. Common tagged single-nucleotide polymorphism (SNP) analysis at the genome level, using SNP microarray technology cannot ascertain the burden of rare variants due to the poor linkage disequilibrium between common and rare variants (Siu et al. 2011). This case-control study utilized whole exome sequencing (WES) to investigate the degree of enrichment and proportion of disease burden accounted for by rare pathogenic variants in genes near common variants known to be associated with POAG and related quantitative traits from GWAS as this has not been adequately addressed in the literature.

Materials and Methods

This study adhered to the principles listed by the revised Declaration of Helsinki and the Australian National Health and Medical Research Council (NHMRC) statement of ethical conduct in research involving humans. Ethical approval was obtained from the Southern Adelaide and Flinders University Clinical Research Ethics Committee. All peripheral blood samples were collected for genomic DNA extraction as a part of the Australian and New Zealand Registry of Advanced Glaucoma (ANZRAG) as previously described (Souzeau et al. 2012).

Participants

Unrelated Caucasian participants with advanced glaucoma and nonglaucoma Caucasian controls were included for the study. All DNA samples were extracted from peripheral blood samples, using the QIAamp[®] DNA blood kit (Qiagen, Hilden, Germany) following the manufacturer's protocol. Our case cohort inclusion criteria selected only participants with the most severe disease and the youngest age (mean = 44.4 years, SD = 10.4 years) of onset within the ANZRAG database. For this study, participants were included if there was glaucomatous visual field loss involving at least two of the four central fixation squares with a pattern standard deviation of <0.5% on a reliable Humphrey 24-2 field (Carl Zeiss, Dublin, CA), or a mean deviation of worse than -15 dB in the worst affected eye. For participants without formal visual field testing, their best-corrected visual acuity had to be worse than 20/200 due to

glaucomatous damage. Furthermore glaucomatous damage had to also be present in the less affected eye as measured by a reliable Humphrey 24-2 field, with corresponding neuroretinal rim thinning. With the ANZRAG database participants with the youngest age of diagnosis were selected for this study. Efforts were made to ensure a comparable split of participants with high and normal IOP. High-tension glaucoma (HTG) was defined as having a maximum recorded IOP of >21 mmHg. Individuals with known *MYOC* mutations were excluded from this study as the genetic cause for their POAG has already been identified.

For the local control cohort, all participants were examined to exclude glaucoma or glaucoma-related phenotypes. Furthermore, in silico analysis included a larger unexamined control cohort from the Australian Osteoporosis Genetics Consortium (Estrada et al. 2012) (AOGC). These controls were female participants with high or low bone mass who were otherwise healthy by self-report.

Data acquisition

Exon capture and enrichment was performed on all glaucoma cases and local controls with the SureSelect Human All Exon V4 enrichment kit (Agilent, Santa Clara, CA, USA) as per manufacturer's protocol and enriched DNA sequenced on a HiSeq2000 (Illumina, San Diego, CA, USA) with paired end 100 bp reads (Macrogen Next Generation Sequencing Services). Local controls served as both technical and phenotypic controls. Raw experimental data were cleaned and joint-called with AOGC exome data that were captured with Nimblegen Human Exome Capture V2 (Roche, Basel, Switzerland) and sequenced on the HiSeq2000 (Illumina, San Diego, CA, USA) (University of Queensland Centre for Clinical Genomics). FASTQ files were aligned to human genome build hg19 with novoalign (version 3.02.08). Duplicate reads were marked with Picard MarkDuplicates (version 1.124). Local indel realignment and base quality recalibration were performed with GATK (McKenna et al. 2010) (version 3.2-2). Variant calling of single-nucleotide variants and small indels was performed with the UnifiedGenotyper module in GATK and variant quality scores were recalibrated according to the GATK "Best Practices Guidelines" (DePristo et al. 2011; Van der Auwera et al. 2013). Annotation of calls was performed using ANNOVAR (Wang et al. 2010) software using refGene, SIFT, PolyPhen2 HVAR, Exome Sequencing Project (ESP), 1000 Genomes and Exome Aggregation Consortium (ExAC) databases.

Data analysis

Using in-house UNIX scripts, only protein coding exonic and splicing site variants were selected for analysis. Genes

were included in the list if they were reported by any published GWAS or meta-analysis paper to be the closest to SNPs associated with POAG or disease endophenotype at $P < 5 \times 10^{-8}$ in any ethnicity. Quality control threshold was set at a Genotype Quality (GQ) score of 20. Focusing on the impact of rare mutations, all variants with known minor allele frequency (MAF) of >1% in dbSNP, 1000 genomes and ExAC were excluded. Pathogenicity filtering removed all synonymous variants and missense variants considered tolerated or only possibly damaging by both SIFT and PolyPhen-2 HVAR, and retained stop site, frameshift and predicted deleterious missense mutations. Raw VCF data from the Exome Aggregation Consortium were annotated with the ANNOVAR pipeline followed by variant selection using the same in-house scripts to target only rare deleterious mutations in the non-Finnish European subgroup to act as a larger in silico population control. Mutation loads per gene were calculated for the glaucoma cases, local control, AOGC control and ExAC control cohorts by summing the minor allele counts of all qualifying variants in the same gene and dividing by the average number of captured alleles for those variants, thereby adjusting for capture rate. Odds ratios were generated between cases and each control cohort separately, and Fisher's exact test used to calculate *P*-values, and Bonferroni correction applied for multiple testing of all analyzed genes ($n = 86$, threshold $P = 5.81 \times 10^{-4}$). Five sets of randomly chosen independent groups of 86 WES-captured genes were selected using a random number generator and analyzed using the same protocol to act as control gene sets. This step further strengthens the positive results by examining the likelihood of false-positive findings.

Validation of variants

All variants within significantly associated genes were independently validated by capillary sequencing (primers listed in Table S1). PCRs were conducted on carrier samples using 40 ng of DNA, 10 μ mol of each corresponding forward and reverse primers, 0.5 units of HotStarTaq DNA polymerase (Qiagen, Hilden, Germany), 2 μ mol of dNTP and PCR buffer in a 20 μ L volume per reaction. Thirty cycles were performed at an annealing temperature of 62°C. Five μ L of each PCR product was incubated with 2 μ L of Shrimp Alkaline Phosphatase (Affymetrix, Santa Clara, CA, USA) and 0.5 μ L of *E. coli* exonuclease I (New England BioLabs, Ipswich, MA, USA) at 37°C for 60 min. Sequencing was carried out on the fluorescence-based capillary electrophoresis system 3130xl Genetic Analyzer (Life Technologies, Carlsbad, CA, USA) using BigDye Terminator V3.1 (Life Technologies, Carlsbad, CA, USA) according to manufacturer's protocol.

Results

A total of 187 unrelated participants with advanced POAG and 103 screened nonglaucoma controls were sequenced with whole exome capture and enrichment. Combined with the large unscreened cohort of AOGC controls ($n = 993$), the total number of controls was 1096. More participants had high-tension glaucoma (HTG, $n = 122$) than normal-tension glaucoma (NTG, $n = 65$). To date, 101 genes have been reported by GWAS studies as being near SNPs that are statistically associated with POAG, central corneal thickness, IOP or optic disc morphology. Of these 101 genes, 86 had qualifying (MAF < 0.01 and predicted pathogenic) variants captured in glaucoma and control cohorts in this study (Table S2).

The sequencing strategy utilized for cases and local controls in this study yielded good read quality, coverage, and depth. Mean percentage of mappable reads was high at 99.4% with an average total on-target reads per sample of 4.12×10^9 and an average depth of 73 reads per target base. Average coverage at $>10\times$ depth was 97.9% of all targeted exonic regions. The average numbers of coding SNPs and indels per participant was 19,605 and 465, respectively. AOGC controls had an average depth of 24 reads per target base and $10\times$ coverage of 75.1%. After filtering for rare predicted pathogenic variants only, there was an average of 159 qualifying variants per participant. A total of 1159 variants were within the 86 included genes in case, local or AOGC control cohorts. There was no significant enrichment (OR = 1.12, $P = 0.51$) in the total carrier rate of qualifying variants in this gene set in glaucoma cases (64.2%, 120 individuals) compared with local and AOGC controls (61.5%, 648 individuals) although there was a trend towards increased mutation load in POAG cases (Table S3).

Four genes (*CARD10*, *CWC27*, *RERE*, and *USP37*) were nominally enriched (uncorrected $P < 0.05$) in the glaucoma cohort (Table 1). Only *CARD10* (caspase recruitment domain containing protein 10) remained

Table 1. Genes at genome-wide significant associated loci that reached nominal significance for enrichment of rare variants in the glaucoma cohort. Mutational load represents the total allele frequency burden of qualifying variants in each gene. Odds ratios are adjusted for capture quality described in the Methods section. P -values are corrected using Bonferroni multiple testing correction.

Gene	POAG load (%)	Control load (%)	Odds ratio	P -value	Corrected P -value
<i>CARD10</i>	2.14	0.14	13.19	6.94×10^{-5}	5.97×10^{-3}
<i>CWC27</i>	1.34	0.14	6.89	9.47×10^{-3}	0.81
<i>RERE</i>	1.60	0.23	5.49	0.0114	0.98
<i>USP37</i>	1.87	0.18	7.32	5.11×10^{-3}	0.44

statistically significant after Bonferroni correction for 86 tested genes ($P = 6.94 \times 10^{-5}$, corrected $P = 5.97 \times 10^{-3}$) with an odds ratio of 13.2 (3.5–50.2). Eight out of 187 POAG cases (4.28%) carried a rare predicted pathogenic variant in *CARD10* confirmed by Sanger sequencing (Figure S1). Comparatively, qualifying variants in *CARD10* were carried by 0.27% (3/1096) of the joint-called controls. A total of five nonsynonymous variants in this gene were identified in the POAG cohort with all but one absent in the controls (Table 2). These variants were predicted to be pathogenic by SIFT or PolyPhen2 as per the filtering criteria. Additionally, the variants were located in highly conserved regions, all having GERP (Genomic Evolutionary Rate Profiling) scores of >2 (mean GERP score = 4.8). When tested against qualifying variants in the unscreened public domain ExAC non-Finnish European cohort ($n = 33370$) filtered with our pipeline, *CARD10* remained significantly enriched in the glaucoma cohort, albeit with a lower odds ratio (OR = 3.3, $P = 3.9 \times 10^{-3}$). *CARD10* variant carriers had younger age at diagnosis (34.9 vs. 44.8 years, $P = 0.36$) and higher IOP (27.4 vs. 24.8, $P = 0.11$) than non-*CARD10* cases, although the difference was not statistically significant due to the small number of *CARD10* carriers. All eight *CARD10* qualifying variants were successfully validated in the carrier cases using capillary sequencing (Figure S1) and submitted to ClinVar database (<http://www.ncbi.nlm.nih.gov/clinvar/>) with accessions SCV000266585–SCV000266589. Subanalyses of the HTG and NTG cohorts with local and AOGC controls showed significant association of *CARD10* variants with HTG subgroup (OR = 15.2, corrected $P = 0.01$). Five control gene sets of 86 randomly selected genes were examined and none contained any gene that was significantly associated with POAG after Bonferroni correction (Tables S4–S8).

Discussion

Genome-wide association studies have been examined for common variants linked to POAG, NTG and various associated endophenotypes; IOP, vertical cup-to-disc ratio (VCDR), neuroretinal rim area, optic disc area and central corneal thickness. Poor linkage disequilibrium exists between common and rare variants; this has been demonstrated experimentally (Siu et al. 2011). For this reason, previous GWAS designs have reduced power to detect signals at single-nucleotide variants (SNVs) with MAF of $<1\%$. This study employed WES to patch the “gap” left by GWAS and to explore whether rare variants (MAF $<1\%$) within 86 putative POAG risk loci were also enriched within a severe POAG cohort. Although the majority of rare variants from the 86 POAG risk loci do

Table 2. Qualifying *CARD10* (NM_014550.3) variants captured in glaucoma cohort and controls. SIFT score <0.05 is predicted damaging. PolyPhen2 HVAR score >0.909 is predicted damaging.

cDNA	Residue	SIFT	PolyPhen2	POAG cases	POAG freq	Control freq	ExAC freq
c.635G>A	p.Arg212His	0.10	0.987	1	2.7×10^{-3}	0	0
c.983C>T	p.Ala328Val	0.02	0.944	3	8.0×10^{-3}	9.1×10^{-4}	3.6×10^{-3}
c.1024G>A	p.Val342Met	0.04	0.025	1	2.7×10^{-3}	0	2×10^{-4}
c.1210C>T	p.Arg404Trp	0.01	0.764	1	2.7×10^{-3}	0	1.5×10^{-5}
c.2485C>T	p.Arg829Trp	0.02	0.01	2	5.3×10^{-3}	0	4.0×10^{-4}
c.3081delC	p.Pro1027fs			0	0	4.6×10^{-4}	0

not show significant enrichment in cases, the plausibility that GWAS candidates may contribute to disease via a common disease rare variant hypothesis has been demonstrated by our study.

A previous candidate gene study of *SIX6* reported an enrichment of rare variants in POAG in that gene (Carnes et al. 2014). The most prevalent variant, with a carrier frequency of 1.6% in cases, identified by Carnes et al. (2014) (rs146737847:G>A) was found in 1.6% of our POAG cases, but also in 1.9% of local screened controls and 1.6% of AOGC controls with an overall OR of 0.98. The other three rare variants found in that study at a carrier frequency of 0.4% were not detected in our cases or controls. In summary, our study found no enrichment of rare variants in *SIX6* in our POAG cases versus controls.

Genome-wide association of *CARD10*

SNPs near *CARD10* were first found to be significantly associated with optic disc area on GWAS in Singaporean Asians, a result replicated in a Dutch Caucasian cohort (Khor et al. 2011). Optic disc area is relevant in POAG as a large optic disc is correlated with increased susceptibility to glaucoma in Caucasian populations (Healey and Mitchell 1999). Meta-analysis of optic disc morphology further implicated common variants in *CARD10* in VCDR but not POAG after adjustment for optic disc area (Springelkamp et al. 2014). This result has since been replicated in a separate case-control study in an Indian population (Philomenadin et al. 2015). Cup-to-disc ratio is fundamental in POAG as it is often used as a diagnostic criterion due to its strong positive correlation with disease incidence (Miglior et al. 2007; Hollands et al. 2013). The SNPs from the original GWAS linking *CARD10* to optic disc area were situated between 3262 bp and 7204 bp upstream of *CARD10* (Khor et al. 2011). These SNPs are in strong linkage disequilibrium with all common SNPs up to and including rs9610775, which is a missense coding variant, p.R289Q in the *CARD10* gene with a population MAF of 13% (Fig. 1). The signal identified in the optic disc area GWAS may possibly relate to *CARD10* coding mutations. The SNPs found to be

significantly associated with VCDR on meta-analysis were 260 kb upstream of *CARD10* with no linkage disequilibrium to the gene (Springelkamp et al. 2014). More likely, these variants are involved in gene expression and regulation. Therefore, the published GWAS data suggests that both coding variants in *CARD10* and its expression level may contribute to optic disc pathology.

GWAS was able to highlight the association between regulatory and common coding SNPs in *CARD10* with crucial optic disc parameters for the development of POAG (Khor et al. 2011; Springelkamp et al. 2014; Philomenadin et al. 2015). This study complements the GWAS findings by implicating rare coding *CARD10* variants that likely disrupt gene function, and are associated with increased POAG risk within our cohort of advanced glaucoma participants. Both common and rare variants concurrently contribute to disease risk via differing mechanisms affecting the same gene - *CARD10*. Although more HTG participants harbored rare *CARD10* variants than NTG participants, the presence of NTG *CARD10* variant carriers suggests that IOP elevation is not required for the potential of *CARD10* to cause glaucoma. No definitive inference can be made from this discrepancy due to the small sample size of the subcohorts. A plausible hypothesis would be that the increased susceptibility to retinal ganglion cell apoptosis in the *CARD10* mutation carrying individuals is compounded by elevated IOP to cause glaucoma at an earlier age. The participant inclusion criteria of this study selected exclusively for advanced glaucoma phenotype and young age of diagnosis resulting in an enrichment of *CARD10* variant carriers that also maintained elevated IOP.

CARD10 function

Discovered in 2001, *CARD10* functions as a signaling protein in the regulation of the NF κ B (nuclear factor kappa B) pathway via activation of membrane-bound G protein-coupled receptors (McAllister-Lucas et al. 2001; Wang et al. 2001). This cascade is initiated through interactions between the N-terminal CARD domains of *CARD10*, *BCL10* and *MALT1*, which forms a CBM complex




Figure 1. Linkage disequilibrium between GWAS SNPs and common SNPs in *CARD10* from HapMap CEU data showing D' values. GWAS SNPs are underlined. *CARD10* gene location is represented by rectangle at the top. The left-most SNP rs9610775 (p.Arg289Gln) represents the boundary of overlap between the GWAS significant LD block and the *CARD10* gene.

(Scudiero et al. 2014). Overexpression of *CARD10* has been demonstrated in multiple human neoplasias including bladder (Man et al. 2014), breast (Zhao et al. 2013), colon (Miao et al. 2012), lung (Li et al. 2012), ovarian (Xie et al. 2014), pancreatic (Du et al. 2014) and renal cancers (Wu et al. 2013). In all cases, the overexpression of *CARD10* is associated with increased cell survival, proliferation and therefore poor prognosis of cancer. The underlying physiology appears to involve enhanced cell cycling via NF κ B facilitated up regulation of cyclin D1, cyclin E, Bcl-2 and phosphorylated I κ B (Miao et al. 2012; Zhao et al. 2013; Xie et al. 2014). Other documented consequences of *CARD10* upregulation include leukocyte activation (Cowan et al. 2014) and inhibition of angiogenesis (Rau et al. 2014). Homozygous knockout of *CARD10* in mice causes nonviability through neural tube defects (Grabiner et al. 2007). Grabiner et al. (2007) further demonstrated that *CARD10* may be required in neural crest cell survival through G protein-coupled receptor induction of NF κ B activation.

Primary open-angle glaucoma is a disease of enhanced retinal ganglion cell apoptosis (Foster et al. 2002). The NF κ B pathway is profoundly involved in regulation of cellular apoptosis. Traditionally, it was only thought to be a promoter of cell survival and proliferation via downstream transcription of anti-apoptotic proteins. As such, overexpression of NF κ B leads to enhanced growth of cancerous cells (Escarcega et al. 2007). However more recent evidence in the central nervous system suggests NF κ B may also play a proapoptotic role depending on the nature of the noxious stimuli (Kaltschmidt et al. 2005). In certain situations such as neuronal ischemia and Parkinson's disease, upregulation of NF κ B led to neuronal death via p53 signaling. Evidence in tumor cells indicates that *CARD10* promotes the antiapoptotic effect of NF κ B signaling. Therefore loss of function or down-regulation of *CARD10* leads to increased apoptosis, especially in mice neural crest cells (Grabiner et al. 2007). Conversely, proapoptotic activity of NF κ B signalling has been recognized in the rat retina. NMDA (N-methyl-d-

aspartate)-induced expression of NF κ B p65 led to retinal ganglion cell death which was ameliorated with knock-down of p65 by antisense oligonucleotides (Kitaoka et al. 2007). Retinal ganglion cell damage from NF κ B is likely due to glial cell activation and interleukin (IL)-1 β secretion (Kitaoka et al. 2007). Given the intimate and complex relationship between *CARD10*, NF κ B and apoptosis, pathogenic coding mutations in *CARD10* are likely to affect apoptosis signaling. The coding mutations identified within the glaucoma cohort in this study have not been previously characterized, however they may augment retinal ganglion cell death via up or down-regulation of NF κ B. This distinction is important to ascertain in order to translate these findings toward therapeutic strategies in glaucoma management.

Strengths and limitations

A strength of this study is the enrichment of the disease cohort with well-selected extreme disease phenotypes, and it is the largest study to examine whole exomes of a well matched case-control cohort of this nature in the glaucoma field. ANZRAG is the largest collection advanced glaucoma patients worldwide. Our controls are hypernormal and have been individually examined to ensure the absence of glaucoma. Finally, all significantly enriched variants were confirmed via capillary sequencing thereby eliminating the possibility of false-positive calls.

This study has some limitations which were largely technical. Whole exome sequencing has enabled the rapid examination of large numbers of genes at a reasonable cost. Its advantage over microarray technology for genetic studies lies in its ability to capture rare coding variants. However this also introduces the challenge of variable capture. AOGC samples were sequenced on a different platform with a different capture probe set to the case cohort. Joint-calling of the two datasets was used to limit the amount of artifact generated *in silico*. However, capture and sequencing platform-specific differences remained. Some difference in capture rate at specific locations was evident and translated to incomplete capture of some variants in the AOGC cohort. Adjustment for the capture rate was performed in the calculation of odds ratios and *P*-values. ExAC samples are more heterogeneous in data acquisition compared with this study cohort. Inability to joint-call our data with ExAC may have overestimated the actual mutation burden in this public domain control cohort. Hence the odds ratio between case and controls falls considerably when ExAC data is included and may represent an underestimation. Alternatively, incomplete coverage of AOGC controls may have contributed to an overestimation of the discovery odds ratio. Nevertheless, a statistically significant difference remained between POAG

and controls in *CARD10* mutational burden even with comparison with ExAC controls.

Conclusion

Overall, the majority of genes near common variants associated with glaucoma and its endophenotypes found in GWAS did not show enrichment of rare variants in POAG. Rare predicted pathogenic variants in *CARD10*, a gene located near SNPs associated with optic disc area and VCDR in genome-wide association studies, were enriched in a cohort of advanced POAG. Further research is required to elucidate the exact influence of mutations in *CARD10* on NF κ B signaling, apoptosis, and the development of POAG.

Acknowledgments

Tiger Zhou is supported by a Lions Medical Research Foundation Scholarship in Medicine. Stuart MacGregor is supported by an Australian Research Council Future Fellowship. Kathryn P. Burdon is a National Health and Medical Research Council (NHMRC) Senior Research Fellow. Jamie E. Craig is a National Health and Medical Research Council (NHMRC) Practitioner Fellow.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Axenovich, T., I. Zorkoltseva, N. Belonogova, L. M. van Koolwijk, P. Borodin, A. Kirichenko, et al. 2011. Linkage and association analyses of glaucoma related traits in a large pedigree from a Dutch genetically isolated population. *J. Med. Genet.* 48:802–809.
- Bailey, J. N., S. J. Loomis, J. H. Kang, R. R. Allingham, P. Gharahkhani, C. C. Khor, et al. 2016. Genome-wide association analysis identifies TXNRD2, ATXN2 and FOXC1 as susceptibility loci for primary open-angle glaucoma. *Nat. Genet.* 48:189–194.
- Burdon, K. P., S. Macgregor, A. W. Hewitt, S. Sharma, G. Chidlow, R. A. Mills, et al. 2011. Genome-wide association study identifies susceptibility loci for open angle glaucoma at TMCO1 and CDKN2B-AS1. *Nat. Genet.* 43:574–578.
- Carnes, M. U., Y. P. Liu, R. R. Allingham, B. T. Whigham, S. Havens, M. E. Garrett, et al. 2014. Discovery and functional annotation of SIX6 variants in primary open-angle glaucoma. *PLoS Genet.* 10:e1004372.
- Chen, Y., Y. Lin, E. N. Vithana, L. Jia, X. Zuo, T. Y. Wong, et al. 2014. Common variants near ABCA1 and in PMM2 are associated with primary open-angle glaucoma. *Nat. Genet.* 46:1115–1119.

- Cowan, C., C. K. Muraleedharan, J. J. 3rd O'Donnell, P. K. Singh, H. Lum, A. Kumar, et al. 2014. MicroRNA-146 inhibits thrombin-induced NF-kappaB activation and subsequent inflammatory responses in human retinal endothelial cells. *Invest. Ophthalmol. Vis. Sci.* 55:4944–4951.
- DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire, C. Hartl, et al. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* 43:491–498.
- Du, S., L. Jia, Y. Zhang, L. Fang, X. Zhang, and Y. Fan. 2014. CARMA3 is upregulated in human pancreatic carcinoma, and its depletion inhibits tumor proliferation, migration, and invasion. *Tumour Biol.* 35:5965–5970.
- Escarcega, R. O., S. Fuentes-Alexandro, M. Garcia-Carrasco, A. Gatica, and A. Zamora. 2007. The transcription factor nuclear factor-kappa B and cancer. *Clin. Oncol. (R. Coll. Radiol.)* 19:154–161.
- Estrada, K., U. Styrkarsdottir, E. Evangelou, Y. H. Hsu, E. L. Duncan, E. E. Ntzani, et al. 2012. Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture. *Nat. Genet.* 44:491–501.
- Fingert, J. H. 2011. Primary open-angle glaucoma genes. *Eye (London)* 25:587–595.
- Foster, P. J., R. Buhrmann, H. A. Quigley, and G. J. Johnson. 2002. The definition and classification of glaucoma in prevalence surveys. *Br. J. Ophthalmol.* 86:238–242.
- Gao, X., W. J. Gauderman, Y. Liu, P. Marjoram, M. Torres, T. Haritunians, et al. 2013. A genome-wide association study of central corneal thickness in Latinos. *Invest. Ophthalmol. Vis. Sci.* 54:2435–2443.
- Gharahkhani, P., K. P. Burdon, R. Fogarty, S. Sharma, A. W. Hewitt, S. Martin, et al. 2014. Common variants near ABCA1, AFAP1 and GMDS confer risk of primary open-angle glaucoma. *Nat. Genet.* 46:1120–1125.
- Grabiner, B. C., M. Blonska, P. C. Lin, Y. You, D. Wang, J. Sun, et al. 2007. CARMA3 deficiency abrogates G protein-coupled receptor-induced NF- κ B activation. *Genes Dev.* 21:984–996.
- Healey, P. R., and P. Mitchell. 1999. Optic disk size in open-angle glaucoma: the Blue Mountains Eye Study. *Am. J. Ophthalmol.* 128:515–517.
- Hoehn, R., T. Zeller, V. J. Verhoeven, F. Grus, M. Adler, R. C. Wolfs, et al. 2012. Population-based meta-analysis in Caucasians confirms association with COL5A1 and ZNF469 but not COL8A2 with central corneal thickness. *Hum. Genet.* 131:1783–1793.
- Hollands, H., D. Johnson, S. Hollands, D. L. Simel, D. Jinapriya, and S. Sharma. 2013. Do findings on routine examination identify patients at risk for primary open-angle glaucoma? The rational clinical examination systematic review. *JAMA* 309:2035–2042.
- Hysi, P. G., C. Y. Cheng, H. Springelkamp, S. Macgregor, J. N. Bailey, R. Wojciechowski, et al. 2014. Genome-wide analysis of multi-ancestry cohorts identifies new loci influencing intraocular pressure and susceptibility to glaucoma. *Nat. Genet.* 46:1126–1130.
- Jiang, X., R. Varma, S. Wu, M. Torres, S. P. Azen, B. A. Francis, et al. 2012. Baseline risk factors that predict the development of open-angle glaucoma in a population: the Los Angeles Latino Eye Study. *Ophthalmology* 119:2245–2253.
- Kaltschmidt, B., D. Widera, and C. Kaltschmidt. 2005. Signaling via NF-kappaB in the nervous system. *Biochim. Biophys. Acta* 1745:287–299.
- Khor, C. C., W. D. Ramdas, E. N. Vithana, B. K. Cornes, X. Sim, W. T. Tay, et al. 2011. Genome-wide association studies in Asians confirm the involvement of ATOH7 and TGFBR3, and further identify CARD10 as a novel locus influencing optic disc area. *Hum. Mol. Genet.* 20:1864–1872.
- Kitaoka, Y., Y. Munemasa, T. Nakazawa, and S. Ueno. 2007. NMDA-induced interleukin-1beta expression is mediated by nuclear factor-kappa B p65 in the retina. *Brain Res.* 1142:247–255.
- van Koolwijk, L. M., W. D. Ramdas, M. K. Ikram, N. M. Jansonius, F. Pasutto, P. G. Hysi, et al. 2012. Common genetic determinants of intraocular pressure and primary open-angle glaucoma. *PLoS Genet.* 8:e1002611.
- Kwon, Y. H., J. H. Fingert, M. H. Kuehn, and W. L. Alward. 2009. Primary open-angle glaucoma. *N. Engl. J. Med.* 360:1113–1124.
- Leske, M. C., A. Heijl, M. Hussein, B. Bengtsson, L. Hyman, and E. Komaroff. 2003. Factors for glaucoma progression and the effect of treatment: the early manifest glaucoma trial. *Arch. Ophthalmol.* 121:48–56.
- Li, Z., L. Qu, Q. Dong, B. Huang, H. Li, Z. Tang, et al. 2012. Overexpression of CARMA3 in non-small-cell lung cancer is linked for tumor progression. *PLoS ONE* 7:e36903.
- Li, Z., R. R. Allingham, M. Nakano, L. Jia, Y. Chen, Y. Ikeda, et al. 2015. A common variant near TGFBR3 is associated with primary open angle glaucoma. *Hum. Mol. Genet.* 24:3880–3892.
- Lu, Y., V. Vitart, K. P. Burdon, C. C. Khor, Y. Bykhovskaya, A. Mirshahi, et al. 2013. Genome-wide association analyses identify multiple loci associated with central corneal thickness and keratoconus. *Nat. Genet.* 45:155–163.
- Macgregor, S., A. W. Hewitt, P. G. Hysi, J. B. Ruddle, S. E. Medland, A. K. Henders, et al. 2010. Genome-wide association identifies ATOH7 as a major gene determining human optic disc size. *Hum. Mol. Genet.* 19:2716–2724.
- Mackey, D. A., and A. W. Hewitt. 2014. Genome-wide association study success in ophthalmology. *Curr. Opin. Ophthalmol.* 25:386–393.
- Man, X., J. He, C. Kong, Y. Zhu, and Z. Zhang. 2014. Clinical significance and biological roles of CARMA3 in human bladder carcinoma. *Tumour Biol.* 35:4131–4136.

- Manolio, T. A., F. S. Collins, N. J. Cox, D. B. Goldstein, L. A. Hindorff, D. J. Hunter, et al. 2009. Finding the missing heritability of complex diseases. *Nature* 461:747–753.
- McAllister-Lucas, L. M., N. Inohara, P. C. Lucas, J. Ruland, A. Benito, Q. Li, et al. 2001. Bim1, a MAGUK family member linking protein kinase C activation to Bcl10-mediated NF-kappaB induction. *J. Biol. Chem.* 276:30589–30597.
- McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernysky, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.
- Meguro, A., H. Inoko, M. Ota, N. Mizuki, and S. Bahram. 2010. Genome-wide association study of normal tension glaucoma: common variants in SRBD1 and ELOVL5 contribute to disease susceptibility. *Ophthalmology* 117:1331–1338. e5.
- Miao, Z., T. Zhao, Z. Wang, Y. Xu, Y. Song, J. Wu, et al. 2012. CARMA3 is overexpressed in colon cancer and regulates NF-kappaB activity and cyclin D1 expression. *Biochem. Biophys. Res. Commun.* 425:781–787.
- Miglior, S., N. Pfeiffer, V. Torri, T. Zeyen, J. Cunha-Vaz, and I. Adamsons. 2007. Predictive factors for open-angle glaucoma among patients with ocular hypertension in the European Glaucoma Prevention Study. *Ophthalmology* 114:3–9.
- Nag, A., C. Venturini, K. S. Small, T. L. Young, A. C. Viswanathan, D. A. Mackey, et al. 2014. A genome-wide association study of intra-ocular pressure suggests a novel association in the gene FAM125B in the TwinsUK cohort. *Hum. Mol. Genet.* 23:3343–3348.
- Nakano, M., Y. Ikeda, Y. Tokuda, M. Fuwa, N. Omi, M. Ueno, et al. 2012. Common variants in CDKN2B-AS1 associated with optic-nerve vulnerability of glaucoma identified by genome-wide association studies in Japanese. *PLoS ONE* 7:e33389.
- Osman, W., S. K. Low, A. Takahashi, M. Kubo, and Y. Nakamura. 2012. A genome-wide association study in the Japanese population confirms 9p21 and 14q23 as susceptibility loci for primary open angle glaucoma. *Hum. Mol. Genet.* 21:2836–2842.
- Ozel, A. B., S. E. Moroi, D. M. Reed, M. Nika, C. M. Schmidt, S. Akbari, et al. 2014. Genome-wide association study and meta-analysis of intraocular pressure. *Hum. Genet.* 133:41–57.
- Pascalini, D., and S. P. Mariotti. 2012. Global estimates of visual impairment: 2010. *Br. J. Ophthalmol.* 96:614–618.
- Philomenadin, F. S., R. Asokan, N. Viswanathan, R. George, V. Lingam, and S. Sarangapani. 2015. Genetic association of SNPs near ATOH7, CARD10, CDKN2B, CDC7 and SIX1/SIX6 with the endophenotypes of primary open angle glaucoma in Indian population. *PLoS ONE* 10:e0119703.
- Ramdas, W. D., L. M. van Koolwijk, M. K. Ikram, N. M. Jansonius, P. T. de Jong, A. A. Bergen, et al. 2010. A genome-wide association study of optic disc parameters. *PLoS Genet.* 6:e1000978.
- Ramdas, W. D., L. M. van Koolwijk, H. G. Lemij, F. Pasutto, A. J. Cree, G. Thorleifsson, et al. 2011. Common genetic variants associated with open-angle glaucoma. *Hum. Mol. Genet.* 20:2464–2471.
- Rau, C. S., J. C. Yang, Y. C. Chen, C. J. Wu, T. H. Lu, S. L. Tzeng, et al. 2014. Lipopolysaccharide-induced microRNA-146a targets CARD10 and regulates angiogenesis in human umbilical vein endothelial cells. *Toxicol. Sci.* 140:315–326.
- Scudiero, I., P. Vito, and R. Stilo. 2014. The three CARMA sisters: so different, so similar: a portrait of the three CARMA proteins and their involvement in human disorders. *J. Cell. Physiol.* 229:990–997.
- Siu, H., Y. Zhu, L. Jin, and M. Xiong. 2011. Implication of next-generation sequencing on association studies. *BMC Genom.* 12:322.
- Souzeau, E., I. Goldberg, P. R. Healey, R. A. Mills, J. Landers, S. L. Graham, et al. 2012. Australian and New Zealand registry of advanced glaucoma: methodology and recruitment. *Clin. Exp. Ophthalmol.* 40:569–575.
- Springelkamp, H., R. Hohn, A. Mishra, P. G. Hysi, C. C. Khor, S. J. Loomis, et al. 2014. Meta-analysis of genome-wide association studies identifies novel loci that influence cupping and the glaucomatous process. *Nat. Commun.* 5:4883.
- Springelkamp, H., A. Mishra, P. G. Hysi, P. Gharahkhani, R. Hohn, C. C. Khor, et al. 2015. Meta-analysis of genome-wide association studies identifies novel loci associated with optic disc morphology. *Genet. Epidemiol.* 39:207–216.
- Strange, A., C. Bellenguez, X. Sim, R. Luben, P. G. Hysi, W. D. Ramdas, et al. 2013. Genome-wide association study of intraocular pressure identifies the GLCC11/ICA1 region as a glaucoma susceptibility locus. *Hum. Mol. Genet.* 22:4653–4660.
- Takamoto, M., T. Kaburaki, A. Mabuchi, M. Araie, S. Amano, M. Aihara, et al. 2012. Common variants on chromosome 9p21 are associated with normal tension glaucoma. *PLoS ONE* 7:e40107.
- Tham, Y. C., X. Li, T. Y. Wong, H. A. Quigley, T. Aung, and C. Y. Cheng. 2014. Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. *Ophthalmology* 121:2081–2090.
- Thorleifsson, G., G. B. Walters, A. W. Hewitt, G. Masson, A. Helgason, A. DeWan, et al. 2010. Common variants near CAV1 and CAV2 are associated with primary open-angle glaucoma. *Nat. Genet.* 42:906–909.
- Ulmer, M., J. Li, B. L. Yaspan, A. B. Ozel, J. E. Richards, S. E. Moroi, et al. 2012. Genome-wide analysis of central corneal thickness in primary open-angle glaucoma cases in the NEIGHBOR and GLAUGEN consortia. *Invest. Ophthalmol. Vis. Sci.* 53:4468–4474.
- Van der Auwera, G. A., M. O. Carneiro, C. Hartl, R. Poplin, G. Del Angel, A. Levy-Moonshine, et al. 2013. From FastQ

- data to high confidence variant calls: the genome analysis toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* 11:11 10 1–11 10 33.
- Vitart, V., G. Bencic, C. Hayward, J. Skunca Herman, J. Huffman, S. Campbell, et al. 2010. New loci associated with central cornea thickness include COL5A1, AKAP13 and AVGR8. *Hum. Mol. Genet.* 19:4304–4311.
- Vithana, E. N., T. Aung, C. C. Khor, B. K. Cornes, W. T. Tay, X. Sim, et al. 2011. Collagen-related genes influence the glaucoma risk factor, central corneal thickness. *Hum. Mol. Genet.* 20:649–658.
- Wang, L., Y. Guo, W. J. Huang, X. Ke, J. L. Poyet, G. A. Manji, et al. 2001. Card10 is a novel caspase recruitment domain/membrane-associated guanylate kinase family member that interacts with BCL10 and activates NF-kappa B. *J. Biol. Chem.* 276:21405–21409.
- Wang, K., M. Li, and H. Hakonarson. 2010. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* 38:e164.
- Wiggs, J. L., J. H. Kang, B. L. Yaspan, D. B. Mirel, C. Laurie, A. Crenshaw, et al. 2011. Common variants near CAV1 and CAV2 are associated with primary open-angle glaucoma in Caucasians from the USA. *Hum. Mol. Genet.* 20:4707–4713.
- Wiggs, J. L., B. L. Yaspan, M. A. Hauser, J. H. Kang, R. R. Allingham, L. M. Olson, et al. 2012. Common variants at 9p21 and 8q22 are associated with increased susceptibility to optic nerve degeneration in glaucoma. *PLoS Genet.* 8: e1002654.
- Wolfs, R. C., C. C. Klaver, R. S. Ramrattan, C. M. van Duijn, A. Hofman, and P. T. de Jong. 1998. Genetic risk of primary open-angle glaucoma. Population-based familial aggregation study. *Arch. Ophthalmol.* 116:1640–1645.
- Wu, G. L., J. L. Yuan, X. D. Huang, J. F. Rong, L. X. Zhang, Y. P. Liu, et al. 2013. Evaluating the expression of CARMA3 as a prognostic tumor marker in renal cell carcinoma. *Tumour Biol.* 34:3431–3435.
- Xie, C., Y. Han, L. Fu, Q. Li, X. Qiu, and E. Wang. 2014. Overexpression of CARMA3 is associated with advanced tumor stage, cell cycle progression, and cisplatin resistance in human epithelial ovarian cancer. *Tumour Biol.* 35:7957–7964.
- Zhao, T., Z. Miao, Z. Wang, Y. Xu, J. Wu, X. Liu, et al. 2013. CARMA3 overexpression accelerates cell proliferation and inhibits paclitaxel-induced apoptosis through NF-kappaB regulation in breast cancer cells. *Tumour Biol.* 34:3041–3047.

Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1. Chromatograms of all eight *CARD10* mutation carriers in the POAG cohort.

Table S1. PCR primers for validation of *CARD10* (NM_014550.3) variants by direct sequencing.

Table S2. List of genes found on GWAS to be associated with POAG or endophenotype.

Table S3. Summary of disease burden in genes near GWAS associated SNPs.

Table S4. 86 randomly selected genes (Set 1 of 5).

Table S5. 86 randomly selected genes (Set 2 of 5).

Table S6. 86 randomly selected genes (Set 3 of 5).

Table S7. 86 randomly selected genes (Set 4 of 5).

Table S8. 86 randomly selected genes (Set 5 of 5).