

On a Modified Iterative Method for the Solutions of Advection Model

S. O. Edeki, *Member, IAENG*, G. O. Akinlabi, and A. S. Osheku

Abstract— Variational Iterative Method (VIM) has been reported in literature as a powerful semi-analytical method for solving linear and nonlinear differential equations; however, it has also been shown to have some weaknesses such as calculation of unneeded terms, and time-consumption regarding repeated calculations for series solution. In this work, a modified VIM is applied for approximate-analytical solution of homogeneous advection model. The result attest to the robustness and efficiency of the proposed method (MVIM).

Index Terms— Exact solutions; modified VIM; HPM; advection model.

I. INTRODUCTION

In pure and applied sciences, Nonlinear Partial Differential Equations (PDEs) stand out for modelling real life problems. An example of such nonlinear model is the advection model of the form:

$$\begin{aligned} u_t(x,t) + uu_x &= f(x,t), \\ u(x,0) &= h(x). \end{aligned} \quad (1.1)$$

The importance of providing solutions (numerical or exact) to linear and nonlinear differential equations cannot be overemphasized. This has led to various methods of solution [1-10].

Ji-Huan He [11] proposed the popular variational iteration method (VIM) for a nonlinear differential equation. VIM has been widely applied [12-16]. However, some of the weaknesses of the VIM are remarked. This include: repeated computations and computations of unneeded terms, and so on. Hence, the VIM modification [17].

In this work, the modified VIM is applied to advection model for approximate-analytical solutions.

II. VIM AND THE MODIFIED VIM [17]

Considering the general nonlinear PDE of the form:

Manuscript received March 20, 2017. Revised March 31, 2017.
S.O. Edeki (e-mail: soedeki@yahoo.com) is with the Department of Mathematics, Covenant University, Canaanland, Ota, Nigeria.

G.O. Akinlabi (e-mail: grace.akinlabi@covenantuniversity.edu.ng) is with the Department of Mathematics, Covenant University, Canaanland, Ota, Nigeria.

A. S. Osheku is with the Department of Mathematics, University of Ibadan, Nigeria.

$$\begin{cases} Lu(x,t) + Ru(x,t) + Nu(x,t) = f(x,t) \\ u(x,t) = h(x) \end{cases} \quad (2.1)$$

where $L = \frac{\partial}{\partial t}$, R is a linear operator whose partial derivatives are w.r.t. x , $Nu(x,t)$ is a nonlinear term associated to (2.1) and $f(x,t)$ is a source term (which may be homogeneous or inhomogeneous), thus by the classical VIM, the solution of (2.1) is expressed as:

$$u_{n+1}(x,t) = u_n(x,t) + \int_0^t \lambda (Lu_n + \overline{Ru_n} + \overline{Nu_n} - f(x,t)) ds \quad (2.2)$$

where λ is a Lagrange multiplier [11, 12] to be identified optimally via variational theory, and the terms: $\overline{Ru_n}$ and $\overline{Nu_n}$ are being considered as restricted variations such that $\delta \overline{Ru_n} = 0$ and $\delta \overline{Nu_n} = 0$. Hence, by calculating the variations w.r.t. u_n using the stationary conditions:

$$\begin{cases} \lambda'(s) = 0, \\ 1 + \lambda(s) = 0. \end{cases} \quad (2.3)$$

The Lagrange multiplier is identified as $\lambda = -1$. Therefore, (2.2) becomes:

$$\begin{cases} u_{n+1}(x,t) = u_n(x,t) \\ - \int_0^t (Lu_n + R(u_n) + N(u_n) - f(x,t)) ds. \end{cases} \quad (2.4)$$

Remark: Using (2.4) for the solution of special kind of nonlinear differential equations involve the calculation of unrequired terms, repeated calculations, and time-consumption, hence, the need for meaningful modification of the VIM. The modified VIM as proposed by [17] gives the iterative formula as follows:

$$u_{n+1}(x,t) = u_n(x,t) - \int_0^t (R(u_n - u_{n-1}) + (G_n - G_{n-1})) ds, \quad (2.5)$$

where

$$\begin{cases} u_{-1} = 0, \\ u_0 = h(x), \\ u_1 = u_0 - \int_0^t (R(u_n - u_{n-1}) + (G_n - G_{n-1}) - f) ds, \\ G_n = Nu_n + O(t^{n+1}). \end{cases} \quad (2.6)$$

III. ILLUSTRATIVE APPLICATION

In this subsection, the modified VIM is applied to homogeneous advection model as follows.

Application: Consider the following homogeneous advection model:

$$\begin{cases} u_t + uu_x = 0, \\ u(x, 0) = -x. \end{cases} \quad (3.1)$$

Procedure:

By applying the modified VIM, it is therefore obvious that $Lu = u_t$, $Ru = 0$ and $Nu = uu_x$. Therefore, we have:

$$u_{n+1} = u_n - \int_0^t (G_n - G_{n-1}) ds, \quad (3.2)$$

$$\begin{cases} u_{-1} = 0, \\ u_0 = -x, \\ u_1 = u_0 - \int_0^t (G_0 - G_{-1}) ds, \\ G_n = u_n (u_n)_x. \end{cases} \quad (3.3)$$

Hence, for $n = 0$, we have:

$$\begin{aligned} G_0 &= u_0 (u_0)_x = x, \\ u_1 &= -x - \int_0^t (x) ds = -x - xt. \end{aligned} \quad (3.4)$$

When $n = 1$, we have:

$$G_1 = u_1 (u_1)_x = (-x - xt)(-1 - t),$$

showing that $G_1 - G_0 = 2xt + xt^2$, as such:

$$\begin{aligned} u_2 &= u_1 - \int_0^t (G_1 - G_0) ds \\ &= -x - xt - \int_0^t (2xt + xt^2) ds \\ &= -x - xt - xt^2 - \frac{xt^3}{3} \\ &\vdots \quad \vdots \end{aligned} \quad (3.5)$$

From (3.5), the solution is therefore deduced as:

$$\begin{aligned} u(x, t) &= -x - xt - xt^2 - xt^3 + \dots \\ &= -x(1 + t^2 + t^3 + \dots) \approx \frac{x}{t-1}. \end{aligned} \quad (3.6)$$

Eq. (3.6) is thus, the exact solution of the solved problem.

IV. CONCLUDING REMARKS

This paper demonstrated the robustness and efficiency of the modified VIM. For illustration, the MVIM is applied for approximate-analytical solution of homogeneous advection

model. Applying this method does not require perturbation, or linearization.

ACKNOWLEDGEMENT

The authors are sincerely thankful to Covenant University, and the anonymous referee(s) for financial support, and constructive suggestions respectively.

REFERENCES

- [1] G.O. Akinlabi and S.O. Edeki "Perturbation Iteration Transform Method for the Solution of Newell-Whitehead-Segel Model Equations", *Journal of Mathematics and Statistics*, arXiv preprint arXiv:1703.06745.
- [2] A.A. Soliman, Numerical simulation of the generalized regularized long wave equation by He's variational iteration method, *Math. Comput. Simulation* **70** (2) (2005) 119–124.
- [3] S.O. Edeki, G.O. Akinlabi and S.A. Adeosun, "Analytic and Numerical Solutions of Time-Fractional Linear Schrödinger Equation", *Comm Math Appl*, **7**(1), (2016): 1–10.
- [4] Y. Khan, N. Faraz, "Application of modified Laplace decomposition method for solving boundary layer equation", *J. King Saud Univ*, **23** (2011):115–119.
- [5] J.H. He, A new approach to nonlinear partial differential equations, *Commun. Nonlinear Sci. Numer. Simulation* **2** (4) (1997): 230–235.
- [6] G.O. Akinlabi and S.O. Edeki "On Approximate and Closed-form Solution Method for Initial-value Wave-like Models", *International Journal of Pure and Applied Mathematics*, **107** (2), (2016): 449–456.
- [7] M.M. Rashidi, "The modified differential transform method for solving MHD boundary-layer equations", *Comput Phys Commun*, **180**, (2009).
- [8] S.O. Edeki, G.O. Akinlabi, S.A. Adeosun, "On a modified transformation method for exact and approximate solutions of linear Schrödinger equations", *AIP Conference proceedings* 1705, 020048 (2016); doi: 10.1063/1.4940296.
- [9] S.O. Edeki, G.O. Akinlabi, S.A. Adeosun, "Approximate-analytical Solutions of the Generalized Newell-Whitehead-Segel Model by He's Polynomials Method", *World Congress on Engineering 2017, WCE2017*, London, U.K. (Accepted).
- [10] S.O. Edeki, and G.O. Akinlabi, "He-Laplace Method for the Solutions of the Navier-Stokes Model", *World Congress on Engineering 2017, WCE2017*, London, U.K. (Accepted).
- [11] J.H. He, "Homotopy perturbation technique", *Computer Methods in Applied Mechanics and Engineering*, **178** (3/4): (1999): 257-262.
- [12] M.A. Abdou, A.A. Soliman, Variational iteration method for solving Burger's and coupled Burger's equations, *J. Comput. Appl. Math.* **181** (2) (2005) 245–251.
- [13] J.H. He, Variational iteration method for autonomous ordinary differential systems, *Appl. Math. Comput.* **114** (2–3) (2000) 115–123.
- [14] A.A. Soliman, A numerical simulation and explicit solutions of KdV-Burgers' and Lax's seventh-order KdV equations, *Chaos, Solitons & Fractals* **29** (2) (2006) 294–302.
- [15] M. Shakil, T. Khan, H.A. Wahab, S. Bhatti, "A Comparison of Adomian Decomposition Method (ADM) and Homotopy Perturbation Method (HPM) for Nonlinear Problems", *International Journal of Research in Applied, Natural and Social Sciences*, **1** (3), (2013): 37-48.
- [16] H. He, X.H. Wu, Construction of solitary solution and compacton-like solution by variational iteration method, *Chaos, Solitons & Fractals* **29** (1) (2006) 108–113.
- [17] T. A. Abassy, M. A. El-Tawil, H. El Zoheiry, Toward a modified variational iteration method, *Computational and Applied Mathematics, Journal of Computational and Applied Mathematics* **207** (2007) 137–147.