

References

- [1] Emmert MY, Salzberg SP, Seifert B, Rodriguez H, Plass A, Hoerstrup SP, Grünenfelder J, Falk V. Is off-pump superior to conventional coronary artery bypass grafting in diabetic patients with multivessel disease? *Eur J Cardiothorac Surg* 2011;224:233–9.
- [2] Herlitz J, Malmberg K, Karlsson BW, Rydén L, Hjalmarson A. Mortality and morbidity during a five-year follow-up in diabetics with myocardial infarction. *Acta Med Scand* 1988;224:31–8.
- [3] Fava S, Azzopardi J, Agius-Muscat H. Outcome of unstable angina in patients with diabetes mellitus. *Diabet Med* 1997;14:209–13.
- [4] The BARI Investigators. Seven-year outcome in the Bypass Angioplasty Revascularization Investigation (BARI) by treatment and diabetic status. *J Am Coll Cardiol* 2000;35:1122–9.
- [5] Abizaid A, Costa MA, Centemero M, Abizaid AS, Legrand VM, Limet RV, Schuler G, Mohr FW, Lindeboom W, Sousa AG, Sousa JE, van Hout B, Hugenholtz PG, Unger F, Serruys PW, Arterial Revascularization Therapy Study Group. Clinical and economic impact of diabetes mellitus on percutaneous and surgical treatment of multivessel coronary disease patients: insights from the Arterial Revascularization Therapy Study (ARTS) trial. *Circulation* 2001;104:533–8.

* Corresponding author. Tel.: +44 1895828550; fax: +44 1895828992.
E-mail address: drrajashahzad@hotmail.com

doi:10.1016/j.ejcts.2011.03.012

Reply to the Letter to the Editor

Reply to Raja SG

Maximilian Y. Emmert*, Volkmar Falk, Sacha P. Salzberg
Clinic for Cardiac Surgery, University Hospital Zurich, Zurich, Switzerland

Received 2 March 2011; accepted 8 March 2011; Available online
14 April 2011

Keywords: Diabetes; Coronary artery disease; Off-pump; Coronary artery bypass grafting; Surgery; Myocardial revascularisation

We thank Dr Raja for his interest in our manuscript [1].

The strategy for management of blood glucose in the perioperative period was performed in a standardised fashion, according to our institutional standards, and is in line with the recently published guidelines on the perioperative management of diabetes in cardiac and vascular surgery [2,3]. In general, perioperative normoglycaemia was aimed at, and a minimal disturbance of the patients' habitual blood glucose balance was anticipated. Hypoglycaemia was avoided whenever possible, whereas hyperglycaemia was immediately corrected to a level of 6–10 mmol l⁻¹. The evening before the operation, oral antidiabetics were stopped and only the standard dose of long-acting insulin (the patient was used to) was applied subcutaneously. On the day of operation, also any type of oral antidiabetics was avoided and no baseline insulin was applied to the patient. By contrast, the blood glucose level was measured in the early morning and was corrected prior to the operation using actrapid, if necessary. During the operation, the blood glucose level was continuously monitored; if it was too high, it was corrected using an actrapid perfusor (1–6 IE h⁻¹) and if too low, glucose 5% was infused (200 ml h⁻¹). After the operation, the therapy was changed to the habitual settings as soon as possible.

Bilateral internal mammary artery usage was not associated with increased risk for sternal wound infection in our cohort. In our opinion, this is due to the skeletonised technique [4] of harvest, which is standard at our institution and to the usage of local antibiotic prophylaxis in many patients [5].

As stated in our limitations, the study period was quite long, with most on-pump coronary artery bypass grafting (CABG) patients being from the early part of the study, whereas most of OPCAB patients were from the later part of the study period [6]. Therefore, a certain difference was visible with regard to the choice of grafts, as the usage of arterial grafts or even total arterial grafting constitutes nowadays the standard of care at our institution for patients under the age of 70 years, whereas, in earlier days, the usage of saphenous vein grafts was a common approach to revascularise these patients.

References

- [1] Raja SG. Feasibility, safety and efficacy of multivessel off-pump coronary artery bypass grafting in diabetics. *Eur J Cardiothorac Surg* 2011;40:1549–50.
- [2] Joshi GP, Chung F, Vann MA, Ahmad S, Gan TJ, Goulson DT, Merrill DG, Twersky R. Society for Ambulatory Anesthesia consensus statement on perioperative blood glucose management in diabetic patients undergoing ambulatory surgery. *Anesth Analg* 111:1378–87.
- [3] Wong J, Zoungas S, Wright C, Teede H. Evidence-based guidelines for perioperative management of diabetes in cardiac and vascular surgery. *World J Surg* 34:500–13.
- [4] Peterson MD, Borger MA, Rao V, Peniston CM, Feindel CM. Skeletonization of bilateral internal thoracic artery grafts lowers the risk of sternal infection in patients with diabetes. *J Thorac Cardiovasc Surg* 2003;126:1314–9.
- [5] Nakano J, Okabayashi H, Hanyu M, Soga Y, Nomoto T, Arai Y, Matsuo T, Kai M, Kawatou M. Risk factors for wound infection after off-pump coronary artery bypass grafting: should bilateral internal thoracic arteries be harvested in patients with diabetes? *J Thorac Cardiovasc Surg* 2008;135:540–5.
- [6] Emmert MY, Salzberg SP, Seifert B, Rodriguez H, Plass A, Hoerstrup SP, Grünenfelder J, Falk V. Is off-pump superior to conventional coronary artery bypass grafting in diabetic patients with multivessel disease? *Eur J Cardiothorac Surg* 2010.

* Corresponding author. Address: Department of Cardiovascular Surgery, University Hospital Zurich, Raemi Street 100, 8091 Zurich, Switzerland. Tel.: +41 44 255 1111; fax: +41 44 255 1111.
E-mail addresses: maximilian.emmert@usz.ch, maximilian_emmert@web.de (M.Y. Emmert).

doi:10.1016/j.ejcts.2011.03.013

Letter to the Editor

Safety of tranexamic acid in pediatric cardiac surgery: what we do not know

David Faraoni*

Department of Anesthesiology, Centre Hospitalo-Universitaire Brugmann – HUDE RF, Brussels, Belgium

Received 6 February 2011; accepted 7 March 2011; Available online
14 April 2011

Keywords: Tranexamic acid; Antifibrinolytic therapy; Transfusion; Pediatric cardiac surgery; Congenital heart disease; Seizures

I have read with interest the article of Martin et al. and I would like to make some comments [1]. The risks, benefits,