

washing and wearing gowns and gloves are important infection control measures, whereas the aerosol dissemination of influenza virus necessitates masks to prevent spread. A large prospective study using viral cultures as well as serology is needed to assess more accurately the scope of this problem in older persons.

**Ann R. Falsey, Edward E. Walsh, and Robert F. Betts**

*Departments of Infectious Diseases, Medicine, and Pediatrics, Infectious Disease Unit, University of Rochester School of Medicine and Dentistry, New York*

#### References

1. Chanock RM, Parrot RH. Acute respiratory disease in infancy and childhood: present understanding and prospects for prevention. *Pediatrics* 1965;6:21-39
2. Johnson KM, Bloom HH, Mufson MA, Chanock RM. Natural reinfection of adults by respiratory syncytial virus: possible relation to mild upper respiratory diseases. *N Engl J Med* 1962;267:68-71
3. Finger R, Anderson JT, Dicker RC, Harrison B, Doan R, Downing A, Corey L. Epidemic infections caused by respiratory syncytial virus in institutionalized young adults. *J Infect Dis* 1987;155:1335-1339

### Hepatitis C Virus in Patients with Cryoglobulinemia Type II

COLLEAGUES—The association between cryoglobulinemia and hepatitis B virus (HBV) infection has been described [1]. Cryoglobulins have been found in 6%–43% of patients with chronic liver diseases, whether or not the liver disease was due to HBV [2]. Such cryoglobulins are most often of the mixed type without a monoclonal component [3]. This association is even stronger when the analysis is reversed: Two-thirds of patients with mixed essential cryoglobulinemia have a liver disease or abnormalities in liver function tests [1, 2]. Type II cryoglobulinemia is characterized by the presence in the cryoprecipitate of a monoclonal immunoglobulin that binds to Fc portions of polyclonal IgG; most often this monoclonal component is an IgM $\kappa$ . Of those with this type of cryoglobulinemia, >50% have some liver disease, including cryptogenic chronic hepatitis or HBV chronic hepatitis [1, 2]. Although this high incidence has not been found everywhere [1], the association between chronic liver disease and cryoglobulins of type II is unlikely to be fortuitous.

Serologic markers for HBV and hepatitis C virus (HCV) antibodies [4] were measured in 10 patients with cryoglobulinemia type II (all: IgM $\kappa$ /polyclonal IgG). Six patients had positive markers for hepatitis B, and three had HCV antibodies detected by enzyme-linked immunosorbent assay (Ortho HCV ELISA; Ortho Pharmaceutical, Raritan, NJ; optical density at 490 nm: 1.7, 1.02, and >2 for a cutoff of 0.45).

Reprints and correspondence: Dr. J. A. Schifferli, Department of Medicine, Hôpital Cantonal Universitaire, 24, rue Micheli-du-Crest, 1211 Geneva 4, Switzerland.

*The Journal of Infectious Diseases* 1990;162:569-570  
© 1990 by The University of Chicago. All rights reserved.  
0022-1899/90/6202-0048\$01.00

4. Englund JA, Sullivan CJ, Jordan C, Dehner LP, Vercellotti GM, Balfour HH. Respiratory syncytial virus infection in immunocompromised adults. *Ann Intern Med* 1988;1:203-208
5. Levenson RM, Kanter OS. Fatal pneumonia in an adult due to respiratory syncytial virus. *Arch Intern Med* 1987;147:791-792
6. Garvie DG, Gray J. Outbreak of respiratory syncytial virus infection in the elderly. *Br Med J* 1980;281:1253-1254
7. Hart JC. An outbreak of respiratory syncytial virus infection in an old people's home. *J Infect* 1984;8:259-261
8. Sorvillo FJ, Huie SF, Strassburg MA, Butsumyo A, Shandra WX, Fanin SL. An outbreak of respiratory syncytial virus pneumonia in a nursing home for the elderly. *J Infect* 1984;9:252-256
9. Hendry MR, Burns JC, Walsh EE, Graham BS, Wright PF, Hemming VG, Rodriguez WJ, Kim HW, Prince GA, McIntosh K, Chanock RM, Murphy BR. Strain-specific serum antibody responses in infants undergoing primary infection with respiratory syncytial virus. *J Infect Dis* 1988;157:640-647
10. Mathur U, Bentley DW, Hall CB. Concurrent respiratory syncytial virus and influenza A infections in the institutionalized elderly and chronically ill. *Ann Intern Med* 1980;93:49-52

Only 1 of 3 patients with antibodies to HCV had hepatitis B markers: a 33-year-old man with a history of drug addiction. He had antibodies to hepatitis B surface antigen (HBsAg) but was negative for HBsAg. He had moderately elevated transaminases, and liver biopsy showed chronic fibrosing hepatitis with a negative stain for hepatitis B core antigen by immunofluorescence. The most likely diagnosis was chronic HCV hepatitis [5].

The second patient, a 44-year-old man, had membranoproliferative glomerulonephritis, which was treated successfully with steroids and cyclophosphamide, and persistently elevated transaminase levels of 4 years duration. He declined liver biopsy. The last patient was 68 years old and, except for purpura, had no complaints. Liver function tests showed a slight increase in  $\gamma$ -glutamyltransferase (59 units/l; normal, <42 units/l) and alkaline phosphatase (131 units/l; normal, <125 units/l); however, transaminases were normal.

Although the number of patients is small, our observation suggests an association between HCV and cryoglobulinemia type II. The emergence of a monoclonal rheumatoid factor has been linked to chronic viral infection (Epstein-Barr virus [6], HBV [1]), which might participate in the activation process leading to the uncontrolled proliferation of a B cell clone. Hepatitis C virus may have a similar effect.

**Manuel Pascual, Luc Perrin, Emilio Giostra,  
and Jürg A. Schifferli**

*Department of Medicine, Hôpital Cantonal Universitaire,  
Geneva, Switzerland*

#### References

1. Montagnino G. Reappraisal of the clinical expression of mixed cryoglobulinaemia. *Springer Semin Immunopathol* 1988;10:1-19
2. Zarski JP, Rougier D, Aubert H, Renversez JC, Cordonnier D, Stroebner P, Rachail M. Association cryoglobuline et maladie hépatique: fré-

- quence, nature et caractères immuno-chimiques de la cryoglobulinémie. *Gastroenterol Clin Biol* 1984;8:845-850
3. Garcia-Bragado F, Biosca M, Villar M, de la Figuera M, Rodrigo MJ, Vilardell M. Maladie hépatiques et cryoglobulinémie mixte. *Gastroenterol Clin Biol* 1985;9:456-457
  4. Choo QL, Kuo G, Weiner AJ, Overby LR, Bradley DW, Houghton M. Isolation of a cDNA clone derived from blood-borne non-A, non-B viral hepatitis genome. *Science* 1989;244:359-362
  5. Alter HJ, Purcell RH, Shih JW, Melpolder JC, Houghton M, Choo QL, Kuo G. Detection of antibody to hepatitis C virus in prospectively followed transfusion recipients with acute and chronic non-A, non-B hepatitis. *N Engl J Med* 1989;321:1494-1500
  6. Fiorini GF, Sinico RA, Winearls C, Custode P, De Guili-Morghen C, D'Amico G. Persistent Epstein-Barr virus infection in patients with type II essential mixed cryoglobulinaemia. *Clin Immunol Immunopathol* 1988;47:262-269

### ***Campylobacter jejuni* versus *Campylobacter coli* in Developing Countries: How Accurate Are Prevalence Estimates?**

COLLEAGUES—*Campylobacter* species are an important cause of infectious diarrhea, especially endemic diarrhea among children of developing countries and adults traveling to these areas. Because *Campylobacter coli* is generally thought to occur in <5% of human *Campylobacter* infections, most clinical laboratories do not attempt to distinguish between *C. coli* and *Campylobacter jejuni* [1]. Unspiciated *Campylobacter* isolates from diarrhea surveys are commonly reported as "*Campylobacter* spp.," "*C. jejuni/coli*," or simply "*C. jejuni*." As a result, the frequency of *C. coli* infections and its relative importance as a diarrheal agent are unknown for many areas of the world.

During 1984-1989, 513 diarrheic and 379 nondiarrheic stool specimens from adult US citizens living in or visiting Lima, Peru, were tested for presence of *Campylobacter* and other common bacterial enteropathogens. On the whole, *Campylobacter* organisms were a less important cause of diarrhea than anticipated, being isolated from 6.1% of cases and 2.1% of controls, accounting for only 17% of all

enteropathogens isolated from patients with diarrhea. *Campylobacter* isolates were speciated according to their ability to hydrolyze hippurate [1]. The isolation frequency for *C. jejuni* was 2.5% from stools of diarrheic individuals and 1.3% from control stools. *C. coli*, however, was less frequently isolated from cases (0.8%) than from controls (1.6%), suggesting *C. jejuni* may be the only important cause of *Campylobacter*-associated diarrhea in adult travelers to Peru. Furthermore, because 36% of *Campylobacter* isolates were *C. coli*, the true frequency of *C. jejuni*-associated diarrhea could have been overestimated by 144% if it had been assumed that most isolates were *C. jejuni*.

These limited results indicate American adults living in or traveling to Peru are at relatively low risk of developing *Campylobacter*-associated diarrhea. Not surprisingly, most of the risk is attributable to *C. jejuni*. However, the high proportion of *C. coli* isolates among all *Campylobacter* species isolated suggests that infection or transient colonization by this organism may be much more common in many areas of the world than previously considered. For Peru and other countries where *C. coli* is prevalent, the unexpectedly high proportion of *C. coli* among all *Campylobacter* isolates could lead to inflated estimates of the relative importance of *C. jejuni* if each species is not considered separately. To establish whether this phenomenon occurs in other geographic regions, we encourage diarrheal disease investigators to report *Campylobacter* isolates according to species.

**G. Pazzaglia, R. Batchelor, J. Escamilla,  
M. Bernal, and G. Nunez**

*Department of Bacteriology and Epidemiology, United States Naval Medical Research Institute Detachment, Lima, Peru*

Informed consent was obtained from participants; the US Navy's guidelines for use of human subjects were followed.

The opinions and assertions contained herein are the private ones of the authors and are not to be construed as official or reflecting the views of the US or Peruvian Navy Departments.

Financial support: US Naval Medical Research and Development Command (62770A 3M162770.A870.AR-522, 61102A 3M161102.BS13.AK-511); Surgeon General's Office of the Peruvian Navy.

Correspondence: LCDR G. Pazzaglia, MSC, USN, NAMRID, APO Miami, FL 34031. Reprints: Editorial Assistant, NAMRID, APO Miami, FL 34031.

**The Journal of Infectious Diseases** 1990;162:570  
This article is in the public domain.  
0022-1899/90/6202-0049

#### **Reference**

1. Morris GK, Patton CM. *Campylobacter*. In: Lennette EH, ed. Manual of clinical microbiology. 4th ed. Washington, DC: American Society for Microbiology, 1985