

ESMO Minimum Clinical Recommendations for diagnosis, treatment and follow-up of soft tissue sarcomas

Incidence

- The crude incidence of soft tissue sarcoma in the European Union is 1.0–3.0/100 000 per year, the mortality 0.6–0.8/100 000 per year. They are diagnosed at any age but are more frequent in older patients with a peak incidence at the age of 50 years.

Diagnosis

- Histological diagnosis and evaluation of grades are preferentially made on an incisional surgical biopsy or on the completely resected tumor. Tru-cut biopsies are also used but should be restricted to experienced centers.
- Gastrointestinal stromal tumors should be confirmed by staining for CD117.
- Specific types of small round cell tumors (extra-osseous Ewing's sarcoma, embryonal rhabdomyosarcoma) should be identified by immunohistochemistry and cytogenetics and be treated accordingly.

Staging and risk assessment

- Staging is performed by physical exam and the appropriate radiological techniques. To exclude lung metastases a CT scan is recommended in operable patients.
- For staging according to the UICC/AJCC 2002 system, tumor size is categorized as small (≤ 5 cm, T1) or large (T2) and is complemented by information about location (superficial: Ta or deep: Tb) and histological grade (using either of the grading systems G1-4 or G1-3 or low/high) as further parameters that are required for the stage grouping are shown in Table 1.

Treatment plan

- Soft tissue sarcoma requires a multidisciplinary approach by an experienced team.

Surgery

- Surgery is the main treatment for localized disease. The tumor should be removed by wide excision or by compartmental resection including the cutaneous scar and tractus of a previous biopsy. The width of the resection might be decreased in case of resistant anatomic planes such as muscular fasciae, periosteum, perineurium, if not infiltrated.
- After wide excision of high-grade sarcomas, adjuvant radiation therapy is recommended [III, B].

- In case of radical surgery obtained by compartmental excision or amputation at a large distance from the primary tumor, adjuvant radiation therapy is not necessary.
- Re-operation is recommended in case of previous marginal or intralésional resection [III, B].
- In case of completely resectable lung metastases, surgery has to be considered [III, B].

Radiation therapy

- Radiation therapy should be administered postoperatively at a dose of 60–65 Gy with a shrinking field technique in case of wide resection [II, B].
- In selected patients preoperative radiotherapy may be an option [III, B].

Chemotherapy

Localized disease:

- Preoperative chemotherapy is not standard practice for operable patients [III, B]. It can be considered together with radiotherapy in patients with borderline resectable tumors.
- Adjuvant chemotherapy is not standard practice even though it might improve distant and local control. Its impact on overall survival is still debated [II, A]. It may be considered in younger patients with large and high-grade tumors [II, C].
- In non-resectable tumors confined to an extremity, chemotherapy with or without radiotherapy or isolated hyperthermic limb perfusion with chemotherapy and/or cytokines offers an alternative to amputation [III, B].

Metastatic disease:

- Chemotherapy is the standard treatment for metastatic disease. Doxorubicin with or without ifosfamide is commonly

Table 1.

Stage	Primary tumor	Lymph nodes	Distant metastases	Grading G ₁₋₄ (or G ₁₋₃ or low / high)
IA	T _{1a} or b	N ₀ or x	M ₀	G ₁₋₂ (or G ₁ or low)
IB	T _{2a} or b	N ₀ or x	M ₀	G ₁₋₂ (or G ₁ or low)
IIA	T _{1a} or b	N ₀ or x	M ₀	G ₃₋₄ (or G ₂₋₃ or high)
IIB	T _{2a}	N ₀ or x	M ₀	G ₃₋₄ (or G ₂₋₃ or high)
III	T _{2b}	N ₀ or x	M ₀	G ₃₋₄ (or G ₂₋₃ or high)
IV	Any T Any T	N ₁ Any N	M ₀ M ₁	Any G Any G

used. Doxorubicin alone seems to be equivalent to its combinations with other agents with regard to survival, despite a higher response rate with combination regimens [II, B].

- Imatinib (STI 571, a tyrosine kinase receptor inhibitor) is the treatment of choice for gastro-intestinal stromal tumors (GIST).

Response evaluation

- Response evaluation of chemotherapy for metastatic disease should be performed after 2 or 3 cycles with the radiological exams that were positive before treatment.

Follow-up

- Early detection of recurrence might influence the possibility of a curative therapy. The patients should be followed every 3 months with history and physical examination. MRI of the site of resection of the primary tumor is proposed twice a year for the first 2–3 years and then once a year. For patients with high-grade tumors a chest X-ray is recommended every 3 to 4 months in the first 2–3 years, twice a year up to the 5th year, and once a year thereafter [IV, B].

Note

Levels of Evidence [I–V] and Grades of Recommendation [A–D] as used by the American Society of Clinical Oncology are given in square brackets. Statements without grading were considered justified standard clinical practice by the experts and the ESMO faculty.

Literature

1. Wylie JP, O'Sullivan B, Catton C, Gutierrez E. Contemporary radiotherapy for soft tissue sarcoma. *Semin Surg Oncol* 1999; 17: 33–46.
2. Sarcoma meta-analysis collaboration. Adjuvant chemotherapy for localized resectable soft-tissue sarcoma: a meta-analysis of individual data. *Lancet* 1997; 350: 1647–1654.
3. Eggermont AMM, Schraffordt Koops H, Liénard D et al. Isolated limb perfusion with high-dose tumor necrosis factor- α in combination with interferon- γ and melphalan for non resectable extremity soft tissue sarcomas: a multicentric trial. *J Clin Oncol* 1996; 14: 2653–2665.
4. O'Byrne K, Steward WP. The role of chemotherapy in the treatment of adult soft tissue sarcomas. *Oncology* 1999; 56: 13–23.
5. Frustaci S, Gherlinzoni F, De Paoli A et al. Adjuvant chemotherapy for adult soft tissue sarcomas of the extremities and girdles: results of the Italian Randomized Cooperative Trial. *J Clin Oncol* 2001; 19: 1238–1247.
6. Weiss SW. *Histological typing of soft tissue tumors*. Heidelberg: Springer-Verlag/World Health Organization 1994.
7. Van Unnik JA, Coindre JM, Contesso C et al. Grading of soft tissue sarcomas: experience of the EORTC soft tissue and bone sarcoma group. *Eur J Cancer* 1993; 29 A: 2089–2093.
8. Flugstad DL, Wilke CP, McNutt MA et al. Importance of surgical resection in the successful management of soft tissue sarcoma. *Arch Surg* 1999; 134: 856–861.
9. Lewis JJ, Leung D, Espat J et al. Effect of reresection in extremity soft tissue sarcoma. *Ann Surg* 2000; 231: 655–663.
10. Van Geel AN, Pastorino V, Jauch KW et al. Surgical treatment of lung metastases. The EORTC-STBSG study of 255 patients. *Cancer* 1996; 77: 675–682.

Coordinating authors for the ESMO Guidelines Task Force: S. Leyvraz¹ & S. Jelic²

¹Invited author, Fondation du Centre Pluridisciplinaire d'Oncologie, CHUV, Lausanne, Switzerland; ²Assigned task force member, Institute of Oncology and Radiology of Serbia, Department of Medical Oncology, Pasterova 14, 11000 Belgrade, Serbia

Approved by the ESMO Guidelines Task Force: August 2003, last update December 2004.

Correspondence to:
ESMO Guidelines Task Force
ESMO Head Office
Via La Santa 7
CH-6962 Lugano
Switzerland