

*Pure Appl. Chem.*, Vol. 73, No. 2, pp. 377–380, 2001.

© 2001 IUPAC

INTERNATIONAL UNION OF PURE AND APPLIED CHEMISTRY

INORGANIC CHEMISTRY DIVISION

COMMISSION ON NOMENCLATURE OF INORGANIC CHEMISTRY\*

# NAMES FOR MUONIUM AND HYDROGEN ATOMS AND THEIR IONS

## (IUPAC Recommendations 2001)

*Prepared for publication by*

W. H. KOPPENOL

*Laboratorium für Anorganische Chemie, Eidgenössische Technische Hochschule, Zürich, Switzerland*

\*Membership of the Commission during the preparation of this report (1997–2001) was as follows:

**Titular Members:** A. Sargeson (Chairman, 1987–1997); H. Kaesz (Chairman, 1998–present); W. H. Koppenol (Secretary, 1994–1997); C. A. Stewart (Secretary, 1998–1999); T. Damhus (Secretary, 2000–present); M. W. G. de Bolster (1987–1997); N. Connelly (1998–present); T. Damhus (1998–present); A. Dress (1996–present); R. M. Hartshorn (2000–present); A. T. Hutton (2000–present); H. Kaesz (1994–1997); A. Salzer (1991–1999); C. A. Stewart (1996–1999); **Associate Members:** G. Bergerhoff (1991–1997); J. B. Casey (1992–present); J. de Oliveira Cabral (1994–1997); P. Royo-Garcia (1994–present); Y. Yamamoto (1996–present); **National Representatives:** M. A. Bennett (Australia, 1992–present); R. Bhattacharyya (India, 1998–2001); J. Cernak (Slovakia, 2000–present); C. S. Chin (Korea, 1992–present); N. Coville (South Africa, 1998–2001); L. I. Elding (Sweden, 1994–1999); C. A. L. Filgueiras (Brazil, 1998–present); P. Fodor-Csányi (Hungary, 1991–present); E. W. Godly (United Kingdom, 1987–1997); V. Kolotov (Russia, 1994–1997); N. Külcü (Turkey, 2000–present); R. S. Laitinen (Finland, 1994–present); K. Mackay (New Zealand, 1987–1999); D. Meyerstein (Israel, 1994–present); H. Nakazawa (Japan, 1996–1999); E. Nordlander (Sweden, 2000–present); Florian P. Pruchnik (Poland, 1998–present); Prof. J. Reedijk (Netherlands, 1987–1999); M. Zulu (Republic of South Africa, 1989–1997).

---

*Republication or reproduction of this report or its storage and/or dissemination by electronic means is permitted without the need for formal IUPAC permission on condition that an acknowledgment, with full reference to the source along with use of the copyright symbol ©, the name IUPAC, and the year of publication, are prominently visible. Publication of a translation into another language is subject to the additional condition of prior approval from the relevant IUPAC National Adhering Organization.*

# Names for muonium and hydrogen atoms and their ions

## (IUPAC Recommendations 2001)

*Abstract:* Muons are short-lived species with an elementary positive or negative charge and a mass 207 times that of the electron. These recommendations concern positive muons, given the short lifetime of negative muons. A positive muon mimics a light hydrogen nucleus, and names are given in analogy to existing names for hydrogen-containing compounds. A particle consisting of a positive muon and an electron ( $\mu^+e^-$ ) is named “muonium” and has the symbol Mu. Examples: “muonium chloride,”  $\text{MuCl}$ , is the equivalent of deuterium chloride,  ${}^2\text{HCl}$  or  $\text{DCl}$ ; “muoniomethane”,  $\text{CH}_3\text{Mu}$ , is the product of the muonation of methane; and  $\text{NaMu}$  is “sodium muonide.”

### INTRODUCTION

The name “muon” is used in physics for the short-lived species that belong to the lepton family and that are designated with the nuclear symbols  $\mu^+$  and  $\mu^-$ , each having a mass 207 times that of the electron [1]. Muons are produced artificially from high-energy proton beams at specialized facilities. A negative muon can replace an electron in the 1s orbital of an atom, in which case this atom is called a “muonic” atom:  $\text{H}^+\mu^-$  is “muonic hydrogen”. Replacement of an electron by a muon in other atoms is possible. Negative muons have a shorter life time than positive muons and are currently thought not to be chemically relevant [1]. This recommendation is therefore concerned with *positive* muons.

During its lifetime of ca. 2  $\mu\text{s}$  a positive muon can take on the role of a “light” hydrogen nucleus, and in the literature one finds the name “muonium” and the symbol “Mu” for the particle  $\mu^+e^-$  [1,2]. Muonium is formed near the end of the track of a positive muon when it abstracts an electron [1], and can be regarded as a light isotope of hydrogen. Although chemical reactions of muonium atoms have been studied for more than two decades, the nomenclature of muonium and related species has not been addressed by IUPAC. In this recommendation, names are given for muonium ions in analogy to the existing names for hydrogen and its isotopes [3].

### NOMENCLATURE

A summary of muonium and hydrogen nomenclature is given in Table 1.

These recommendations lead to the following related terms:

- “muonido” indicates muonium as a ligand (*cf.* hydrido);
- “muonio” is the prefix (*cf.* deuterio) used in substitutive nomenclature to denote replacement of hydrogen by muonium; this process is called “muonation”;
- “muonation” is the equivalent of protonation.

**Table 1** Comparison of names and symbols of muonium species with hydrogen equivalents [3].

Isotope	Particle	Name	Symbol
Muonium	$\mu^+$	muon	$\text{Mu}^+$
	$\mu^+e^-$	muonium	$\text{Mu}^\bullet$
	$\mu^+(e^-)_2$	muonide	$\text{Mu}^-$
Protium	$p^+$	proton	$^1\text{H}^+$
	$p^+e^-$	protium	$^1\text{H}^\bullet$
	$p^+(e^-)_2$	protide	$^1\text{H}^-$
Deuterium	$d^+$	deuteron	$^2\text{H}^+ (\text{D}^+)$
	$d^+e^-$	deuterium	$^2\text{H}^\bullet (\text{D}^\bullet)$
	$d^+(e^-)_2$	deuteride	$^2\text{H}^- (\text{D}^-)$
Tritium	$t^+$	triton	$^3\text{H}^+ (\text{T}^+)$
	$t^+e^-$	tritium	$^3\text{H}^\bullet (\text{T}^\bullet)$
	$t^+(e^-)_2$	tritide	$^3\text{H}^- (\text{T}^-)$
Hydrogen (naturally occurring mixture of isotopes)		hydron	$\text{H}^+$
		hydrogen	$\text{H}^\bullet$
		hydride	$\text{H}^-$

## Examples

- MuCl is named “muonium chloride”, which yields positive muons (or muonium ions),  $\text{Mu}^+$ , and chloride ions,  $\text{Cl}^-$ , upon dissociation. A hydrogen equivalent would be “deuterium chloride”,  $^2\text{HCl}$  or  $\text{DCl}$ .
- $\text{CH}_3\text{Mu}$ , “muoniomethane”, is a product of muonation of methane, while tritiation leads to “tritiomethane”.
- $\text{NaMu}$  is named “sodium muonide”.
- $^1\text{HMuO}$  is named “muonium protium oxide”; for an unspecified isotope of hydrogen the name changes to “hydrogen muonium oxide”. A substitutive name is “muoniooxidane”.  $\text{MuO}^-$  is named “muoniumoxide(1-),” or “muoniooxidanide”.
- As the systematic name of  $^1\text{H}_2$  is “diprotium”,  $^1\text{HMu}$  is named “muonium protium”,  $\text{HMu}$  “hydrogen muonium” and  $\text{Mu}_2$  “dimuonium”. Note that the particle  $\mu^+\mu^-$  would be named “muonic muonium” [2].
- Addition of muonium to a double bond results in the formation of a radical. For example, the reaction of  $\text{Mu}^\bullet$  with benzene could yield “6-muoniocyclohexa-2,4-dien-1-yl” or “4-muoniocyclohexa-2,5-dien-1-yl”.

**REFERENCES AND NOTES**

1. D. C. Walker. *Acc. Chem. Res.* **18**, 167–173 (1985).
2. IUPAC. *Quantities, Units, and Symbols in Physical Chemistry*, 2<sup>nd</sup> ed., I. M. Mills, T. Cvitaš, K. Homann, N. Kallay, and K. Kuchitsu (Eds.), p. 93, Blackwell Scientific Publications, Oxford (1993). Please note that in the 1988 edition of this book, on p. 85, muonium is incorrectly defined as a particle with the composition  $\mu^+\mu^-$ .
3. J. F. Bunnett and R. A. Y. Jones. *Pure Appl. Chem.* **60**, 1115–1116 (1988).