

J Antimicrob Chemother 2010; **65**: 1826–1837

doi:10.1093/jac/dkq203

Advance publication 9 June 2010

Molecular analysis of methicillin-resistant *Staphylococcus pseudintermedius* of feline origin from different European countries and North America

Kristina Kadlec¹, Stefan Schwarz^{1*}, Vincent Perreten², Ulrika Grönlund Andersson³, Maria Finn³, Christina Greko³, Arshnee Moodley⁴, Stephen A. Kania⁵, Linda A. Frank⁵, David A. Bemis⁵, Alessia Franco⁶, Manuela Iurescia⁶, Antonio Battisti⁶, Birgitta Duim⁷, Jaap A. Wagenaar⁷, Engeline van Duijkeren⁷, J. Scott Weese⁸, J. Ross Fitzgerald⁹, Alexandra Rossano² and Luca Guardabassi⁴

¹Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut, Neustadt-Mariensee, Germany; ²Institute of Veterinary Bacteriology, University of Berne, Berne, Switzerland; ³National Veterinary Institute, SVA, Uppsala, Sweden; ⁴Department of Veterinary Disease Biology, Faculty of Life Sciences, University of Copenhagen, Frederiksberg, Denmark; ⁵University of Tennessee, Knoxville, USA; ⁶Istituto Zooprofilattico Sperimentale delle Regioni Lazio e Toscana, Rome, Italy; ⁷Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands; ⁸University of Guelph, Guelph, Canada; ⁹The Roslin Institute and Centre for Infectious Diseases, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, UK

*Corresponding author. Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Höltystr. 10, 31535 Neustadt-Mariensee, Germany. Tel: +49-5034-871-241; Fax: +49-5034-871-246; E-mail: stefan.schwarz@fli.bund.de

Keywords: MLST, PFGE, *spa* typing, genotyping, antimicrobial resistance, cats, zoonosis

Sir,
Staphylococcus pseudintermedius is the most frequent causative agent of canine pyoderma and may also be associated with wound infections, urinary tract infections and otitis externa in dogs.¹ Although more rarely, *S. pseudintermedius* causes infections in cats and has also been identified in infections of humans.^{1–3} The latter observation highlights the zoonotic potential of *S. pseudintermedius*. *S. pseudintermedius* appears to be able to readily accumulate antimicrobial resistance genes^{1,4} and, in recent years, a rapid increase in methicillin resistance has been observed.^{5,6} A recent multicentre study on methicillin-

resistant *S. pseudintermedius* (MRSP) of canine origin, obtained in different European countries as well as in the USA and Canada, revealed that most of the canine strains exhibited resistance to virtually all classes of antimicrobial agents approved for veterinary use.⁴ This represents a major therapeutic challenge for veterinarians in Europe and North America. Moreover, this multicentre study revealed that among MRSP from dogs two distinct, dominant clones—as identified by multilocus sequence typing (MLST), *spa* typing, SCCmec typing and SmaI PFGE—have disseminated across Europe and North America.⁴ In contrast to the wealth of data available for MRSP isolates of canine origin, little data are available about MRSP from cats.

The aim of the present study was to characterize MRSP isolates from cats in different countries for their genetic relationships and antimicrobial resistance phenotypes and genotypes. Twelve epidemiologically unrelated MRSP isolates of feline origin were identified during 2006–08 in five different countries (Table 1). Eleven isolates were from clinical disease conditions, including septicaemia, urinary tract infections, nephritis, rhinitis, wound infection and pneumonia. The remaining isolate was obtained from a nasal swab of an apparently healthy cat (Table 1). All isolates were confirmed to be *S. pseudintermedius* by MboI digestion of a PCR-amplified internal fragment of the *pta* gene.⁴ For a better comparison with data of canine MRSP isolates, MLST,⁷ *spa* typing,⁸ SmaI PFGE and SCCmec typing was performed as recently described.⁴ MICs of 17 antimicrobial agents were determined using the VetMIC™ microdilution panels (National Veterinary Institute, Uppsala, Sweden) as previously described and evaluated using the breakpoints of the CLSI.⁴ MICs of rifampicin, mupirocin and quinupristin/dalfopristin were determined by Etest® (AB Biodisk, Solna, Sweden). Antibiotic resistance genes were detected using either a microarray or specific PCR assays as described previously.⁴

All 12 feline *S. pseudintermedius* proved to be MRSP by oxacillin MICs of >16 mg/L and carriage of the *mecA* gene. Despite the diverse geographical origins, the 11 European MRSP isolates shared the same MLST type ST71, *spa* type t02 and SCCmec type II–III. SCCmec type II–III is a hybrid of SCCmec II (2A) from *Staphylococcus epidermidis* and SCCmec III from *Staphylococcus aureus*.⁹ PFGE analysis identified three different patterns J, N and O among these European feline MRSP isolates. The single MRSP isolate from Canada harboured an SCCmec type V element and exhibited *spa* type t23, MLST type ST100 and PFGE pattern B. A comparison with the data of the multicentre study on MRSP in dogs⁴ revealed that PFGE patterns B, N and O were exclusive to feline MRSP isolates whereas pattern J was the dominant PFGE pattern among European canine MRSP.⁴ The Canadian feline MRSP isolate also differed largely in its resistance phenotype and genotype from the European feline MRSP isolates. It was only resistant to β -lactam antibiotics via *mecA* and *bla_Z*, and to tetracyclines via *tet(M)*. In contrast, the European isolates

Table 1. Characteristics of the 12 MRSP isolates from cats investigated in this study

Isolate no.	Country ^a	Disease condition	SCCmec type	<i>spa</i> type	MLST type	PFGE pattern	Resistance phenotype ^b	Resistance genotype
E028	I	septicaemia	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E029	I	nephritis	II–III	t02	ST71	N	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E031	I	septicaemia	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E037	I	septicaemia	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E047	CH	rhinitis	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E051	CH	wound infection	II–III	t02	ST71	J	BLA, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E053	CH	urinary tract infection	II–III	t02	ST71	J	BLA, TET, ML, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E060	CH	urinary tract infection	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E072	CH	urinary tract infection	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E076	NL	urinary tract infection	II–III	t02	ST71	O	BLA, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E114	D	healthy	II–III	t02	ST71	J	BLA, TET, ML, CHL, TMP, GEN, KAN, STR, FQ	<i>mecA, blaZ, tet(K), erm(B), cat_{pC221}, dfrG, aac(6′)-Ie/aph(2′)-Ia, aph(3′)-III, ant(6′)-Ia</i>
E132	CAN	pneumonia	V	t23	ST100	B	BLA, TET	<i>mecA, blaZ, tet(M)</i>

^aI, Italy; CH, Switzerland; NL, the Netherlands; D, Germany; CAN, Canada.

^bBLA, β -lactam antibiotics; CHL, chloramphenicol; FQ, fluoroquinolones; GEN, gentamicin; KAN, kanamycin; ML, macrolides/lincosamides; STR, streptomycin; TET, tetracyclines; TMP, trimethoprim.

exhibited three different expanded resistance phenotypes and genotypes (Table 1). All isolates were resistant to β -lactam antibiotics (*mecA*, *bla_Z*), macrolides/lincosamides [*erm(B)*], gentamicin/kanamycin [*aac(6')-Ie/aph(2')-Ia*], kanamycin [*aph(3')-III*], streptomycin [*ant(6')-Ia*], trimethoprim [*dhfrG*] and ciprofloxacin. Moreover, all but one isolate from Switzerland and all but two isolates from Switzerland and the Netherlands were resistant to chloramphenicol (*cat_{PC221}*) and to tetracyclines [*tet(K)*], respectively. However, all feline MRSP isolates were susceptible to mupirocin, rifampicin, quinupristin/dalfopristin, linezolid and vancomycin, which are important for decolonization of humans or represent 'antimicrobial agents of last resort' for the treatment of methicillin-resistant *S. aureus* (MRSA) infections in humans.

A comparison with the results of the genetic analysis of canine MRSP showed that the Canadian feline MRSP differed in all characteristics, except SCCmec type V, from the dominant canine MRSP clone in North America, which is characterized by ST68 (MLST)-C (PFGE)-t06 (*spa*)-V (SCCmec).⁴ In contrast, 9 of the 11 European feline MRSP isolates were identified as members of the previously described dominant clonal lineage among canine MRSP in Europe, which is characterized by ST71 (MLST)-J (PFGE)-t02 (*spa*)-II-III (SCCmec).⁴ This observation strongly suggested an exchange of MRSP isolates between dogs and cats in Europe. Whether the feline MRSP isolate from Canada represents a member of a new MRSP clone with a particular tropism for cats remains to be determined.

Acknowledgements

We are grateful to the institutions and diagnostic laboratories that provided MRSP isolates. We thank Fabiola Feltrin, Serena Lorenzetti, Angela Ianzano, Cinzia Onorati, Mike Schiwiek and Kerstin Meyer for expert technical assistance.

Funding

This study is the result of a self-funded project conducted by all the participating institutions.

Transparency declarations

None to declare.

References

- Weese JS, van Duijkeren E. Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine. *Vet Microbiol* 2010; **140**: 418–29.
- Van Hoovels L, Vankeerberghen A, Boel A et al. First case of *Staphylococcus pseudintermedius* infection in a human. *J Clin Microbiol* 2006; **44**: 4609–12.
- Campanile F, Bongiorno D, Borbone S et al. Characterization of a variant of the SCCmec element in a bloodstream isolate of *Staphylococcus intermedius*. *Microb Drug Resist* 2007; **13**: 7–10.
- Perreten V, Kadlec K, Schwarz S et al. Clonal spread of methicillin-resistant *Staphylococcus pseudintermedius* in Europe and North America: an international multicentre study. *J Antimicrob Chemother* 2010; **65**: 1145–54.

5 Loeffler A, Linek M, Moodley A et al. First report of multiresistant, *mecA*-positive *Staphylococcus intermedius* in Europe: 12 cases from a veterinary dermatology referral clinic in Germany. *Vet Dermatol* 2007; **18**: 412–21.

6 Bemis DA, Jones RD, Frank LA et al. Evaluation of susceptibility test breakpoints used to predict *mecA*-mediated resistance in *Staphylococcus pseudintermedius* isolated from dogs. *J Vet Diagn Invest* 2009; **21**: 53–8.

7 Bannoehr J, Ben Zakour NL, Waller AS et al. Population genetic structure of the *Staphylococcus intermedius* group: insights into *agr* diversification and the emergence of methicillin-resistant strains. *J Bacteriol* 2007; **189**: 8685–92.

8 Moodley A, Stegger M, Ben Zakour NL et al. Tandem repeat sequence analysis of staphylococcal protein A (*spa*) gene in methicillin-resistant *Staphylococcus pseudintermedius*. *Vet Microbiol* 2009; **135**: 320–6.

9 Descloux S, Rossano A, Perreten V. Characterization of new staphylococcal cassette chromosome *mec* (SCCmec) and topoisomerase genes in fluoroquinolone- and methicillin-resistant *Staphylococcus pseudintermedius*. *J Clin Microbiol* 2008; **46**: 1818–23.

J Antimicrob Chemother 2010

doi:10.1093/jac/dkq208

Advance publication 9 June 2010

Molecular characterization of plasmids encoding CTX-M-15 extended-spectrum β -lactamase associated with the ST131 *Escherichia coli* clone in Belgium

Isabel Montesinos¹, Hector Rodriguez-Villalobos^{2*}, Ricardo De Mendonça¹, Pierre Bogaerts³, Ariane Deplano¹ and Yuri Glupczynski³

¹Laboratoire de Bactériologie, Hôpital Universitaire Erasme, Université Libre de Bruxelles, B-1070 Bruxelles, Belgium;

²Laboratoire de Bactériologie, Cliniques universitaires Saint-Luc, Université catholique de Louvain, B-1200 Bruxelles, Belgium;

³Laboratoire de Bactériologie, Cliniques universitaires de Mont-Godinne, Université catholique de Louvain, B-5530 Yvoir, Belgium

*Corresponding author. Tel: +32-2-764-67-32; Fax: +32-2-764-94-40; E-mail: hector.rodriguez@uclouvain.be

Keywords: ESBLs, B2 phylogroup, pEK499, pC15

Sir,

Previous studies performed in a large number of Belgian hospitals showed the dissemination of a major clone of the virulent B2 group O25b:H4-ST131 CTX-M-15-producing *Escherichia coli*.¹ This *E. coli* ST131 clone has been reported worldwide and represents a major public health problem.² The present study sought to characterize *bla*_{CTX-M-15}-containing plasmids associated with ST131 *E. coli* CTX-M-15 isolates recovered in Belgian hospitals. This specific clone was detected from clinical specimens of patients hospitalized at the Erasme hospital in Brussels since 2001, as well as in 18 other Belgian hospitals