

Letters to the Editor

European Journal of Orthodontics 32 (2010) 228
doi:10.1093/ejo/cjq001

Sir,

We write to comment on the article by Kirveskari and Jämsä (2009). Therein, the authors reported that among adult females, who considered their work physically stressful to the cervicobrachial musculature, ‘meticulous occlusal adjustment aimed at elimination of all occlusal interferences’ (p. 491) at the initiation of the investigation and during annual follow-ups decreased later demand for therapy of symptoms located in the head, face and/or cervicobrachial regions. These findings complement results from an earlier randomized controlled trial in which it was shown that regular biannual occlusal adjustment over a period of 4 years among healthy children and adolescents led to a statistically significant reduction of the incidence of symptoms and signs of temporomandibular disorders (TMD) (Kirveskari *et al.*, 1998).

First of all, the authors need to be congratulated on their methodologically sound longitudinal studies in which participants were monitored over a period of 48 months. Based on their findings, Kirveskari and Jämsä (2009) consider ‘occlusal interferences as a structural health risk for symptoms in the head and neck region’ (p. 492). While we agree that the results indicate that the occlusion may have an influence on the neuromuscular system of individual patients, we consider the notion ‘that occlusal interferences are a health risk’ (p. 494) an over-interpretation that is likely to obfuscate a more physiological explanation. In fact, we believe that the outcome of both investigations may be construed as follows:

It is well-established in occupational physiology that repetitive, long-lasting, low-intensity muscle loading, which selectively and continuously activates small type I motor units (Cinderella hypothesis), may lead to muscle pain due to metabolic exhaustion and damage of single motor units (Zennaro *et al.*, 2004; Visser and van Dieën, 2006; Staal *et al.*, 2007). It is conceivable that a similar mechanism may also occur within masticatory muscles of susceptible patients during sustained motor activity, such as prolonged tooth grinding, jaw clenching or non-physiological daily tooth contacts (cf. Chen *et al.*, 2007).

There is good evidence that based on the inherent functional heterogeneity of the masticatory muscles and their resulting differential activation behaviour (Blanksma and van Eijden, 1995; Schindler *et al.*, 2005, 2006; Farella *et al.*, 2009), even minimal positional changes of the mandible, as induced, for instance, by occlusal adjustment, build-up of tooth surfaces (Kirveskari and Jämsä, 2009, p. 491) or insertion of oral

splints (Schindler *et al.*, 2000), alter the recruitment patterns within the masticatory muscles (Schindler *et al.*, 2005, 2006). These biomechanical modifications, in turn, are likely to reduce stress concentrations in particular muscle regions and, as a consequence, unload localized painful areas (Türp and Schindler, 2003). Hence, systematic (and periodic) alterations of occlusal surfaces at specific time intervals, as carried out in the two investigations discussed here (Kirveskari *et al.*, 1998; Kirveskari and Jämsä, 2009), may have decreased the duration and amount of stereotyped loading of motor units of susceptible subjects and, therefore, reduced the incidence of muscle pain.

The idea that any therapeutic increase or decrease of the mandibular position (of course, within a physiological range) may be useful for regional pain reduction in the temporomandibular joints and/or the masticatory muscles [and the adjacent cervicobrachial musculature (cf. Ciancaglini *et al.*, 1999; Sipilä *et al.*, 2002; Rantala *et al.*, 2003)] may serve as an explanatory model not only for the reported effects after occlusal adjustment (Kirveskari *et al.*, 1998; Kirveskari and Jämsä, 2009) but also for any other therapeutically induced alteration of the three-dimensional intermaxillary relationship. Therefore, and considering the ‘practically universal presence of interferences’, it is appreciated that Kirveskari and Jämsä (2009) judiciously eschew generalized ‘prophylactic elimination of occlusal interferences’.

Hans J. Schindler
Department of Prosthodontics
Dental School
University of Heidelberg
Germany

Jens C. Türp
Department of Reconstructive Dentistry and
Temporomandibular Disorders
Dental School
University of Basel
Switzerland

References

- Blanksma N G, van Eijden T M 1995 Electromyographic heterogeneity in the human temporalis and masseter muscles during static biting, open/close excursions, and chewing. *Journal of Dental Research* 74: 1318–1327

- Chen C Y, Palla S, Erni S, Sieber M, Gallo L M 2007 Nonfunctional tooth contact in healthy controls and patients with myogenous facial pain. *Journal of Orofacial Pain* 21: 185–193
- Ciancaglini R, Testa M, Radaelli G 1999 Association of neck pain with symptoms of temporomandibular dysfunction in the general adult population. *Scandinavian Journal of Rehabilitation Medicine* 31: 17–22
- Farella M, Palumbo A, Milani S, Avecone S, Gallo L M, Michelotti A 2009 Synergist coactivation and substitution pattern of the human masseter and temporalis muscles during sustained static contractions. *Clinical Neurophysiology* 120: 190–197
- Kirveskari P, Jämsä T 2009 Health risk from occlusal interferences in females. *European Journal of Orthodontics* 31: 490–495
- Kirveskari P, Jämsä T, Alanen P 1998 Occlusal adjustment and the incidence of demand for temporomandibular disorder treatment. *Journal of Prosthetic Dentistry* 79: 433–438
- Rantala M A 2003 Temporomandibular joint related painless symptoms, orofacial pain, neck pain, headache, and psychosocial factors among non-patients. *Acta Odontologica Scandinavica* 61: 217–222
- Schindler H J, Rong Q, Spieß W E L 2000 Der Einfluss von Aufbissschienen auf das Rekrutierungsmuster des Musculus temporalis. *Deutsche zahnärztliche Zeitschrift* 55: 575–581
- Schindler H J, Türp J C, Blaser R, Lenz J 2005 Differential activity patterns in the masseter muscle under simulated clenching and grinding forces. *Journal of Oral Rehabilitation* 32: 552–563
- Schindler H J, Rues S, Türp J C, Lenz J 2006 Heterogeneous activation of the medial pterygoid muscle during simulated clenching. *Archives of Oral Biology* 51: 498–504
- Sipilä K 2002 Temporomandibular disorders, occlusion, and neck pain in subjects with facial pain: a case-control study. *Cranio: the Journal of Craniomandibular Practice* 20: 158–164
- Staal J B, de Bie R A, Hendriks E J M 2007 Aetiology and management of work-related upper extremity disorders. *Best Practice Research Clinical Rheumatology* 21: 123–133
- Türp J C, Schindler H J 2003 Zum Zusammenhang zwischen Okklusion und Myoarthropathien: Einführung eines integrierenden neurobiologischen Modells. *Schweizer Monatsschrift für Zahnmedizin* 113: 964–977
- Visser B, van Dieën J H 2006 Pathophysiology of upper extremity muscle disorders. *Journal of Electromyography and Kinesiology* 16: 1–16
- Zennaro D, Laubli T, Krebs D, Krueger H, Klipstein A 2004 Trapezius muscle motor unit activity in symptomatic participants during finger tapping using properly and improperly adjusted desks. *Human Factors* 46: 252–266