

- ⁶ Kabir ARML, Rahman AKM, Mannan MA, Chanda SK, Chowdhury AT. Prevalence of wheeze and asthma in children of a coastal community of Bangladesh. *Bangladesh J Child Health* 1999;**23**(3/4):43–47.
- ⁷ Usherwood TP, Scrimgeour A, Barber JH. Questionnaire to measure perceived symptoms and disability in asthma. *Arch Dis Child* 1990;**65**:779–81.
- ⁸ Bangladesh Bureau of Statistics. *Statistical Pocketbook, Bangladesh 98*. Statistical Division, Ministry of Planning, Government of the People's Republic of Bangladesh, Dhaka. June 1999.
- ⁹ Mitchell I, Inglis H, Simpson H. Viral infections in wheezy bronchitis and asthma in children. *Arch Dis Child* 1972;**51**:707–11.
- ¹⁰ The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. *Lancet* 1998;**351**:1225–32.
- ¹¹ Hoque M, Barua PC, Khan AH, Hassan Q, Kabir ARML (eds). *Prevalence and Risk Factors of Measles in Under Five Children in Rural Bangladesh*. Dhaka: Institute of Child and Mother Health, 1999–2000. Sponsored by The Ministry of Health and Family Welfare, Government of Bangladesh.
- ¹² The extent of the tuberculosis problem. In: *National Guidelines for Tuberculosis. 2nd Edn*. Dhaka: TB and Leprosy Services, 1995.
- ¹³ Talukder K, Huda SN, Hassan MQ, Rahman MQ (eds). *The Relationship Between School Achievement and Health Status of School Children in 20 Primary Schools in Rural Bangladesh*. Dhaka: Institute of Child and Mother Health, 1999–2000. Sponsored by the Ministry of Health and Family Welfare, Government of Bangladesh.
- ¹⁴ Shaheen SO, Abay P, Hall AJ *et al*. Measles and atopy in Guinea-Bissau. *Lancet* 1996;**347**:1792–96.
- ¹⁵ Shirakawa T, Enomoto T, Shimaz SI, Hopkin JM. The inverse association between tuberculin responses and atopic disorder. *Science* 1997;**275**:77–79.
- ¹⁶ Becklake MR. International Union Against Tuberculosis and Lung Disease (IUATLD): initiatives in non-tuberculous lung disease. *Tubercle Lung Dis* 1995;**76**:493–504.
- ¹⁷ Lynch NR. Influence of socio-economic level on helminthic infection and allergic reactivity in tropical countries. In: Moqbel R (ed.) *Allergy and Immunity to Helminths: Common Mechanisms or Divergent Pathways*. Bristol: Taylor and Francis, 1992, pp.51–62.
- ¹⁸ National Health and Medical Research Council (NHMRC). *Asthma in Australia. Strategies for Reducing Morbidity and Mortality*. Report of the NHMRC working party on asthma associated deaths. Canberra: AGPS, 1988.
- ¹⁹ Strachan DP. Hay fever, hygiene, and household size. *BMJ* 1989;**299**:1259–60.
- ²⁰ Martinez FD. Role of viral infections in the inception of asthma and allergies during childhood: could they be protective? *Thorax* 1995;**49**:1189–91.
- ²¹ Lin S, Fitzgerald E, Hwang SA, Munsie JP, Stark A. Asthma hospitalization rates and socio-economic status in New York State (1987–1993). *J Asthma* 1999;**36**:239–51.
- ²² Claudio L, Tulton L, Doucette J, Landrigan PJ. Socioeconomic factors and asthma hospitalization rates in New York City. *J Asthma* 1999;**36**:343–50.
- ²³ Amberson M, Voigt T. Ambient air pollution and respiratory disease. *Med J Aust* 1991;**154**:543–51.
- ²⁴ Schmitzberger R, Rhomberg K, Buchele H *et al*. Effects of air pollution on the respiratory tract of children. *Pediatr Pulmonol* 1993;**15**:68–74.
- ²⁵ Mutius von E, Fritsch C, Weiland SK, Roll G, Magnussen H. Prevalence of asthma and allergic disorders among children in united Germany: a descriptive comparison. *BMJ* 1992;**305**:1395–99.
- ²⁶ United States Department of Health and Human Services, Public Health Service. *The Health Consequences of Smoking: Chronic Obstructive Lung Disease*. A report of the Surgeon General. Washington DC: US Government Printing Office, 1984.
- ²⁷ Dodge RR, Burrows B. The prevalence and incidence of asthma and asthma-like symptoms in general population sample. *Am Res Respir Dis* 1980;**122**:567–75.

Commentary: Does the 'hygiene hypothesis' provide an explanation for the relatively low prevalence of asthma in Bangladesh?

Charlotte Braun-Fahrlander

The prevalence of childhood asthma in Bangladesh is much lower than in developed countries, but similar to those of other countries in this region like Pakistan and India.¹ These differences in asthma prevalence may be attributed to a 'western lifestyle', which among other factors is characterized by a higher

level of hygiene, lower rates of infections, and small family size. Among the Bangladeshi children, those living in small families (three or less people) were more likely to suffer from asthma. The observation of an inverse relation between sibship size and atopy formed the basis of what is known today as the 'hygiene hypothesis'.² The immunological extension of this concept is the distinction of Th1 and Th2 lymphocyte populations in laboratory animals and the recognition that 'natural immunity'

Department of Environment and Health, Institute of Social and Preventive Medicine, University of Basel, Steinengraben 49, 4051 Basel, Switzerland.

to bacterial and viral infections induce a Th1 pattern of cytokine release, potentially suppressing the Th2 immune response involved in IgE mediated allergy. Thus, changes in the level of stimulation from the microbial environment associated with improvements in public health and hygiene may have indirectly influenced the postnatal development of immune functions, so as to increase predisposition to chronic allergic conditions during childhood.³

The high rates of respiratory infections, tuberculosis, measles and helminths infections in Bangladeshi children might thus contribute to lower rates of allergy. There is, however, conflicting evidence whether early (viral) infections may enhance or decrease the risk of developing asthma. A recent longitudinal birth cohort study from Germany showed that repeated lower respiratory tract infections early in life were positively associated with subsequent development of asthma, wheeze, and bronchial hyperreactivity. In contrast, early episodes of other infections (particularly viral infections) were inversely related to the development of asthma at age 7.⁴ Reverse causation seems a plausible explanation for the positive association between lower respiratory tract infections and subsequent wheeze and asthma, with lower respiratory tract infections being predictors of, rather than risk factors for, asthma.

Epidemiological evidence supporting the 'hygiene hypothesis' can further be found in studies of Italian military cadets. Respiratory allergies were less frequent in cadets with antibodies against hepatitis A virus and were inversely related to other orofecal and foodborne infections.⁵ These types of infections are likely to play an important role in countries like Bangladesh.

Studies in rural areas of Central Europe have shown that growing up on a farm and more specifically contact to farm animals was associated with a substantial decrease in risk for the development of hay fever and asthma, when children from farming families were compared to their peers living in the same villages.^{6,7} Whether increased exposure to microbial compounds has to occur early in life to affect maturation of the immune system, thereby reducing the risk for development of allergic diseases has recently been investigated in a cross-sectional study in Switzerland, Austria and Germany.⁸ Exposure of children younger than one year, compared with those aged 1–5 years, to stables and consumption of farm milk was associated with lower frequencies of asthma, hay fever and atopic sensitization. The results of these studies provide further evidence for the hygiene hypothesis and may in part explain the lower rates of asthma and allergy found in rural communities as compared to urban ones.

In Bangladesh, asthma was equally prevalent in metropolitan areas, in other urban areas and in rural areas. Thus, the effect of farming seen in European studies was not observed in Bangladesh. In a recent case-control study on indoor exposures and childhood asthma in Nepal, keeping cattle inside of the family home was associated with a risk reduction for asthma whereas keeping cattle outside of the home had no protective effect.⁹ Thus, it is conceivable that there might be variations in asthma prevalence in Bangladesh associated with keeping cattle indoors. Alternatively, if exposure to infections and unhygienic drinking water is the main source of microbial stimulation of Bangladeshi children, and if these exposures are evenly distributed between metropolitan areas, urban areas and rural communities, no variation in asthma prevalence would be expected.

In conclusion, the hygiene hypothesis may at least in part explain the lower prevalence of asthma observed in Bangladesh as compared to developed countries. However, the prevalence of asthma in Bangladesh is substantial and represents an important public health problem which might increase in parallel with the economical development of the country.

References

- ¹ Hassan RM, Kabir ARML, Mahmud AM *et al.* Self-reported asthma symptoms in children and adults in Bangladesh: findings of the National Asthma Prevalence Study. *Int J Epidemiol* 2002;**31**:483–88.
- ² Strachan DP. Hay fever, hygiene, and household size. *BMJ* 1989;**299**:1259–60.
- ³ Martinez FD, Holt PG. Role of microbial burden in aetiology of allergy and asthma. *Lancet* 1999;**354**(Suppl.2):S112–15.
- ⁴ Illi S, von Mutius E, Lau S *et al.* Early childhood infectious diseases and the development of asthma up to school age: a birth cohort study. *BMJ* 2001;**322**:390–95.
- ⁵ Matricardi PM, Rosmini F, Riondino S *et al.* Exposure to foodborne and orofecal microbes versus airborne viruses in relation to atopy and allergic asthma: epidemiological study. *BMJ* 2000;**320**:412–17.
- ⁶ Braun-Fahrlander C, Gassner M, Grize L *et al.* Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. *Clin Exp Allergy* 1999;**29**:28–34.
- ⁷ Von Ehrenstein OS, Von Mutius E, Illi S, Baumann L, Bohm O, von Kries R. Reduced risk of hay fever and asthma among children of farmers [see comments]. *Clin Exp Allergy* 2000;**30**:187–93.
- ⁸ Riedler J, Braun-Fahrlander C, Eder W *et al.* Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet* 2001;**358**:1129–33.
- ⁹ Melsom T, Brinch L, Hessen JO *et al.* Asthma and indoor environment in Nepal. *Thorax* 2001;**56**:477–81.