

ON THE WEAK HOMOLOGICAL DIMENSION OF THE GROUP ALGEBRA OF SOLVABLE GROUPS

URS STAMMBACH

Let KG denote the group algebra of the group G over a field K of characteristic 0. The *weak (homological) dimension* of KG is defined as the largest integer q for which there exist KG -modules B, C such that $\text{Tor}_q^{KG}(B, C) \neq 0$. By [1; p. 352] q is equal to the largest integer for which there exists a KG -module A such that $H_q(G, A) \neq 0$. We write $\text{w.dim } KG = q$.

The following is a corollary of the main result (Theorem 1): *For a solvable group G the weak dimension of KG is equal to the Hirsch number hG of G , $\text{w.dim } KG = hG$.*

As an application we obtain (Theorem 5):

If G is a nilpotent group of finite Hirsch number hG , then hG is equal to the largest integer q for which $H_q(G, K^+) \neq 0$, where K^+ denotes the additive group of K with trivial G -operation.

The starting point of our investigation was a result of K. W. Gruenberg [3], communicated privately to the author, which relates the Hirsch number of a torsion-free nilpotent group to the cohomological dimension of the group. We would like to express our sincere thanks to Gruenberg for showing us his result.

1. The Hirsch number of G and the weak dimension of KG

We define a class C of groups as follows: The group G is in C if and only if there exists a series of normal subgroups $\{N_i\}, i = 0, \dots, k+1$

$$1 = N_{k+1} \subseteq N_k \subseteq \dots \subseteq N_1 \subseteq N_0 = G$$

such that the successive quotients N_i/N_{i+1} for $i = 0, \dots, k$ are either locally finite or abelian. Note that C contains all solvable groups.

Definition. Let G be a group in C , and $\{N_i\}$ a series of normal subgroups with the above property. We define the Hirsch number hG of G to be the sum of the ranks of those quotients N_i/N_{i+1} which are abelian. If this sum is not finite, we write $hG = \infty$.

The fact that hG does not depend on the series of subgroups is well known; also, it follows immediately from our Theorem 1. (In the literature the Hirsch number is usually defined for a slightly different class of groups, see for example [4; p. 150]. However, the above definition is better for our purpose.)

THEOREM 1. *For G in C we have $hG = \text{w.dim } KG$.*

We first prove a special case:

PROPOSITION 2 [2]. *Let N be an abelian group of rank n . Then $\text{w.dim } KN = n$ and $H_n(N, K^+) = K$.*

Proof. We proceed by induction on n .

(a) Let N be an abelian group of rank zero, and let A be a KN -module. If N is finite of order $|N|$, then it is well known that $|N| \cdot H_p(N, A) = 0$ for all $p \geq 1$. On the other hand $H_p(N, A)$ is a K -space. Since K is a field of characteristic 0, it follows that $H_p(N, A) = 0$ for $p \geq 1$. But trivially, $H_0(N, K^+) = K$.

If N is locally finite, then N is the direct limit of its finitely generated, hence finite subgroups N_α . We then have for $p \geq 1$

$$H_p(N, A) = \varinjlim H_p(N_\alpha, A) = 0$$

(see [1; p. 125]). Of course $H_0(N, K^+) = K$.

(b) If $n \geq 1$, then N contains at least one infinite cyclic subgroup C . Consider the extension $C \rightarrow N \rightarrow N/C$. Clearly, $\text{rank}(N/C) = n-1$. The result is now deduced from the Hochschild–Serre spectral sequence [1; p. 350]

$$H_p(N/C, H_q(C, A)) \Rightarrow H_n(N, A).$$

Since C is a free group and since $H_0(C, K^+) = K = H_1(C, K^+)$ we have $\text{w.dim } KC = 1$. By induction we have $\text{w.dim } K(N/C) = n-1$. By the “maximum principle for spectral sequences” it follows that

$$\text{w.dim } KN \leq \text{w.dim } K(N/C) + \text{w.dim } KC = n-1+1 = n.$$

But choosing $A = K^+$, we obtain $H_n(N, K^+) = H_{n-1}(N/C, H_1(C, K^+)) = K$.

Note that it follows from this proposition that $H_n(N, A) = A$ for any trivial KN -module A , for such a module is simply a K -space.

LEMMA 3. *Let N be a normal subgroup of G with quotient group G' . Suppose N is abelian of rank n . Then to any KG' -module A there exists a KG' -module B such that $H_n(N, B) \cong A$ as KG' -module.*

Proof: Let B be any trivial KN -module. By Proposition 2 we have $H_n(N, B) = H_n(N, K^+) \otimes_K B = K \otimes_K B$. In virtue of the fact that N is a normal subgroup of G , the group $H_n(N, K^+) = K$ has a G' -action which is generally non-trivial. We shall denote the KG' -module $H_n(N, K^+)$ by \tilde{K} . The G' -module structure of $\tilde{K} \otimes_K B$ is defined by $x(q \otimes b) = xq \otimes xb$ for $x \in G, q \in K, b \in B$. Denote by \tilde{K}^{-1} the additive group of K together with the “inverse” G' -action, i.e., if $x \in G'$ sends $1 \in \tilde{K}$ into q , then x sends $1 \in \tilde{K}^{-1}$ into $1/q$. It is obvious that \tilde{K}^{-1} is a KG' -module. Define $B = \tilde{K}^{-1} \otimes_K A$. We claim that $\tilde{K} \otimes_K (\tilde{K}^{-1} \otimes_K A) \cong A$ as KG' -modules. Clearly this is true as K -spaces; moreover

$$x(1 \otimes 1 \otimes a) = q \otimes 1/q \otimes xa = xa.$$

LEMMA 4. *Let N be a normal subgroup of G with quotient group G' . Suppose N is abelian of rank n and suppose $hG' = m$, then $\text{w.dim } KG = m+n$.*

Proof. Consider the Hochschild–Serre spectral sequence

$$H_p(G', H_q(N, B)) \Rightarrow H_{p+q}(G, B).$$

Clearly $\text{w.dim } KG \leq n+m$. We show that there is a KG -module B such that $H_{n+m}(G, B) \neq 0$. By hypothesis there is a KG' -module A such that $H_m(G', A) \neq 0$. Take B to be the KG' -module (hence also KG -module) defined in Lemma 3. By the “maximum principle of spectral sequences” we obtain

$$H_{m+n}(G, B) = H_m(G', H_n(N, B)) = H_m(G', A) \neq 0.$$

Proof of Theorem 1. We proceed by induction on $k + h = l$ where k is the length of the given series of subgroups of G , and $h = hG$ is the Hirsch number of G .

(a) If $l = 0$, then G is the trivial group.

(b) Let $l \geq 1$. Divide by N the last nontrivial term N_k in the given series of subgroups, and let $G' = G/N$.

If N is locally finite, it is the direct limit of its finitely generated hence finite subgroups N_α . We therefore have $H_q(N, A) = \lim_{\rightarrow} H_q(N_\alpha, A) = 0$ for any KN -module A and $q \geq 1$. Hence by the Hochschild–Serre spectral sequence we obtain

$$H_q(G, A) = H_q(G', A_N).$$

This shows that $\text{w.dim } KG = \text{w.dim } (G') = h(G') = hG$.

If N is abelian of rank n , then we have $\text{w.dim } KG = \text{w.dim } K(G') + n$ by Lemma 2. Hence by induction $\text{w.dim } KG = h(G') + n = hG$.

Finally, if $hG = \infty$, then G contains subgroups of any Hirsch number; hence $\text{w.dim } KG$ cannot be finite.

Remark. The conjecture that hG is also equal to the cohomological dimension of KG turns out to be false: any infinite, locally finite abelian group is a counter example.

2. Nilpotent groups.

THEOREM 5. *Let G be a nilpotent group with $hG < \infty$. Then hG is equal to the largest integer q for which $H_q(G, K^+)$ (or equivalently $H^q(G, K^+)$) is nontrivial.*

Proof. In view of Theorem 1 we have to show that for $h = hG$, $H_h(G, K^+) \neq 0$. We prove $H_h(G, K^+) = K$ by induction on the length k of the lower central series of G .

(a) If G is abelian, Proposition 2 shows that

$$H_h(G, K^+) = K$$

for $h = hG$.

(b) If G is not abelian denote by G_k the last nonzero term of the lower central series of G . Since $[G, G_k] = 1$, the subgroup G_k must be contained in the centre of G . By the Hochschild–Serre spectral sequence

$$H_p(G/G_k, H_q(G_k, K^+)) \Rightarrow H_n(G, K^+)$$

we obtain for $p = h(G/G_k)$, $q = hG_k$, and $h = hG$

$$H_h(G, K^+) = H_p(G/G_k, H_q(G_k, K^+)) = H_p(G/G_k, K^+) = K,$$

using the fact that G_k is central, Proposition 2 and the induction assumption, respectively.

The cohomology part now follows from the remark that

$$H^q(G, K^+) = \text{Hom}_K(H_q(G, K^+), K^+).$$

References

1. H. Cartan, S. Eilenberg, *Homological Algebra* (Princeton University Press, 1956).
2. A. J. Douglas, "The weak global dimension of the group ring of abelian groups", *J London Math. Soc.* 36 (1961), 371-381.
3. K. W. Gruenberg, Springer Lecture Notes, to appear.
4. W. R. Scott, *Group Theory* (Prentice Hall, 1964).

Cornell University,
Ithaca, N.Y. 14850, U.S.A.

Eidgenössische Technische Hochschule,
Zürich, Switzerland.