



Population and economic impact of the 2013 ACC/AHA guidelines compared with European guidelines to prevent cardiovascular disease

Julien Vaucher^{1*}, Pedro Marques-Vidal², Martin Preisig³, Gérard Waeber^{1†}, and Peter Vollenweider^{1†}

¹Department of Medicine, Internal Medicine, CHUV and Faculty of Biology and Medicine, Bâtiment des Instituts, Etude CoLaus, 19, rue du Bugnon, Lausanne 1005, Switzerland; ²Institute of Social and Preventive Medicine (IUMSP), CHUV and Faculty of Biology and Medicine, Lausanne, Switzerland; and ³Department of Psychiatry, CHUV and Faculty of Biology and Medicine, Lausanne, Switzerland

Received 6 January 2014; revised 28 January 2014; accepted 29 January 2014; online publish-ahead-of-print 25 February 2014

Recently, the American College of Cardiology (ACC) and the American Heart Association (AHA) issued new guidelines on management of cardiovascular (CV) risk in primary prevention.¹ They developed a new CV risk calculator [new pooled cohort atherosclerotic CV disease (CVD) risk equation] targeting individuals between 40 and 75 years and based on four American population-based cohorts.^{2,3} Statin treatment is recommended for individuals with an estimated 10-year risk of CVD $\geq 7.5\%$, including stroke. Conversely, the guidelines of the European Society of Cardiology (ESC) advocate the use of the SCORE equation for individuals aged between 40 and 65 years, and initiation of a statin treatment is recommended if the estimated 10-year risk of death from CVD is $\geq 5\%$.⁴ Whether applying the ACC/AHA guidelines has similar public health consequences as applying the ESC guidelines is currently unknown. Hence, we used the data from a large, population-based study, to assess the country-wide population and economic impact of these new guidelines, supposing full adherence to treatment recommendations.

Data from the Swiss CoLaus study, collected between 2003 and 2006 in 3297 participants (1854 women) aged 50–75 years, were used.⁵ Ten-year CVD risk was computed according to the Swiss SCORE (ESC) and the 2013 ACC/AHA risk equations; results were extrapolated to the Swiss population of the same age group. The Swiss SCORE equation has been validated in individuals up to 75 years of age. Daily cost of treatment was estimated using one widely used statin, i.e. atorvastatin.

The results are summarized in Table 1. Irrespective of the risk equation used, the prevalence of high-risk individuals increased considerably with age, exceeding 80% among participants aged over 70 years. Extrapolated to the Swiss population, applying the ACC/AHA guidelines more than doubled the prevalence of high-risk

individuals (2.2-fold in men and 1.9 fold in women) relative to the SCORE function. The biggest differences were observed for age group 50–60 years, where the ACC/AHA guidelines led to a 30-fold increase in the number of high-risk individuals relative to the ESC guidelines. Full compliance with the ACC/AHA guidelines would also lead to an extra cost of treatment of 1.124 million CHF per day (410 million CHF, or 333.7 million €, per year).

We conclude that, relative to the ESC guidelines, the 2013 ACC/AHA guidelines lead to a considerable increase in the number of high-risk individuals susceptible of receiving statin treatment. This increase is particularly strong in the age group 50–60 years. One likely explanation for this discrepancy may be due to differences in the prevalence of CV risk factors between the USA and Europe.^{6,7} Consequently, if fully implemented, the ACC/AHA guidelines might lead to a considerable increase in primary prevention costs of CVD. Further studies are needed to validate the new ACC/AHA risk equation and to assess the cost-effectiveness of the ACC/AHA guidelines in non-US countries.

Authors' contributions

J.V. collected data and wrote most of the article. P.M.V. made the statistical analyses and wrote part of the article; M.P. revised the article for important intellectual content; G.W. and P.V. were the initiators of this study and revised the article for important intellectual content. P.M.V. had full access to the data and is the guarantor of the study.

Funding

The CoLaus study was supported by grants from the Swiss National Science Foundation (grant no: FN 33CSCO-122661 and

The opinions expressed in this article are not necessarily those of the Editors of the *European Heart Journal* or of the European Society of Cardiology.

* Corresponding author. Tel: +41 213140346, Fax: +41 213148037, Email: julien.vaucher@chuv.ch

† G.W. and P.V. contributed equally to this manuscript and are therefore both to be considered as last authors.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2014. For permissions please email: journals.permissions@oup.com

Table 1 Simulations of the population impact and of daily costs related to treatment with atorvastatin in Switzerland according to the European Society of Cardiology (ESC) or 2013 American College of Cardiology and the American Heart Association (ACC/AHA) guidelines

| ESC | Swiss population | | | Population at risk ^a | | | Ratio ACC/AHA to ESC ^b | | | Daily cost of treatment ^c |
|----------------|------------------|-----------|-----------|---------------------------------|----------------|-----------|-----------------------------------|-------|-------|--------------------------------------|
| | Men | Women | Total | Men | Women | Total | Men | Women | Total | Atorvastatin ^d |
| (50–60) | 561 013 | 551 105 | 1 112 118 | 8976 | — ^e | 8976 | — | — | — | 13 |
| (60–70) | 429 528 | 448 861 | 878 389 | 204 026 | 29 176 | 233 202 | — | — | — | 350 |
| (70–75) | 176 448 | 205 307 | 381 755 | 175 389 | 181 491 | 356 881 | — | — | — | 536 |
| All | 1 814 130 | 1 841 332 | 3 655 462 | 388 391 | 210 667 | 599 059 | — | — | — | 899 |
| ACC/AHA | | | | | | | | | | |
| (50–60) | 561 013 | 551 105 | 1 112 118 | 274 335 | 23 698 | 298 033 | 30.6 | NA | 33.2 | 447 |
| (60–70) | 429 528 | 448 861 | 878 389 | 419 649 | 169 221 | 588 869 | 2.1 | 5.8 | 2.5 | 884 |
| (70–75) | 176 448 | 205 307 | 381 755 | 176 448 | 204 486 | 380 934 | 1.0 | 1.1 | 1.1 | 572 |
| All | 1 166 989 | 1 205 273 | 2 372 262 | 870 432 | 397 405 | 1 267 836 | 2.2 | 1.9 | 2.1 | 2023 |

^aAccording to the ACC/AHA (new pooled cohort atherosclerotic CV disease risk equation) or ESC (Swiss SCORE equation) guidelines.

^bRatio of the number of subjects at risk according to ACC/AHA guidelines to the number of subjects at risk according to ESC guidelines. NA, not assessable.

^cExpressed in 1000 CHF. To obtain €, multiply by 0.814; to obtain USD, multiply by 1.110. Currency exchange rates as of 3 January 2014 were applied.

^dFixed daily price independent of dosing (10–80 mg).

^eSCORE equation do not predict CV risk for women under 60 years.

33CSC0-139468); GlaxoSmithKline; the Faculty of Biology and Medicine of Lausanne, Switzerland; and the Swiss Society of Internal Medicine.

Conflict of interest: G.W. and P.V. received an unrestricted grant from GSK to build the CoLaus study. The other authors report no other conflict of interest.

References

- Stone NJ, Robinson J, Lichtenstein AH, Merz CN, Blum CB, Eckel RH, Goldberg AC, Gordon D, Levy D, Lloyd-Jones DM, McBride P, Schwartz JS, Shero ST, Smith SC Jr, Watson K, Wilson PW. 2013 ACC/AHA Guideline on the Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013; doi: 10.1161/x01.cir.0000437738.63853.7a. [Epub ahead of print].
- Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB Sr, Gibbons R, Greenland P, Lackland DT, Levy D, O'Donnell CJ, Robinson J, Schwartz JS, Shero ST, Smith SC Jr, Sorlie P, Stone NJ, Wilson PW. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013; doi: 10.1161/x01.cir.0000437741.48606.98. [Epub ahead of print].
- Goff DC Jr, Lloyd-Jones DM, Bennett G, O'Donnell CJ, Coady S, Robinson J, D'Agostino RB Sr, Schwartz JS, Gibbons R, Shero ST, Greenland P, Smith SC Jr, Lackland DT, Sorlie P, Levy D, Stone NJ, Wilson PW. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013; doi: 10.1016/j.jacc.2013.11.005. [Epub ahead of print].
- Conroy RM, Pyörälä K, Fitzgerald AP, Sans S, Menotti A, De Backer G, De Bacquer D, Ducimetiere P, Jousilahti P, Keil U, Njolstad I, Oganov RG, Thomsen T, Tunstall-Pedoe H, Tverdal A, Wedel H, Whincup P, Wilhelmsen L, Graham IM, Group Sp. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003; **24**:987–1003.
- Firmann M, Mayor V, Vidal PM, Bochud M, Pecoud A, Hayoz D, Paccaud F, Preisig M, Song KS, Yuan X, Danoff TM, Stirnadel HA, Waterworth D, Mooser V, Waeber G, Vollenweider P. The CoLaus study: a population-based study to investigate the epidemiology and genetic determinants of cardiovascular risk factors and metabolic syndrome. *BMC Cardiovasc Disord* 2008; **8**:6.
- Cooney MT, Dudina AL, Graham IM. Value and limitations of existing scores for the assessment of cardiovascular risk: a review for clinicians. *J Am Coll Cardiol* 2009; **54**: 1209–1227.
- Farzadfar F, Finucane MM, Danaei G, Pelizzari PM, Cowan MJ, Paciorek CJ, Singh GM, Lin JK, Stevens GA, Riley LM, Ezzati M. Global Burden of Metabolic Risk Factors of Chronic Diseases Collaborating G. National, regional, and global trends in serum total cholesterol since 1980: systematic analysis of health examination surveys and epidemiological studies with 321 country-years and 3.0 million participants. *Lancet* 2011; **377**:578–586.