

Review article

Banded together: a review of the factors favouring group living in a social carnivore, the banded mongoose *Mungos mungo* (Carnivora: Herpestidae)

Markus Gusset

Department of Animal Behaviour, Institute of Zoology,
University of Zurich, 8057 Zurich, Switzerland,
e-mail: mgusset@bluewin.ch

Keywords: carnivores; egalitarianism; group living; mongooses; sociality.

As most species of the order Carnivora are solitary (Gittleman 1989), there is ongoing debate over the evolution and maintenance of sociality in carnivores. It has been hypothesised that five factors reduce the costs and/or increase the benefits of tolerating conspecifics (Creel and Macdonald 1995): (1) high abundance and renewability of prey; (2) constraints on dispersal and benefits of philopatry; (3) improved acquisition and defence of resources; (4) enhanced anti-predatory defence; and (5) alloparental care.

Mainly because of differences in feeding ecology, it has been speculated that selective forces behind the evolution and maintenance of sociality in mongooses (Herpestidae) may differ from those in carnivores in general (Rood 1975, 1986, Gorman 1979, Waser 1981, Waser and Waser 1985). Characteristically, most social mongooses are insectivorous, small and diurnal and live in open habitats (Gorman 1979, Rood 1986), which is in accordance with a recent analysis of genetic and behavioural data on the origin of sociality in mongooses (Veron et al. 2004).

Here, to shed further light on the factors favouring group living in carnivores, the five hypotheses outlined above are reviewed in relation to one such insectivorous, small, diurnal social mongoose living in open habitats, the banded mongoose *Mungos mungo* (Gmelin 1788).

Hypothesis 1: high abundance and renewability of prey When foraging, banded mongooses scatter and mainly feed on insects and other invertebrates, supplementing their diet with small vertebrates (Neal 1970, Rood 1975, 1982, Hiscocks and Perrin 1991). Both the abundance and renewal rate of their main prey are high (Waser 1981). High abundance and renewability of insects, together with relatively low competition in catching them, were probably a precondition for the evolution of group living in mongooses (Rood 1975, 1986, Gorman 1979, Waser 1981, Waser and Waser 1985).

Hypothesis 2: constraints on dispersal and benefits of philopatry Dispersal involves substantial costs in

banded mongooses (Cant et al. 2001), thus posing considerable constraints on leaving the natal pack. In line with this, banded mongooses exhibit low variance in within-group relatedness (Waldick et al. 2003), with no apparent avoidance or negative consequences of inbreeding (Gilchrist et al. 2004, Gilchrist 2006a). Despite a loose linear dominance hierarchy (without apparent effects on a variety of demographic and social variables) within a pack, there is no evidence of effective pre- or post-parturition reproductive suppression of subordinates in either sex (Cant 2000, De Luca and Ginsberg 2001, Gilchrist et al. 2004). As a result, there is a good chance for philopatric banded mongooses to reproduce in their natal pack.

Hypothesis 3: improved acquisition and defence of resources Banded mongooses cooperatively attack prey, take prey away from other predators and vocally attract other pack members to rich food patches (Simpson 1964, Rood 1975, Messeri et al. 1987). Aggressive territorial conflicts between banded mongoose packs are common, particularly during oestrus periods, with larger packs dominating smaller ones (Rood 1975, 1986, Cant et al. 2002, Gilchrist and Otali 2002, Müller and Manser 2007). Individuals in larger packs might be more likely to engage in extra-group copulations observed during such encounters (Rood 1975, Cant et al. 2002, Gilchrist et al. 2004).

Hypothesis 4: enhanced anti-predatory defence Banded mongooses cooperatively harass and even kill predators, bunch together around pups when disturbed in the open, and rescue other pack members from predators (Simpson 1964, Neal 1970, Rood 1975, 1982, 1983, 1986). Furthermore, members guard the pack through vigilance, emit alarm calls when danger threatens, and rush to help individuals calling in distress (Neal 1970, Rood 1975, Messeri et al. 1987). Group living in mongooses probably evolved as an anti-predatory response by small, diurnal animals living in open habitats (Rood 1975, 1986, Gorman 1979).

Hypothesis 5: alloparental care Within a banded mongoose pack, several females come into oestrus and subsequently produce their litters synchronously (Neal 1970, Rood 1974, 1975, 1986, Waser et al. 1995, Cant 2000, Gilchrist et al. 2004, Gilchrist 2006a,b). Lactating females apparently nurse any of the pups without discrimination and pack members cooperate in raising the pups around the communal den (Neal 1970, Rood 1974, 1975, 1982, 1986, Viljoen 1980, Waser et al. 1995, Cant 2003). Once the pups start to accompany the adults on their daily foraging trips, pack members “escort” and help provision the pups in a remarkably stable associa-

tion (Gilchrist 2004, Hodge 2005, Bell 2007, Gilchrist and Russell 2007).

Banded mongooses are a rare example of an egalitarian mammalian society, in that they both lack a distinct dominance hierarchy and have an apparently low reproductive skew. As a consequence of egalitarianism, an individual's chance of successful reproduction on reaching adulthood is good in its natal pack, especially as there is no apparent avoidance and negative consequence of inbreeding (Gilchrist et al. 2004, Gilchrist 2006a). Egalitarianism might thus be regarded as an incentive to stay that, similar to high abundance and renewability of prey, reduces an individual's costs of tolerating conspecifics and at the same time allows it to enjoy the equitably distributed benefits of living in a group (i.e., improved acquisition and defence of resources, enhanced anti-predatory defence, and alloparental care).

Therefore, increased (immediate and often mutualistic) direct and indirect fitness benefits accruing to non-dispersers (resulting from low variance in within-group relatedness; Waldick et al. 2003), together with considerable constraints on dispersal (Cant et al. 2001), are likely to be an additional factor favouring group living in banded mongooses as a plural breeder with communal care. Among social carnivores (Packer et al. 2001), the rare combination of plural breeding and low reproductive skew is more prevalent in species with relatively low costs of reproduction, which holds true for banded mongooses (Gittleman 1989, Creel and Creel 1991, Creel and Macdonald 1995).

In conclusion, it is remarkable that all factors proposed to reduce the costs and/or increase the benefits of tolerating conspecifics, and thus to favour group living, in carnivores (Creel and Macdonald 1995) are found to act in concert on a single species, the banded mongoose.

Acknowledgements

Financial support for this work was provided through a Swiss National Science Foundation grant to M. Manser. I am grateful to N. Burgener and two anonymous referees for helpful comments.

References

- Bell, M.B.V. 2007. Cooperative begging in banded mongoose pups. *Curr. Biol.* 17: 717–721.
- Cant, M.A. 2000. Social control of reproduction in banded mongooses. *Anim. Behav.* 59: 147–158.
- Cant, M.A. 2003. Patterns of helping effort in co-operatively breeding banded mongooses (*Mungos mungo*). *J. Zool.* 259: 115–121.
- Cant, M.A., E. Otali and F. Mwanguhya. 2001. Eviction and dispersal in co-operatively breeding banded mongooses (*Mungos mungo*). *J. Zool.* 254: 155–162.
- Cant, M.A., E. Otali and F. Mwanguhya. 2002. Fighting and mating between groups of a cooperatively breeding mammal, the banded mongoose. *Ethology* 108: 541–555.
- Creel, S.R. and N.M. Creel. 1991. Energetics, reproductive suppression and obligate communal breeding in carnivores. *Behav. Ecol. Sociobiol.* 28: 263–270.
- Creel, S. and D. Macdonald. 1995. Sociality, group size, and reproductive suppression among carnivores. *Adv. Stud. Behav.* 24: 203–257.
- De Luca, D.W. and J.R. Ginsberg. 2001. Dominance, reproduction and survival in banded mongooses: towards an egalitarian social system? *Anim. Behav.* 61: 17–30.
- Gilchrist, J.S. 2004. Pup escorting in the communal breeding banded mongoose: behaviour, benefits, and maintenance. *Behav. Ecol.* 15: 952–960.
- Gilchrist, J.S. 2006a. Reproductive success in a low skew, communal breeding mammal: the banded mongoose, *Mungos mungo*. *Behav. Ecol. Sociobiol.* 60: 854–863.
- Gilchrist, J.S. 2006b. Female eviction, abortion, and infanticide in banded mongooses (*Mungos mungo*): implications for social control of reproduction and synchronized parturition. *Behav. Ecol.* 17: 664–669.
- Gilchrist, J.S. and E. Otali. 2002. The effects of refuse-feeding on home-range use, group size, and intergroup encounters in the banded mongoose. *Can. J. Zool.* 80: 1795–1802.
- Gilchrist, J.S. and A.F. Russell. 2007. Who cares? Individual contributions to pup care by breeders vs. non-breeders in the cooperatively breeding banded mongoose (*Mungos mungo*). *Behav. Ecol. Sociobiol.* 61: 1053–1060.
- Gilchrist, J.S., E. Otali and F. Mwanguhya. 2004. Why breed communally? Factors affecting fecundity in a communal breeding mammal: the banded mongoose (*Mungos mungo*). *Behav. Ecol. Sociobiol.* 57: 119–131.
- Gittleman, J.L. 1989. Carnivore group living: comparative trends. In: (J.L. Gittleman, ed.) *Carnivore behavior, ecology, and evolution*. Cornell University Press, Ithaca. pp. 183–207.
- Gorman, M.L. 1979. Dispersion and foraging of the small Indian mongoose, *Herpestes auropunctatus* (Carnivora: Viverridae) relative to the evolution of social viverrids. *J. Zool.* 187: 65–73.
- Hiscocks, K. and M.R. Perrin. 1991. A dietary comparison between two sympatric viverrids, *Helogale parvula* (Sundevall, 1846) and *Mungos mungo* (Gmelin, 1788). *J. Afr. Zool.* 105: 307–312.
- Hodge, S.J. 2005. Helpers benefit offspring in both the short and long-term in the cooperatively breeding banded mongoose. *Proc. R. Soc. B Biol. Sci.* 272: 2479–2484.
- Messeri, P., E. Masi, R. Piazza and F. Dessi-Fulgheri. 1987. A study of the vocal repertoire of the banded mongoose, *Mungos mungo* (Gmelin). *Ital. J. Zool.* 22 (Suppl.): 341–373.
- Müller, C.A. and M.B. Manser. 2007. 'Nasty neighbours' rather than 'dear enemies' in a social carnivore. *Proc. R. Soc. B Biol. Sci.* 274: 959–965.
- Neal, E. 1970. The banded mongoose, *Mungos mungo* Gmelin. *E. Afr. Wildl. J.* 8: 63–71.
- Packer, C., A.E. Pusey and L.E. Eberly. 2001. Egalitarianism in female African lions. *Science* 293: 690–693.
- Rood, J.P. 1974. Banded mongoose males guard young. *Nature* 248: 176.
- Rood, J.P. 1975. Population dynamics and food habits of the banded mongoose. *E. Afr. Wildl. J.* 13: 89–111.
- Rood, J.P. 1982. Ecology and social organization of the banded and dwarf mongoose. *Natl. Geog. Soc. Res. Rep.* 14: 571–576.
- Rood, J.P. 1983. Banded mongoose rescues pack member from eagle. *Anim. Behav.* 31: 1261–1262.
- Rood, J.P. 1986. Ecology and social evolution in the mongooses. In: (D.I. Rubenstein and R.W. Wrangham, eds.) *Ecological aspects of social evolution*. Princeton University Press, Princeton. pp. 131–152.
- Simpson, C.D. 1964. Notes on the banded mongoose, *Mungos mungo* (Gmelin). *Arnoldia* 1: 1–8.
- Veron, G., M. Colyn, A.E. Dunham, P. Taylor and P. Gaubert. 2004. Molecular systematics and origin of sociality in mongooses (Herpestidae, Carnivora). *Mol. Phylogenet. Evol.* 30: 582–598.
- Viljoen, S. 1980. Early postnatal development, parental care and

- interaction in the banded mongoose *Mungos mungo*. S. Afr. J. Zool. 15: 119–120.
- Waldick, R.C., P. Johnson and J. Pemberton. 2003. Identification and characterization of 14 polymorphic microsatellite loci for a member of the Herpestidae (*Mungos mungo*). Mol. Ecol. Notes 3: 236–238.
- Waser, P.M. 1981. Sociality or territorial defense? The influence of resource renewal. Behav. Ecol. Sociobiol. 8: 231–237.
- Waser, P.M. and M.S. Waser. 1985. *Ichneumia albicauda* and the evolution of viverrid gregariousness. Z. Tierpsychol. 68: 137–151.
- Waser P.M., L.F. Elliott, N.M. Creel and S.R. Creel. 1995. Habitat variation and mongoose demography. In: (A.R.E. Sinclair and P. Arcese, eds.) Serengeti II: dynamics, management, and conservation of an ecosystem. University of Chicago Press, Chicago. pp. 421–447.