

High-energy radiation and particles in the environments of young stellar objects

Manuel Güdel

Institute of Astronomy, ETH Zurich, 8093 Zurich, Switzerland
email: guedel@astro.phys.ethz.ch

Abstract. High-energy radiation and particles profoundly affect circumstellar disk gas and solids. We discuss stellar high-energy sources and summarize their effects on circumstellar disks.

Keywords. Young stellar objects, X-rays, high-energy particles, ionization

1. Circumstellar disks subject to high-energy radiation and particles

Stellar X-rays may heat circumstellar disks to > 1000 K at AU distances, as suggested by, e.g. H_2 $2.12\mu\text{m}$ (Bary *et al.* 2003) and CO fundamental+overtone emission (Najita *et al.* 2003). X-rays may also ionize disks more efficiently than cosmic rays at similar distances, thus driving accretion through the magnetorotational instability (Glassgold *et al.* 1997). They may further dominate disk photoevaporation in the 10–40 AU range, resulting in mass loss rates of order $10^{-9} M_{\odot} \text{yr}^{-1}$ (Ercolano *et al.* 2009).

What high-energy photons do reach the disk surface at all? Average neutral gas column densities around classical T Tauri stars (CTTS, $N_{\text{H}} \approx 10^{21} - 10^{22} \text{cm}^{-2}$, Güdel *et al.* 2007a) typically exceed those of weak-lined T Tauri stars, suggesting excess gas located relatively close to the star. This gas easily absorbs EUV radiation, as do disk winds (Hollenbach & Gorti 2009), questioning the role of EUV radiation in CTTS disks. Absorption may be even more severe in some strong accretors where accretion streams seem to absorb all X-rays below $\approx 1 - 2$ keV (Güdel *et al.* 2007b).

Increasing X-ray absorption with increasing disk inclination provides, nevertheless, direct evidence for disk ionization (Kastner *et al.* 2005). Further support comes from fluorescent 6.4 keV $K\alpha$ emission from “cold” iron at the disk surface during strong flares (Imanishi *et al.* 2001). Fluorescence may reach extremely high levels in protostars even outside any obvious flaring (Skinner *et al.* 2007).

The [Ne II] $12.81\mu\text{m}$ line, frequently detected in *Spitzer* spectra (Pascucci *et al.* 2007), may be another diagnostic for X-ray/EUV disk irradiation (Glassgold *et al.* 2007). A statistically significant correlation with the stellar X-ray luminosity is, however, dominated by scatter (Figure 1a); obviously, a number of further parameters (e.g., disk properties) are relevant. For example, the presence of jets leads to a large increase in [Ne II] luminosity (Fig. 1b) as also shown in spatially resolved observations (van Boekel *et al.* 2009).

Glauser *et al.* (2009) found dust crystallinity to anti-correlate with the central star’s X-ray luminosity. Although X-rays carry insufficient momentum to induce lattice displacements, energetic (several keV) particles in the stellar wind are held responsible for amorphizing circumstellar dust.

2. What high-energy sources?

Apart from “traditional” magnetic coronae, additional high-energy sources have been identified. While accretion “suppresses” coronal X-rays by a factor of ≈ 2 (Telleschi *et al.*




Figure 1. *Left (a):* [Ne II] luminosity vs. X-ray luminosity for CTTS; jet sources are shown by diamonds, disks without jets by circles (filled: detections; open: upper limits). *Right (b):* Kaplan-Meier estimator for the [Ne II] luminosity of CTTS disks without jets, CTTS jet sources, and transition disks. From Güdel *et al.* (2009).

2007), it adds a “soft excess” at cool (coronal) temperatures (Güdel & Telleschi 2007), perhaps due to an interaction between accretion streams and the corona. High densities inferred from X-ray line triplets (Kastner *et al.* 2002) have successfully been modeled in terms of shocks at the footpoints of accretion streams (Günther *et al.* 2007). CTTS jets also emit X-rays close to the star (Güdel *et al.* 2007b). Their “lamp-post” arrangement may provide ideal illumination of disk surfaces, avoiding absorption by accretion streams.

Large flares are of interest as they produce hard X-rays and may eject more energetic particles. Disks may thus be ionized more efficiently to deeper levels (Ilgner & Nelson 2006). However, as the energetics of X-ray emission appear to be dominated by the large population of *small* flares (Audard *et al.* 2000), hard X-rays should be continuously present, adding as yet unrecognized ionization power to the circumstellar environment.

References

- Audard, M., Güdel, M., Drake, J. J., & Kashyap, V. L. 2000, *ApJ* 541, 396
 Bary, J. S., Weintraub, D. A., & Kastner, J. H. 2003, *ApJ* 586, 1136
 Ercolano, B., Clarke, C. J., & Drake, J. J. 2009, *ApJ* 699, 1639
 Glassgold, A. E., Najita, J. R., & Igea, J. 1997, *ApJ* 480, 344
 Glassgold, A. E., Najita, J. R., & Igea, J. 2007, *ApJ* 656, 515
 Glauser, A. *et al.* 2009, *A&A*, in press
 Güdel, M. & Telleschi, A. 2007, *A&A* 474, L25
 Güdel, M. *et al.* 2007a, *A&A* 468, 353
 Güdel, M., *et al.* 2007b, *A&A* 468, 515
 Güdel *et al.* 2009, *A&A*, submitted
 Günther, H. M., Schmitt, J. H. M. M., Robrade, J., & Liefke, C. 2007, *A&A*, 466, 1111
 Hollenbach, D. & Gorti, U. 2009, *ApJ*, 703, 1203
 Ilgner, M. & Nelson, R. P. 2006, *A&A* 455, 731
 Imanishi, K., Koyama, K., & Tsuboi, Y. 2001, *ApJ* 557, 747
 Kastner, J. H. *et al.* 2002, *ApJ* 567, 434
 Kastner, J. H. *et al.* 2005, *ApJS* 160, 511
 Najita, J., Carr, J. S., & Mathieu, R. D. 2003, *ApJ*, 589, 931
 Pascucci, I. *et al.* 2007, *ApJ*, 663, 383
 Skinner, S. L., Simmons, A. E., Audard, M., & Güdel, M. 2007, *ApJ* 658, 1144
 Telleschi, A., Güdel, M., Briggs, K. R., Audard, M., & Palla, F. 2007, *A&A* 468, 425
 van Boekel, R., Güdel, M., Henning, Th., Lahuis, F., & Pantin, E. 2009, *A&A*, 497, 137