

- untwisting behavior in patients with dilated cardiomyopathy: analysis by two-dimensional speckle tracking. *Heart* 2008;**95**:290–296.
32. van Dalen BM, Soliman OI, Vletter WB, ten Cate FJ, Geleijnse ML. Age-related changes in the biomechanics of left ventricular twist measured by speckle tracking echocardiography. *Am J Physiol Heart Circ Physiol* 2008;**295**:H1705–H1711.
  33. Ng AC, Tran dT, Newman M, Allman C, Vidaic J, Leung DY. Comparison of left ventricular dyssynchrony by two-dimensional speckle tracking versus tissue Doppler imaging in patients with non-ST-elevation myocardial infarction and preserved left ventricular systolic function. *Am J Cardiol* 2008;**102**:1146–1150.
  34. Zhang Q, Fung JW, Yip GW, Chan JY, Lee AP, Lam YY, Wu LW, Wu EB, Yu CM. Improvement of left ventricular myocardial short-axis, but not long-axis function or torsion after cardiac resynchronisation therapy: an assessment by two-dimensional speckle tracking. *Heart* 2008;**94**:1464–1471.
  35. Nishikage T, Nakai H, Mor-Avi V, Lang RM, Salgo IS, Settlemier SH, Husson S, Takeuchi M. Quantitative assessment of left ventricular volume and ejection fraction using two-dimensional speckle tracking echocardiography. *Eur J Echocardiogr* 2009;**10**:82–88.
  36. Okamoto K, Takeuchi M, Nakai H, Nishikage T, Salgo IS, Husson S, Otsuji Y, Lang RM. Effects of aging on left atrial function assessed by 2D speckle tracking echocardiography. *J Am Soc Echocardiogr* 2009;**22**:70–75.
  37. Jacobs LD, Salgo IS, Goonewardena S, Weinert L, Coon P, Bardo D, Gerard O, Allain P, Zamorano JL, de Isla LP, Mor-Avi V, Lang RM. Rapid online quantification of left ventricular volume from real-time three-dimensional echocardiographic data. *Eur Heart J* 2006;**27**:460–468.
  38. Meunier J. Tissue motion assessment from 3D echographic speckle tracking. *Phys Med Biol* 1998;**43**:1241–1254.
  39. Chen X, Xie H, Erkamp R, Kim K, Jia C, Rubin JM, O'Donnell M. 3-D correlation-based speckle tracking. *Ultrason Imaging* 2005;**27**:21–36.
  40. de Isla LP, Vivas D, Zamorano JL. Three-dimensional speckle tracking. *Current Cardiovasc Imaging Reports* 2008;**1**:25–29.
  41. Mor-Avi V, Sugeng L, Weinert L, MacEneaney P, Caiani EG, Koch R, Salgo IS, Lang RM. Fast measurement of left ventricular mass with real-time three-dimensional echocardiography: comparison with magnetic resonance imaging. *Circulation* 2004;**110**:1814–1818.
  42. Lang RM, Mor-Avi V, Sugeng L, Nieman PS, Sahn DJ. Three-dimensional echocardiography: the benefits of the additional dimension. *J Am Coll Cardiol* 2006;**48**:2053–2069.
  43. Corsi C, Lang RM, Veronesi F, Weinert L, Caiani EG, MacEneaney P, Lamberti C, Mor-Avi V. Volumetric quantification of global and regional left ventricular function from real-time three-dimensional echocardiographic images. *Circulation* 2005;**112**:1161–1170.

## CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehp173

Online publish-ahead-of-print 19 May 2009

### Ischaemic mitral regurgitation

Raban V. Jeger\* and Hans Peter Brunner-La Rocca

Department of Cardiology, University Hospital Basel, Petersgraben 4, Basel, Switzerland

\*Corresponding author. Tel: +41 61 265 2525, Fax: +41 61 265 4598, Email: rjeger@uhbs.ch

A 79-year-old woman with a history of treated hypertension and permanent atrial fibrillation was referred because of increasing shortness of breath at minimal exercise. Blood pressure was 110/49 mmHg, heart rate 80 b.p.m. (peak exercise rate 146 b.p.m.), and brain natriuretic peptide 86 pg/mL. Myocardial perfusion scan [Panel A: stress (upper rows) and resting (lower rows) single photon emission computed tomography images (left: short-axis views, right: long-axis views)], spirometry, and chest computer tomography were unremarkable. Resting transthoracic echocardiography revealed slight concentric hypertrophy of both ventricles, no wall-motion abnormalities, diastolic left-ventricular dysfunction grade 2, no valvular disease, and moderate pulmonary hypertension. Resting right heart catheter showed mildly elevated filling pressures and only mild pulmonary hypertension [Panel B: resting right heart catheter with pulmonary capillary wedge pressure (PCWP) mean 14 mmHg, v-wave 18 mmHg (arrow; caliper 40 mmHg)]. However, with minimal exercise (handgrip), PCWP increased significantly with marked v-waves, suggestive of exercise-induced mitral regurgitation [Panel C: exercise right heart catheter (hand grip): PCWP mean 30 mmHg, v-wave 46 mmHg (arrow; caliper 100 mmHg)]. Subsequent coronary angiography showed a severe stenosis of the left circumflex artery [Panel D: coronary angiography (ap view): high-grade stenosis of the left circumflex artery (arrow)] that was treated by percutaneous coronary intervention including stent implantation [Panel E: coronary angiography (ap view): left circumflex artery after implantation of a bare-metal stent (arrow)]. After the intervention, the symptoms disappeared. Two months later, the patient still does not report any shortness of breath. In conclusion, shortness of breath may be caused by various pathologies related to myocardial ischaemia.

