

Nephrol Dial Transplant (2006) 21: 859–864
doi:10.1093/ndt/gfk054
Advance Access publication 23 January 2006



Editorial Review

Novel anti-inflammatory drugs in hypertension

Matthias Hermann and Frank Ruschitzka

Cardiovascular Center, Cardiology, University Hospital Zurich, Switzerland

Keywords: hypertension; inflammation; cyclooxygenase; cardiovascular risk; NSAID; coxib

‘Atherosclerosis is a chronic inflammatory disease induced by cholesterol ...’

Rudolf Virchow (1821–1902)

As suggested by Virchow more than a century ago, inflammation plays a pivotal role in atherogenesis and potentially also in the pathogenesis of hypertension and its sequelae. As such, the current review focuses on the role of inflammation in hypertension and emerging therapeutical approaches.

Inflammation and cardiovascular risk

Clinical data

Over the last years chronic low-grade inflammation has emerged as an important new cardiovascular risk factor. Inflammatory responses within the vasculature might release pro-inflammatory cytokines that increase levels of C-reactive protein (CRP). First evidence for a role of inflammation in the development of atherosclerosis comes from studies demonstrating its prognostic value in unstable angina [1]. The inflammation hypothesis was further supported by data from observational studies demonstrating that CRP is a predictor of first cardiovascular events and might be an even stronger prognostic factor than low-density lipoprotein (LDL)-Cholesterol for coronary artery disease [2,3]. CRP appears to be a stable analyte over time and has been subject to numerous recent studies. Thus, on the basis of several studies, the Centers for Disease Control and Prevention and the American Heart Association recommended using highly sensitive (hs)-CRP and implicated the relative risk categories (low, average, high), corresponding

to approximate tertiles of values (<1.0, 1.0 to 3.0 and >3.0 mg/l, respectively) for individual cardiovascular risk assessment, in particular for those patients presenting with an intermediate risk [4]. However, a single inflammatory marker may or may not estimate all aspects of the underlying inflammatory processes, especially as they may affect cardiovascular disease (CVD) risk, and different markers may differ in their specificity for CVD. Indeed, there are several additional markers with a potential role as useful predictors of cardiovascular risk, such as serum amyloid-A, cytokines (e.g. interleukin (IL)-6), acute phase reactants, adhesion molecules and fibrinogen.

Interestingly, a recent publication demonstrated that significant gender and race differences in CRP levels exist which contribute to differences in cardiovascular outcome [5]. CRP plasma levels were assessed in 2749 white and black subjects aged 30–65 years and participating in the Dallas Heart Study. Black subjects had higher CRP levels than white subjects and women had higher CRP levels than men, suggesting that overall recommendations for CRP cut-off levels for risk assessment should be adjusted for race and gender. The clinical relevance of CRP as a significant predictor of coronary artery disease (CAD) was further challenged by Danesh *et al.* [6], demonstrating that CRP was a weaker marker than traditional risk factors like total cholesterol or smoking in the Reykjavik prospective study. However, in the study population, total plasma cholesterol levels were higher and CRP levels lower than those reported in the US population.

Experimental data

Experimental data demonstrate that CRP may exert effects on the vasculature. Indeed, CRP induces ICAM-1, VCAM-1 and MCP-1 expression in endothelial cells [7,8], upregulates angiotensin II type 1 receptors in human vascular smooth muscle cells [9] and CRP opsonization of low-density lipoprotein (LDL) mediates LDL uptake by macrophages [10]. CRP inhibits the expression of NO synthase via translational and post-translational mechanisms, resulting in reduced

Correspondence and offprint requests to: Matthias Hermann, Cardiovascular Center, Cardiology, University Hospital Zurich, Switzerland. Email: mhermann@gmx.de

NO bioavailability, and stimulates the release of ET-1, IL-6 and vasoconstrictive peptides in human endothelial cells [11,12]. Interestingly, CRP levels correlate to endothelial dysfunction in patients with angiographically normal coronary arteries, established CAD and acute coronary syndromes [13–15]. A recent publication demonstrated specific receptors for CRP on human aortic endothelial cells [17]. After binding to FC gamma receptors I and II, CD64 and CD32 respectively, CRP reduces eNOS activity and prostacyclin levels and increases interleukin-8, ICAM-1 and VCAM-1. These pro-inflammatory effects were prevented with specific antibodies to CD32 and CD64. However, CRP elicits potential vasoprotective effects such as endothelium-independent vasorelaxing effects in human vessels *in vitro* [16].

In addition, opposing effects of native and modified CRP have been reported. Indeed, native CRP promotes while modified CRP attenuates the development of atherosclerotic lesions in ApoE knockout mice [18]. Verma *et al.* [12] showed that CRP inhibited both basal and vascular endothelial growth factor (VEGF)-stimulated angiogenesis, which would in turn inhibit plaque neovascularization. Furthermore, recent publications suggest, that CRP independent mechanisms like azide and lipopolysaccharide, but not CRP itself, might cause some of the observed effects on endothelial cells [76,77].

Inflammation and endothelial function

Endothelial function has evolved as a surrogate for the development and progression of atherosclerotic disease. Indeed, impaired endothelial function, characterized by reduced bioavailability of NO, has prognostic value for future cardiovascular events [19] and has been demonstrated for all established and also for newer cardiovascular risk factors like inflammation.

Acute and chronic systemic inflammation attenuates endothelial function. Experimental inflammation after infusion of salmonella thyphi vaccine induces endothelial dysfunction within 8 h in healthy volunteers [20]. Evidence for a potential role of chronic inflammation in atherogenesis comes from studies in patients with rheumatoid arthritis (RA). These patients show an increase in both cardiovascular morbidity and mortality [21]. Furthermore, striking similarities exist between the paradigm of inflammation in the pathogenesis of both atherosclerotic vascular disease and RA [22,23]. The shared features include involvement of cytokines, such as TNF- α and IL-6, raised concentrations of CRP, fibrinogen and amyloid-A, increased local expression of adhesion molecules and endothelin-1 [24]. These similarities suggest that inflammatory mechanisms also involve the vessel wall and facilitate the development of atherosclerotic lesions. Indeed, we recently demonstrated early endothelial dysfunction in patients with RA [25] providing an explanation for the excess cardiovascular morbidity in RA patients in whom traditional cardiovascular risk factors are usually absent [26].

Hypertension and vascular inflammation

Chronic inflammation presents with activation of the cyclooxygenase (COX) system, increased production of ROS and increased synthesis of CRP and pro-inflammatory cytokines, such as TNF- α and IL-6. An increasing body of evidence suggests that low-grade inflammation and oxidative stress account in part for hypertension-induced endothelial dysfunction and that CRP levels are associated with future development of hypertension [27–29]. Similar results have been provided by Engstrom *et al.* [30], showing that increased levels of inflammation-sensitive plasma proteins (fibrinogen, alpha-1-antitrypsin, haptoglobin, ceruloplasmin and orosomucoid) are associated with an increased incidence of hypertension. Incident hypertension, the initiation of antihypertensive treatment, and self-reported systolic blood pressure of at least 140 mmHg or a diastolic blood pressure of at least 90 mmHg are increased in women with baseline CRP levels above 3.5 mg/l even after adjustment for cardiovascular risk factors [29]. Cross-sectional analysis of more than 15 000 women demonstrated a linear relationship between increasing blood pressure and increasing CRP levels [27]. During follow-up over 8 years both parameters were strong predictors for cardiovascular events, and the predictive values of CRP and elevated blood pressure in combination are additive. Interestingly, although low-dose aspirin has no effect on CRP levels, it has been shown to be effective in primary prevention in men. This effect, however, is greatest in persons with the highest levels of hs-CRP and declines in direct relation to CRP levels [31].

In untreated human hypertension, CRP levels have recently been found to be increasingly dependent on systolic blood pressure levels [32–35]. Most importantly, CRP increase is independently associated with other classical cardiovascular risk factors [32] and might be an independent risk factor for the development of hypertension after correction for age, sex, body mass index, family history of hypertension, fasting glycemia, sedentary behaviour, and alcohol consumption [34]. In addition, CRP is a strong risk factor for ischaemic stroke, independent of the severity of the underlying atherosclerotic disease [36,37]. A recent study investigated the relationship between IL-6, TNF- α , and hs-CRP with arterial stiffness in untreated hypertensive patients [38]. The authors demonstrated that hs-CRP, IL-6 and TNF- α are significantly related to pulse wave velocity, a marker of aortic stiffness, and augmentation index, a manifestation of wave reflection, in essential hypertension. These data suggest a pivotal role for inflammation in the development of vascular disease, hypertension in particular. However, initiation of inflammatory processes expands from hypertension towards the more complex metabolic syndrome, including pathologic glucose and lipid levels, visceral obesity and hypertension. Each component of the metabolic syndrome can induce vascular inflammation,

thus increasing cardiovascular risk. Therefore, a rationale for assessment of inflammation status and anti-inflammatory treatment in addition to anti-hypertensive treatment in atherosclerosis and hypertension is given and may increase clinical benefit.

Anti-inflammatory drugs and hypertension

Selective and non-selective COX-inhibitors

The impact of specific anti-inflammatory drugs, selective and non-selective COX-inhibitors (coxibs and NSAIDs), on cardiovascular homeostasis and the incidence of cardiovascular events have been discussed intensively over the last months.

Indeed, increased cardiovascular events have been observed with rofecoxib in the Vioxx Gastrointestinal Outcomes Research trial (VIGOR [39]) and Adenomatous Polyp Prevention on Vioxx (APPROVE [40]) trial as well as with celecoxib in the Adenoma Prevention with Celecoxib (APC) trial [41], pointing towards a potential class effect for coxibs. This 'class effect' for an increase in the risk of cardiovascular disease is thought to be related to preferential inhibition of prostacyclin over thromboxane and thus a tendency towards pro-thrombosis. However, data suggest differential effects of coxibs respect to cardiovascular risk. Thus, other mechanisms come into play and might counterbalance this pro-thrombotic effect. These include a differential impact on oxidative stress, inflammation markers, endothelial function, renal function and morphology and tissue factor expression and activity with potential beneficial effects for celecoxib but not for rofecoxib [42–46].

In addition to putative pro-thrombotic effects of these drugs, the impact on blood pressure levels and blood pressure control are important. Several studies revealed differences between commonly used drugs and within the group of selective COX-2 inhibitors, the coxibs. Rofecoxib is clearly associated with an increase in blood pressure and new onset of hypertension while celecoxib is comparable to other NSAIDs [47,48]. In patients with osteoarthritis, treatment with rofecoxib but not celecoxib or naproxen induced a significant increase in 24 h systolic BP. However, a destabilization of hypertension control occurred to some extent with all three drugs, while this phenomenon was seen more often in patients treated with rofecoxib than with the other therapies [49]. Similar data come from a recent meta-analysis of 19 randomized controlled trials involving coxibs and non-selective NSAIDs [50]. Selective COX-2 inhibitors were associated with blood pressure elevation compared with placebo and nonselective NSAIDs; however, there was a higher incidence of developing hypertension observed with rofecoxib compared with celecoxib.

Involved mechanisms might include a disturbance of sodium and water retention similar to that observed with classical non-selective NSAIDs. Early studies in healthy subjects showed that high doses of celecoxib,

rofecoxib and NSAIDs induce a slight decrease in water, sodium and potassium excretion, but had no impact on blood pressure levels [51,52]. In users of NSAIDs, development of peripheral oedema and increases in blood pressure are associated with the development of chronic heart failure [53]. Further data on differences between coxibs come from a large observational study on the incidence of chronic heart failure during COX-inhibition. In over 38 000 patients who were started on rofecoxib, celecoxib or non-selective NSAIDs, rofecoxib and NSAIDs showed an adjusted rate-ratio of 1.8 and 1.4 respectively, while no increased risk of admission for congestive heart failure was observed for celecoxib [54].

Interestingly, ACE-inhibitors, beta blockers and diuretics, and to a lesser extent angiotensin II receptor blockers and calcium channel blockers, rely on the synthesis of vasodilator prostaglandins to exert their effects [55,56]. This is of particular interest, since prostacyclin levels can be reduced not only with selective COX-2 inhibitors but also with higher doses of classical NSAIDs [57]. This might translate into clinical practice, as recent data suggest that rofecoxib appears to interfere with the anti-hypertensive effects of ACE-inhibitors and beta-blockers, but not of calcium channel blockers [47].

While anti-hypertensive treatment is known to improve endothelial function as a surrogate for atherosclerosis, clinical and experimental studies also demonstrate that anti-inflammatory treatment has beneficial effects on vascular function. In patients with severe CAD, treatment with celecoxib reduces plasma levels of hs-CRP and oxidized-LDL, as markers of low-grade chronic inflammation and oxidative stress. This effect was accompanied by improved endothelial function [42]. Similar results were obtained in patients with arterial hypertension who were treated with celecoxib. Endothelial function improved within 3 h and remained so during the treatment course [43]. A comparative study of celecoxib, rofecoxib and diclofenac on vascular function in salt-sensitive hypertensive rats showed that celecoxib, but not rofecoxib or diclofenac, improves endothelial function [44]. In addition, oxidative stress and inflammation markers were reduced by celecoxib alone. In a similar model, celecoxib improved proteinuria, renal and pre-glomerular vessel morphology and reduced the number of macrophages and CRP-mRNA in renal cortex of the hypertensive animals [45].

ACE-inhibitors and angiotensin receptor blockers

Several large randomized trials demonstrated a reduction in the incidence of recurrent cardiovascular events and mortality with ACE-inhibitor therapy [58–60]. In addition, ACE-inhibitors and angiotensin receptor blockers exert anti-inflammatory effects particularly by inhibiting pro-inflammatory effects of angiotensin II [61]. Olmesartan treatment significantly reduced serum levels of hs-CRP, TNF- α , IL-6, and monocyte chemoattractant protein-1 after 6 weeks of therapy in patients with

essential hypertension, suggesting that this anti-inflammatory action of angiotensin II receptor antagonists may contribute to their beneficial cardiovascular effects. The stimulation of cytokines by angiotensin II seems to be mediated via an activation of nuclear factor- κ B (NF- κ B) dependent pathway, which can be blocked, by losartan and candesartan [62,63].

Statins

Statin treatment significantly reduces the number of cardiovascular events, particularly in secondary prevention [64,65]. For better guidance of cholesterol levels to reduce cardiovascular risk with statins, monitoring of CRP might be helpful, because statins lower LDL cholesterol and CRP. The relationships between LDL cholesterol and CRP levels after treatment with atorvastatin or pravastatin and the risk of recurrent myocardial infarction or death from coronary causes were evaluated among 3745 patients with acute coronary syndromes [66]. Both statins lowered cholesterol levels, although atorvastatin was more likely than pravastatin to result in low levels of LDL cholesterol and CRP. Patients who had LDL cholesterol levels of less than 70 mg/dl and CRP levels of less than 1 mg/l after statin therapy had the lowest rate of recurrent events, demonstrating that patients who have low CRP levels after statin therapy have better clinical outcomes than those with higher CRP levels, regardless of the resultant level of LDL cholesterol. To further investigate the impact of statin treatment in the primary prevention of cardiovascular disease in patients with low LDL-cholesterol but elevated hs-CRP levels, the JUPITER trial has recently been initiated [67].

PPAR gamma activators

Other widely described drugs for patients with cardiovascular risk factors like hypertension, metabolic syndrome and type 2 diabetes, peroxisome proliferator-activated receptor (PPAR)-alpha and -gamma activators, seem to have anti-inflammatory properties. Indeed, PPAR-alpha activators like fenofibrate inhibit myocardial inflammation in angiotensin II-infused and salt-induced hypertensive rats [68,69]. In a similar model, angiotensin II-induced hypertensive rats received the PPAR-gamma activators pioglitazone or rosiglitazone. Both drugs reduced upregulated cell cycle proteins and pro-inflammatory mediators like NF- κ B, VCAM and PECAM and improved endothelial function, and attenuated blood pressure increase and expression of angiotensin II type 1 receptors [70]. The anti-inflammatory effects of pioglitazone seem to be independent of glucose control, as demonstrated recently in a prospective study in 192 patients with type 2 diabetes [71]. Therefore, in view of the evolving epidemic of the metabolic syndrome with arterial hypertension and hyperglycaemia, additional effects of drugs for the control of single components of the metabolic syndrome are important.

Cannabinoid receptor antagonist

This might hold also for a very new class of drugs interacting with the endocannabinoid system, the cannabinoid-1 receptor in particular. Originally developed to help smoking cessation, rimonabant has favourable effects on body weight, waist circumference, insulin resistance and dyslipidaemia [72]. In normotensive animals no effect on blood pressure levels is observed but in hypertensive animals, rimonabant lowers blood pressure and reduces cardiac contractility [73]. In addition, rimonabant has potent anti-inflammatory activity, as demonstrated in several animal models [74,75]. However, data from large-scale human studies show that effects on blood pressure are only mild and data on impact on inflammation markers are as yet unavailable.

Conclusion

Inflammation plays a pivotal role in the development and progression of atherosclerosis. Like other risk factors, arterial hypertension is associated with a pro-inflammatory status. However, the 'inflammation in atherosclerosis hypothesis' remains still to be challenged in prospective clinical trials. Indeed, unless anti-inflammatory treatment strategies provide benefit for patients, inflammation may thus face the fate of some other cardiovascular risk factors that, notwithstanding their potential pathophysiological relevance, are considered as lost in translation into clinical practice.

Conflict of interest statement. Dr Ruschitzka has served as a consultant or received speaker fees from Aventis, Bayer, Merck, Novartis and Pfizer.

References

1. Liuzzo G, Biasucci LM, Gallimore JR *et al.* The prognostic value of C-reactive protein and serum amyloid A protein in severe unstable angina. *N Engl J Med* 1994; 331: 417–424
2. Ridker PM, Hennekens CH, Buring JE *et al.* C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000; 342: 836–843
3. Ridker PM, Rifai N, Rose L *et al.* Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med* 2002; 347: 1557–1565
4. Pearson TA, Mensah GA, Alexander RW *et al.* Markers of inflammation and cardiovascular disease: application to clinical and public health practice: a statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 2003; 107: 499–511
5. Khera A, McGuire DK, Murphy SA *et al.* Race and gender differences in C-reactive protein levels. *J Am Coll Cardiol* 2005; 46: 464–469
6. Danesh J, Wheeler JG, Hirschfield GM *et al.* C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *N Engl J Med* 2004; 350: 1387–1397
7. Pasceri V, Willerson JT, Yeh ET. Direct proinflammatory effect of C-reactive protein on human endothelial cells. *Circulation* 2000; 102: 2165–2168

8. Pasceri V, Cheng JS, Willerson JT *et al.* Modulation of C-reactive protein-mediated monocyte chemoattractant protein-1 induction in human endothelial cells by anti-atherosclerosis drugs. *Circulation* 2001; 103: 2531–2534
9. Wang CH, Li SH, Weisel RD *et al.* C-reactive protein upregulates angiotensin type 1 receptors in vascular smooth muscle. *Circulation* 2003; 107: 1783–1790
10. Zwaka TP, Hombach V, Torzewski J. C-reactive protein-mediated low density lipoprotein uptake by macrophages: implications for atherosclerosis. *Circulation* 2001; 103: 1194–1197
11. Venugopal SK, Devaraj S, Yuhanna I *et al.* Demonstration that C-reactive protein decreases eNOS expression and bioactivity in human aortic endothelial cells. *Circulation* 2002; 106: 1439–1441
12. Verma S, Wang CH, Li SH *et al.* A self-fulfilling prophecy: C-reactive protein attenuates nitric oxide production and inhibits angiogenesis. *Circulation* 2002; 106: 913–919
13. Teragawa H, Fukuda Y, Matsuda K *et al.* Relation between C-reactive protein concentrations and coronary microvascular endothelial function. *Heart* 2004; 90: 750–754
14. Fichtlscherer S, Rosenberger G, Walter DH *et al.* Elevated C-reactive protein levels and impaired endothelial vasoreactivity in patients with coronary artery disease. *Circulation* 2000; 102: 1000–1006
15. Fichtlscherer S, Zeiher AM. Endothelial dysfunction in acute coronary syndromes: association with elevated C-reactive protein levels. *Ann Med* 2000; 32: 515–518
16. Sternik L, Samee S, Schaff HV *et al.* C-reactive protein relaxes human vessels in vitro. *Arterioscler Thromb Vasc Biol* 2002; 22: 1865–1868
17. Devaraj S, Du Clos TW, Jialal I. Binding and internalization of C-reactive protein by Fcγ receptors on human aortic endothelial cells mediates biological effects. *Arterioscler Thromb Vasc Biol* 2005; 25: 1359–1363
18. Schwedler SB, Amann K, Wernicke K *et al.* Native C-reactive protein increases whereas modified C-reactive protein reduces atherosclerosis in apolipoprotein E-knockout mice. *Circulation* 2005; 112: 1016–1023
19. Schachinger V, Britten MB, Zeiher AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation* 2000; 101: 1899–1906
20. Hingorani AD, Cross J, Kharbanda RK *et al.* Acute systemic inflammation impairs endothelium-dependent dilatation in humans. *Circulation* 2000; 102: 994–999
21. Mutru O, Laakso M, Isomaki H *et al.* Cardiovascular mortality in patients with rheumatoid arthritis. *Cardiology* 1989; 76: 71–77
22. Ross R. Atherosclerosis – an inflammatory disease. *N Engl J Med* 1999; 340: 115–126
23. Pasceri V, Yeh ET. A tale of two diseases: atherosclerosis and rheumatoid arthritis. *Circulation* 1999; 100: 2124–2126
24. van der Wal AC, Becker AE, van der Loos CM *et al.* Site of intimal rupture or erosion of thrombosed coronary atherosclerotic plaques is characterized by an inflammatory process irrespective of the dominant plaque morphology. *Circulation* 1994; 89: 36–44
25. Hurlimann D, Forster A, Noll G *et al.* Anti-tumor necrosis factor-α treatment improves endothelial function in patients with rheumatoid arthritis. *Circulation* 2002; 106: 2184–2187
26. Kharbanda RK, Walton B, Allen M *et al.* Prevention of inflammation-induced endothelial dysfunction: a novel vasculo-protective action of aspirin. *Circulation* 2002; 105: 2600–2604
27. Blake GJ, Rifai N, Buring JE *et al.* Blood pressure, C-reactive protein, and risk of future cardiovascular events. *Circulation* 2003; 108: 2993–2999
28. Cai H, Harrison DG. Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress. *Circ Res* 2000; 87: 840–844.
29. Sesso HD, Buring JE, Rifai N *et al.* C-reactive protein and the risk of developing hypertension. *JAMA* 2003; 290: 2945–2951.
30. Engstrom G, Janzon L, Berglund G *et al.* Blood pressure increase and incidence of hypertension in relation to inflammation-sensitive plasma proteins. *Arterioscler Thromb Vasc Biol* 2002; 22: 2054–2058
31. Ridker PM, Cushman M, Stampfer MJ *et al.* Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 1997; 336: 973–979
32. Schillaci G, Pirro M, Gemelli F *et al.* Increased C-reactive protein concentrations in never-treated hypertension: the role of systolic and pulse pressures. *J Hypertens* 2003; 21: 1841–1846.
33. Dodson PM, Shine B. Retinal vein occlusion: C-reactive protein and arterial hypertension. *Acta Ophthalmol (Copenh)* 1984; 62: 123–130
34. Bautista LE, Lopez-Jaramillo P, Vera LM *et al.* Is C-reactive protein an independent risk factor for essential hypertension? *J Hypertens* 2001; 19: 857–861
35. Choi H, Cho DH, Shin HH *et al.* Association of high sensitivity C-reactive protein with coronary heart disease prediction, but not with carotid atherosclerosis, in patients with hypertension. *Circ J* 2004; 68: 297–303
36. Curb JD, Abbott RD, Rodriguez BL *et al.* C-reactive protein and the future risk of thromboembolic stroke in healthy men. *Circulation* 2003; 107: 2016–2020
37. Cao JJ, Thach C, Manolio TA *et al.* C-reactive protein, carotid intima-media thickness, and incidence of ischemic stroke in the elderly: the Cardiovascular Health Study. *Circulation* 2003; 108: 166–170
38. Mahmud A, Feely J. Arterial stiffness is related to systemic inflammation in essential hypertension. *Hypertension* 2005; 46(5): 1118–1222
39. Bombardier C, Laine L, Reicin A *et al.* Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis. VIGOR Study Group. *N Engl J Med* 2000; 343: 1520–1528
40. Bresalier RS, Sandler RS, Quan H *et al.* Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *N Engl J Med* 2005; 352: 1092–1102
41. Solomon SD, McMurray JJ, Pfeffer MA *et al.* Cardiovascular risk associated with celecoxib in a clinical trial for colorectal adenoma prevention. *N Engl J Med* 2005; 352: 1071–1080
42. Chenevard R, Hurlimann D, Bechir M *et al.* Selective COX-2 inhibition improves endothelial function in coronary artery disease. *Circulation* 2003; 107: 405–409
43. Widlansky ME, Price DT, Gokce N *et al.* Short- and long-term COX-2 inhibition reverses endothelial dysfunction in patients with hypertension. *Hypertension* 2003; 42: 310–315
44. Hermann M, Camici G, Fratton A *et al.* Differential effects of selective cyclooxygenase-2 inhibitors on endothelial function in salt-induced hypertension. *Circulation* 2003; 108: 2308–2311
45. Hermann M, Shaw S, Kiss E *et al.* Selective COX-2 inhibitors and renal injury in salt-sensitive hypertension. *Hypertension* 2005; 45: 193–197
46. Steffel J, Hermann M, Greutert H *et al.* Celecoxib decreases endothelial tissue factor expression through inhibition of c-Jun terminal NH2 kinase phosphorylation. *Circulation* 2005; 111: 1685–1689
47. Whelton A, White WB, Bello AE *et al.* Effects of celecoxib and rofecoxib on blood pressure and edema in patients > or =65 years of age with systemic hypertension and osteoarthritis. *Am J Cardiol* 2002; 90: 959–963
48. White WB, Kent J, Taylor A *et al.* Effects of celecoxib on ambulatory blood pressure in hypertensive patients on ACE inhibitors. *Hypertension* 2002; 39: 929–934

49. Sowers JR, White WB, Pitt B *et al.* The Effects of cyclooxygenase-2 inhibitors and nonsteroidal anti-inflammatory therapy on 24-hour blood pressure in patients with hypertension, osteoarthritis, and type 2 diabetes mellitus. *Arch Intern Med* 2005; 165: 161–168
50. Aw TJ, Haas SJ, Liew D *et al.* Meta-analysis of cyclooxygenase-2 inhibitors and their effects on blood pressure. *Arch Intern Med* 2005; 165: 490–496
51. Rossat J, Maillard M, Nussberger J *et al.* Renal effects of selective cyclooxygenase-2 inhibition in normotensive salt-depleted subjects. *Clin Pharmacol Ther* 1999; 66: 76–84
52. Catella-Lawson F, McAdam B, Morrison BW *et al.* Effects of specific inhibition of cyclooxygenase-2 on sodium balance, hemodynamics, and vasoactive eicosanoids. *J Pharmacol Exp Ther* 1999; 289: 735–741
53. Feenstra J, Heerdink ER, Grobbee DE *et al.* Association of nonsteroidal anti-inflammatory drugs with first occurrence of heart failure and with relapsing heart failure: the Rotterdam Study. *Arch Intern Med* 2002; 162: 265–270
54. Mamdani M, Juurlink DN, Lee DS *et al.* Cyclo-oxygenase-2 inhibitors versus non-selective non-steroidal anti-inflammatory drugs and congestive heart failure outcomes in elderly patients: a population-based cohort study. *Lancet* 2004; 363: 1751–1756
55. Whelton A. Renal aspects of treatment with conventional nonsteroidal anti-inflammatory drugs versus cyclooxygenase-2-specific inhibitors. *Am J Med* 2001; 110 [Suppl 3A]: 33S–42S
56. Johnson AG, Nguyen TV, Day RO. Do nonsteroidal anti-inflammatory drugs affect blood pressure? A meta-analysis. *Ann Intern Med* 1994; 121: 289–300
57. Yamanari H, Nakamura K, Kakishita M *et al.* Effects of cyclooxygenase inhibition on endothelial function in hypertensive patients treated with angiotensin-converting enzyme inhibitors. *Clin Cardiol* 2004; 27: 523–527
58. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. Heart Outcomes Prevention Evaluation Study Investigators. *Lancet* 2000; 355: 253–259
59. Yusuf S, Sleight P, Pogue J *et al.* Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 2000; 342: 145–153
60. Fox KM. Efficacy of perindopril in reduction of cardiovascular events among patients with stable coronary artery disease: randomised, double-blind, placebo-controlled, multicentre trial (the EUROPA study). *Lancet* 2003; 362: 782–788
61. Fliser D, Buchholz K, Haller H. Antiinflammatory effects of angiotensin II subtype 1 receptor blockade in hypertensive patients with microinflammation. *Circulation* 2004; 110: 1103–1107
62. Kranzhofer R, Schmidt J, Pfeiffer CA *et al.* Angiotensin induces inflammatory activation of human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol.* 1999; 19: 1623–1629
63. Skurk T, van Harmelen V, Hauner H. Angiotensin II stimulates the release of interleukin-6 and interleukin-8 from cultured human adipocytes by activation of NF-kappaB. *Arterioscler Thromb Vasc Biol* 2004; 24: 1199–1203
64. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994; 344: 1383–1389
65. Ridker PM, Rifai N, Pfeffer MA *et al.* Inflammation, pravastatin, and the risk of coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events (CARE) Investigators. *Circulation* 1998; 98: 839–844
66. Ridker PM, Cannon CP, Morrow D *et al.* C-reactive protein levels and outcomes after statin therapy. *N Engl J Med* 2005; 352: 20–28
67. Ridker PM. Rosuvastatin in the primary prevention of cardiovascular disease among patients with low levels of low-density lipoprotein cholesterol and elevated high-sensitivity C-reactive protein: rationale and design of the JUPITER trial. *Circulation* 2003; 108: 2292–2297
68. Ogata T, Miyauchi T, Sakai S *et al.* Myocardial fibrosis and diastolic dysfunction in deoxycorticosterone acetate-salt hypertensive rats is ameliorated by the peroxisome proliferator-activated receptor-alpha activator fenofibrate, partly by suppressing inflammatory responses associated with the nuclear factor-kappa-B pathway. *J Am Coll Cardiol* 2004; 43: 1481–1488
69. Diep QN, Benkirane K, Amiri F *et al.* PPAR alpha activator fenofibrate inhibits myocardial inflammation and fibrosis in angiotensin II-infused rats. *J Mol Cell Cardiol* 2004; 36: 295–304
70. Diep QN, El Mabrouk M, Cohn JS *et al.* Structure, endothelial function, cell growth, and inflammation in blood vessels of angiotensin II-infused rats: role of peroxisome proliferator-activated receptor-gamma. *Circulation* 2002; 105: 2296–2302
71. Pfutzner A, Marx N, Lubben G *et al.* Improvement of cardiovascular risk markers by pioglitazone is independent from glycemic control: results from the pioneer study. *J Am Coll Cardiol* 2005; 45: 1925–1931
72. Van Gaal LF, Rissanen AM, Scheen AJ *et al.* Effects of the cannabinoid-1 receptor blocker rimonabant on weight reduction and cardiovascular risk factors in overweight patients: 1-year experience from the RIO-Europe study. *Lancet* 2005; 365: 1389–1397
73. Batkai S, Pacher P, Osei-Hyiaman D *et al.* Endocannabinoids acting at cannabinoid-1 receptors regulate cardiovascular function in hypertension. *Circulation* 2004; 110: 1996–2002
74. Croci T, Landi M, Galzin AM *et al.* Role of cannabinoid CB1 receptors and tumor necrosis factor-alpha in the gut and systemic anti-inflammatory activity of SR 141716 (rimonabant) in rodents. *Br J Pharmacol* 2003; 140: 115–122
75. Massa F, Marsicano G, Hermann H *et al.* The endogenous cannabinoid system protects against colonic inflammation. *J Clin Invest* 2004; 113: 1202–1209
76. Taylor KE, Giddings JC, van den Berg CW. C-reactive protein-induced *in vitro* endothelial cell activation is an artefact caused by azide and lipopolysaccharide. *Arterioscler Thromb Vasc Biol* 2005; 25(6): 1225–1230
77. Clapp BR, Hirschfield GM, Storry C *et al.* Inflammation and endothelial function: direct vascular effects of human C-reactive protein on nitric oxide bioavailability. *Circulation* 2005; 111(12): 1530–1536

Received for publication: 28.11.05

Accepted in revised form: 14.12.05