

Cite this article as: Zientara A, Genoni M, Schwegler I. Type B dissection reveals a rare anomaly of left cervical arch of Haughton D type. Eur J Cardiothorac Surg 2015;47:1109.

## Type B dissection reveals a rare anomaly of left cervical arch of Haughton D type

Alicja Zientara\*, Michele Genoni and Igor Schwegler

Department of Cardiac and Vascular Surgery, Triemli Hospital, Zürich, Switzerland


\* Corresponding author. Department of Cardiac Surgery, City Hospital Triemli, Birmensdorferstrasse 497, 8063 Zürich, Switzerland. Tel: +41-44-4661650; fax: +41-44-4662745; e-mail: alicja\_zientara@web.de (A. Zientara).

Received 1 May 2014; received in revised form 26 June 2014; accepted 27 June 2014


**Keywords:** Left cervical arch aneurysm • Vascular malformations • Thoracic aorta • Mediastinum

A 32-year-old woman with a history of lumbar-vertebral syndrome presented with acute back pain. The imaging showed a rare anomaly of a left cervical arch of Haughton D type complicated by

a dissection arising from the arch aneurysm (Fig. 1) to both iliac arteries (Fig. 2A and B).



**Figure 1:** Computed tomography; arrow points to the left subclavian artery arising from the arch aneurysm (\*greatest extent of 41 mm).



**Figure 2:** (A and B) Computed tomography reconstruction of the aortic arch; arrows show the course of the dissecting membrane, which starts from the aneurysm. LACC: left common carotid artery; RACC: right common carotid artery; TB: truncus brachiocephalicus; RSA: right subclavian artery.