

An Anosov action on the bundle of Weyl chambers

HANS-CHRISTOPH IM HOF

*Mathematisches Institut der Universität Basel,
Rheinsprung 21, CH-4051 Basel, Switzerland*

(Received 15 October 1984 and revised 1 March 1985)

Abstract. We introduce an Anosov action on the bundle of Weyl chambers of a riemannian symmetric space of non-compact type, which for rank one spaces coincides with the geodesic flow.

Introduction

The geodesic flow of a riemannian symmetric space of non-compact type is an Anosov flow if and only if the rank of the space is one. In this note we shall introduce an action, which for spaces of rank one coincides with the geodesic flow, but which is Anosov for all riemannian symmetric spaces of non-compact type. The basic idea in defining this action is to view geodesics not as particular curves of the space, but as totally geodesic flat subspaces. So we are led to consider flats, i.e. maximal totally geodesic flat subspaces, and Weyl chambers as particular subsets of flats. The bundle of all Weyl chambers will be the phase space of the action. The geodesic flow is given by moving geodesic rays along their supporting geodesics. Similarly, the action to be defined here consists of parallel translating Weyl chambers within their supporting flats. If the rank of the space is one, then flats are geodesics, Weyl chambers are geodesic rays, and our action coincides with the geodesic flow. For spaces of rank higher than one, i.e. spaces with higher-dimensional flats, this action is no longer a flow, but an action of a higher-dimensional abelian Lie group.

Let us recall the definition of an *Anosov action* (cf. [5]; for the ergodicity of Anosov actions, see [8]). An action φ of a Lie group L on a riemannian manifold M is said to be *Anosov* if the orbits under φ foliate M , and if there exists at least one Anosov element in L , i.e. an element $a \in L$ such that the tangent bundle $T(M)$ admits a $d\varphi_a$ -invariant splitting

$$T(M) = E^s \oplus E^c \oplus E^u$$

into three sub-bundles: the *central bundle* E^c tangent to the orbits under φ , the *strong stable bundle* E^s , on which the differential $d\varphi_a$ contracts more than on E^c , and the *strong unstable bundle* E^u , on which $d\varphi_a$ expands more than on E^c .

Given such a splitting, the *stable* and *unstable bundles* are defined to be the sums $E^s \oplus E^c$ and $E^u \oplus E^c$, respectively. Finally, the foliations of $T(M)$ tangent to E^s , E^u , $E^s \oplus E^c$, and $E^u \oplus E^c$ are called respectively the *strong stable*, *strong unstable*, *stable*, and *unstable foliation* induced by φ_a .

The first part of this note is purely group-theoretical. There we shall define our action and prove its Anosov character. Geometrical interpretations will be given in the second part, where we shall introduce the bundle of Weyl chambers and relate our action to the geodesic flow. Finally the stable and unstable foliations will find an interpretation in terms of horocycles.

The content of this note is part of the author's Habilitationsschrift [6]. I wish to thank Ernst Ruh and Mikhael Gromov for their encouragement and criticism. For financial support I am indebted to the Sonderforschungsbereich 'Theoretische Mathematik' at the University of Bonn, the Stiftung Volkswagenwerk, and the IHES in Bures-sur-Yvette, France.

1. The action of A on G/M

Here we state and prove our main result in group-theoretical terms. Let G be a real semisimple Lie group of non-compact type and finite centre. A maximal compact subgroup $K \subset G$ gives rise to a Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, where \mathfrak{g} and \mathfrak{k} denote the respective Lie algebras of G and K , and \mathfrak{p} is the orthogonal complement to \mathfrak{k} with respect to the Killing form on \mathfrak{g} . Let $\mathfrak{a} \subset \mathfrak{p}$ be a maximal abelian subalgebra and $A = \exp \mathfrak{a} \subset G$ the corresponding subgroup. By $\log: A \rightarrow \mathfrak{a}$ we shall denote the inverse of $\exp: \mathfrak{a} \rightarrow A$. The pair $(\mathfrak{g}, \mathfrak{a})$ gives rise to a root system $\Lambda \subset \text{Hom}(\mathfrak{a}, \mathbb{R})$ and a root space decomposition

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \sum_{\lambda \in \Lambda} \mathfrak{g}_\lambda.$$

An element $a \in A$ is called *regular* if $\lambda(\log a) \neq 0$ for all $\lambda \in \Lambda$, otherwise it is called *singular*. Now consider $M = Z_K(A)$, the centralizer of A in K . By defining $\varphi(a, gM) = \varphi_a(gM) = gaM$ we obtain an action φ of A on the homogeneous space G/M .

THEOREM. *The action $\varphi: A \times G/M \rightarrow G/M$ defined by $\varphi(a, gM) = gaM$ is an Anosov action.*

Proof. The orbits under φ coincide with the fibres of the natural projection $G/M \rightarrow G/MA$, therefore they are the leaves of a foliation. Now we shall prove that all regular elements of A are Anosov elements for φ . At first we fix an arbitrary $a \in A$. We have to compute the differential of

$$\varphi_a: G/M \rightarrow G/M$$

at a point $gM \in G/M$. Let \mathfrak{m} denote the Lie algebra of M . Since all tangent spaces of G/M are identified with $\mathfrak{g}/\mathfrak{m}$ by left translation, it suffices to compute

$$d\varphi_a: \mathfrak{g}/\mathfrak{m} \rightarrow \mathfrak{g}/\mathfrak{m}.$$

Let $R_a: G \rightarrow G$ denote right translation by a . The projection of R_a onto G/M is just φ_a . The differential of R_a is $\text{Ad}(a^{-1}): \mathfrak{g} \rightarrow \mathfrak{g}$, and this in turn projects onto $\mathfrak{g}/\mathfrak{m}$. Let us denote this projection again by $\text{Ad}(a^{-1})$, then we have

$$d\varphi_a = \text{Ad}(a^{-1}): \mathfrak{g}/\mathfrak{m} \rightarrow \mathfrak{g}/\mathfrak{m}.$$

In order to compute $\text{Ad}(a^{-1})$ we use the root space decomposition of \mathfrak{g} with respect

to a . Observing $\mathfrak{g}_0 = \mathfrak{m} \oplus \mathfrak{a}$, we have

$$\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{a} \oplus \sum_{\lambda \in \Lambda} \mathfrak{g}_\lambda,$$

therefore

$$\mathfrak{g}/\mathfrak{m} \cong \mathfrak{a} \oplus \sum_{\lambda \in \Lambda} \mathfrak{g}_\lambda.$$

According to this isomorphism an element $\xi \in \mathfrak{g}/\mathfrak{m}$ will be written as

$$\xi = \xi_0 + \sum_{\lambda \in \Lambda} \xi_\lambda, \tag{1}$$

with $\xi_0 \in \mathfrak{a}$ and $\xi_\lambda \in \mathfrak{g}_\lambda$. The equation

$$\text{Ad}(a^{-1}) = \exp(\text{ad}(-\log a))$$

in $\text{GL}(\mathfrak{g})$ implies

$$\text{Ad}(a^{-1})(\xi) = \xi_0 + \sum_{\lambda \in \Lambda} e^{-\lambda(\log a)} \xi_\lambda \tag{2}$$

for ξ as given by (1).

Assume now that a is regular. Then Λ splits into the two subsets

$$\Lambda^+ = \{\lambda \in \Lambda; \lambda(\log a) > 0\}, \quad \Lambda^- = \{\lambda \in \Lambda; \lambda \log a < 0\}.$$

Defining

$$\mathfrak{n}^+ = \sum_{\lambda \in \Lambda^+} \mathfrak{g}_\lambda, \quad \mathfrak{n}^- = \sum_{\lambda \in \Lambda^-} \mathfrak{g}_\lambda,$$

we obtain

$$\mathfrak{g}/\mathfrak{m} = \mathfrak{a} \oplus \mathfrak{n}^+ \oplus \mathfrak{n}^-. \tag{3}$$

We claim that this decomposition, extended to all tangent spaces of G/M by left translation, induces the desired splitting of $T(G/M)$. Indeed, \mathfrak{a} is tangent to the orbits of φ , and by (2) we get

$$\begin{aligned} d\varphi_a(\xi) &= \xi \in \mathfrak{a}, \\ d\varphi_a(\xi) &= \sum_{\lambda \in \Lambda^+} e^{-\lambda(\log a)} \xi_\lambda \in \mathfrak{n}^+, \\ d\varphi_a(\xi) &= \sum_{\lambda \in \Lambda^-} e^{-\lambda(\log a)} \xi_\lambda \in \mathfrak{n}^-, \end{aligned}$$

for $\xi \in \mathfrak{a}$, $\xi = \sum_{\lambda \in \Lambda^+} \xi_\lambda \in \mathfrak{n}^+$, $\xi = \sum_{\lambda \in \Lambda^-} \xi_\lambda \in \mathfrak{n}^-$, respectively. Thus (3) is invariant under $d\varphi_a$.

In order to establish the necessary estimates we need a metric on G/M . Inverting the sign of the Killing form on \mathfrak{k} we get a positive definite scalar product on \mathfrak{g} which we restrict onto $\mathfrak{g}/\mathfrak{m} \cong \mathfrak{a} \oplus \sum_{\lambda \in \Lambda} \mathfrak{g}_\lambda$. By left translation this induces a riemannian metric on G/M . Since root spaces are orthogonal to each other, we have

$$\|\xi\|^2 = \|\xi_0\|^2 + \sum_{\lambda \in \Lambda} \|\xi_\lambda\|^2,$$

for ξ as given by (1). Therefore our expressions for $d\varphi_a(\xi)$ imply

$$\begin{aligned} \|d\varphi_a(\xi)\| &= \|\xi\| && \text{for } \xi \in \mathfrak{a}; \\ \|d\varphi_a(\xi)\| &\leq e^{-k} \|\xi\| && \text{for } \xi \in \mathfrak{n}^+; \\ \|d\varphi_a(\xi)\| &\geq e^k \|\xi\| && \text{for } \xi \in \mathfrak{n}^-; \end{aligned}$$

where $k = \min \{\lambda(\log a); \lambda \in \Lambda^+\}$ is a positive constant depending only on a . □

Remark. The same arguments applied to a singular element $a \in A$ lead to the decomposition

$$g/m = \left(a \oplus \sum_{\lambda(\log a)=0} g_\lambda \right) \oplus \sum_{\lambda(\log a)>0} g_\lambda \oplus \sum_{\lambda(\log a)<0} g_\lambda,$$

which exhibits a component transverse to a , where $d\varphi_a$ is neither contracting nor expanding. Therefore singular elements of A are not Anosov elements for the action φ .

2. *Geometric interpretations*

Here we give geometric interpretations of the action $\varphi: A \times G/M \rightarrow G/M$ and of the stable and unstable foliations induced by this action. Let X be a riemannian symmetric space of non-compact type. The maximal totally geodesic flat subspaces of X are called *flats*. The common dimension of all flats of X is called the *rank* of X . Each geodesic is contained in at least one flat. A geodesic is called *regular* (*singular*), if it is contained in exactly one flat (in more than one flat). We fix a point $x \in X$ and a flat $F \subset X$ with $x \in F$. The singular geodesics in F through x , are contained in the union of finitely many hyperplanes in F through x , and conversely, any geodesic lying in one of these hyperplanes is singular. The connected components of F after removing all singular geodesics through x are called the *Weyl chambers* of F based at x . Admitting all possible flats $F \subset X$ and all possible base points $x \in F$, we obtain the set CX of all Weyl chambers of X . Let $\pi: CX \rightarrow X$ denote the map which assigns to each Weyl chamber its base point. Following Mostow ([7, § 4]) we call two Weyl chambers *asymptotic*, if each one lies in a tubular neighbourhood of the other. The set of classes of asymptotic Weyl chambers is called the *boundary* of X . Let τ denote the map which assigns to each Weyl chamber its asymptoticity class.

A flat with a distinguished class of asymptotic Weyl chambers is called an *oriented flat*. Let ρ denote the map which assigns to a Weyl chamber c its supporting flat together with the orientation induced by c .

Finally we shall sketch a geometric definition of horocycles (for more details see [6]). Each geodesic ray in X determines a horosphere passing through the initial point of the ray and orthogonal to it (cf. [1, § 3]). For a fixed Weyl chamber c based at x , the *horocycle* associated with c is defined as the intersection of all horospheres determined by the (regular) geodesic rays emanating from x and contained in c . This intersection is a submanifold whose codimension equals the rank of X . Let σ denote the map which assigns to each Weyl chamber its associated horocycle.

Now we shall translate the geometrical picture introduced so far into the group-theoretical language. Let G be the connected component of the isometry group of X ; it is a real semisimple Lie group of non-compact type and trivial centre. Choose a point $x_0 \in X$ and let K be the isotropy group of x_0 . Then K is a maximal compact subgroup of G , and X is identified with the homogeneous space G/K . Furthermore, the choice of x_0 gives rise to a Cartan decomposition $g = \mathfrak{k} \oplus \mathfrak{p}$, where the vector part \mathfrak{p} is identified with the tangent space of X at x_0 .

Now consider a flat F through x_0 . Its tangent space at x_0 is a maximal abelian subalgebra $\mathfrak{a} \subset \mathfrak{g}$, and conversely, each such subalgebra corresponds to a flat through x_0 . The group K operates transitively on the set of all flats through x_0 (cf. [3, Chapter V, Lemma 6.3]).

We now fix a flat F_0 through x_0 . Let \mathfrak{a} denote the corresponding maximal abelian subalgebra of \mathfrak{g} , and set $A = \exp \mathfrak{a} \subset G$. The group A acts on F_0 as its group of translations. The root system Λ with respect to the pair $(\mathfrak{g}, \mathfrak{a})$ defines Weyl chambers in \mathfrak{a} and, via the exponential map, in A . These correspond to the Weyl chambers in F_0 based at x_0 . The Weyl group W of X is defined as the quotient M'/M , where $M = Z_K(A)$ is the centralizer and $M' = N_K(A)$ the normalizer of A in K . It acts simply transitively on the set of Weyl chambers in F_0 based at x_0 (for spaces of compact type, see [3, Chapter VII, Theorem 2.12]). Let c_0 be one of the Weyl chambers in F_0 based at x_0 . The choice of c_0 determines a set $\Lambda^+ \subset \Lambda$ of positive roots. Define $\mathfrak{n} = \sum_{\lambda \in \Lambda^+} \mathfrak{g}_\lambda$ and $N = \exp \mathfrak{n} \subset G$. Finally define $P = MAN$.

The Weyl chamber c_0 and the groups derived from this choice shall be fixed for the rest of this paper.

PROPOSITION. *Let X be a riemannian symmetric space of non-compact type. Then the set of Weyl chambers of X is identified with the homogeneous space G/M , the set of oriented flats with G/MA , the set of horocycles with G/MN , and the boundary of X with G/P . Furthermore, the maps π, ρ, σ , and τ introduced above coincide with the natural projections from G/M to $G/K, G/MA, G/MN$, and G/P , respectively.*

Proof. We shall give a proof for the case of G/M . First we have to show that G operates transitively on CX . Let c be any Weyl chamber, F the flat containing c , and x the base point of c . An isometry $g \in G$ takes the flat F into a flat gF through the point gx , and it preserves the configuration of regular and singular geodesics. Therefore the Weyl chamber c is mapped into a Weyl chamber gc contained in gF and based at gx .

Now we shall find an isometry which takes c into c_0 . There is a $g \in G$ such that $gx = x_0$, thus $x_0 \in gF$. An element $k \in K$ maps gF into F_0 , hence $kgF = F_0$. Now kgc and c_0 are two Weyl chambers in F_0 based at x_0 , therefore there is an element $m' \in M'$ such that $m'kgc = c_0$.

Finally we shall determine the isotropy group of c_0 in G . Suppose $gc_0 = c_0$. Then certainly $gx_0 = x_0$, hence $g \in K$. Moreover $gF_0 = F_0$, thus $g \in M'$. Since $W = M'/M$ acts simply transitively on the set of Weyl chambers in F_0 based at x_0 , it follows that $g \in M$. Conversely, we clearly have $mc_0 = c_0$ for $m \in M$.

The proof for G/MA is similar to the one for G/M , so we shall omit it.

The case of G/MN is treated in [4] in purely group-theoretical terms. There, the horocycle associated with $c = gc_0$ is defined as the orbit of gx_0 under the group gNg^{-1} . The relation with our geometric notion of horocycles is established in [6].

The identification of the boundary G/P with the set of classes of asymptotic Weyl chambers is proved in [7, § 4].

The statements about the natural projections are evident. □

In addition to the obvious left action of G on the spaces $G/M, G/MA, G/MN,$ and $G/P,$ there is a right action of the Weyl group W on these spaces. Let m_w denote a representative for an element $w \in W.$ Then the effect of w on a Weyl chamber gM is given by $gM \mapsto gm_wM.$ The action of W on the other spaces is defined similarly, and so all these W -actions are compatible with natural projections.

Now recall the action $\varphi: A \times G/M \rightarrow G/M$ given by $\varphi(a, gM) = gaM.$ The orbit of a Weyl chamber $gM \in G/M$ under φ is $\{gaM \in G/M; a \in A\},$ it coincides with the fibre $\rho^{-1}(gMA)$ of the projection $\rho: G/M \rightarrow G/MA.$ In other words, the effect of φ consists of translating a given Weyl chamber along its supporting flat to another Weyl chamber of the same flat and asymptotic to the given one. In the rank one case, Weyl chambers coincide with geodesic rays, and these in turn may be identified with unit tangent vectors. Moreover, since the group A is one-dimensional it admits a parametrization $A = \{a(t) \in G; t \in \mathbb{R}\}.$ Therefore the action φ may be viewed as an action of \mathbb{R} on the unit tangent bundle of $X.$ The effect of φ consists in this case of translating a given geodesic ray along its supporting geodesic to another ray of the same geodesic and pointing to the same direction. This is exactly the effect of the geodesic flow. (For a group-theoretical description of the geodesic flow, see [2].)

Finally we shall determine the leaves of the stable and strong stable foliations induced by $\varphi.$ First we shall consider a special case. Let a_0 be a regular element of A with $\lambda(\log a_0) > 0$ for all $\lambda \in \Lambda^+.$ For such an element, the contracting subspace of $\mathfrak{g}/\mathfrak{m} \cong \mathfrak{a} \oplus \sum_{\lambda \in \Lambda} \mathfrak{g}_\lambda$ is $\mathfrak{n} = \sum_{\lambda \in \Lambda^+} \mathfrak{g}_\lambda.$ Therefore the leaf of the strong stable foliation through $c_0 = M \in G/M$ equals the orbit

$$\{nM \in G/M; n \in N\}$$

of c_0 under $N.$ This orbit coincides with the fibre $\sigma^{-1}(MN)$ of the projection $\sigma: G/M \rightarrow G/MN,$ i.e. it consists of all Weyl chambers asymptotic to c_0 and based at the points of the horocycle $Z_0 = Nx_0 = MN \in G/MN.$

The stable foliation is tangent to $\mathfrak{a} \oplus \mathfrak{n}.$ Its leaf through c_0 equals the orbit

$$\{anM \in G/M; a \in A, n \in N\}$$

of c_0 under $AN,$ it coincides with the fibre $\tau^{-1}(P)$ of the projection $\tau: G/M \rightarrow G/P,$ i.e. it consists of all Weyl chambers asymptotic to $c_0.$

Now consider the case of an arbitrary regular element $a \in A.$ There is a unique Weyl group element $w \in W$ such that $a = m_w a_0 m_w^{-1},$ where $m_w \in M'$ represents $w,$ and a_0 satisfies $\lambda(\log a_0) > 0$ for all $\lambda \in \Lambda^+.$ The contracting subspace with respect to the action of a is

$$\sum_{\lambda(\log a) > 0} \mathfrak{g}_\lambda = \text{Ad}(m_w)\mathfrak{n}.$$

Therefore the leaf of the strong stable foliation through c_0 equals the orbit

$$\{m_w n m_w^{-1} M \in G/M; n \in N\}$$

of c_0 under $m_w N m_w^{-1}.$ This is no longer a set of asymptotic Weyl chambers, it is obtained from the set $\{m_w n M \in G/M; n \in N\}$ of Weyl chambers asymptotic to $m_w c_0$ and based at the points of the horocycle $m_w Z_0$ by applying w^{-1} from the right.

The leaf of the stable foliation through c_0 equals the orbit

$$\{m_w a n m_w^{-1} M \in G/M; a \in A, n \in N\}$$

of c_0 under $m_w A N m_w^{-1} = A m_w N m_w^{-1}$. It consists of the right w^{-1} -images of all Weyl chambers asymptotic to $m_w c_0$.

The unstable foliations with respect to a coincide with the stable foliations with respect to a^{-1} , so they need no further attention.

REFERENCES

- [1] P. Eberlein & B. O'Neill. Visibility manifolds. *Pacific J. Math.* **46** (1973), 45–109.
- [2] I. M. Gel'fand & S. V. Fomin. Geodesic flows on manifolds of constant negative curvature. *Uspehi Mat. Nauk (N.S.)* **7** (1952), 118–137; *Amer. Math. Soc. Translations (2)* **1** (1955), 49–65.
- [3] S. Helgason. *Differential Geometry and Symmetric Spaces*. Academic Press: New York 1962.
- [4] S. Helgason. Duality and Radon transform for symmetric spaces. *Amer. J. Math.* **85** (1963), 667–692.
- [5] M. W. Hirsch. Foliations and non-compact transformation groups. *Bull. Amer. Math. Soc.* **76** (1970), 1020–1023.
- [6] H. C. Im Hof. Die Geometrie der Weylkammern in symmetrischen Räumen vom nichtkompakten Typ. Habilitationsschrift, Bonn, 1979.
- [7] G. D. Mostow. *Strong Rigidity of Locally Symmetric Spaces*. Princeton University Press: Princeton, 1973.
- [8] C. Pugh & M. Shub. Ergodicity of Anosov actions. *Invent. Math.* **15** (1972), 1–23.