

Colour measurement of colostrum for estimation of colostral IgG and colostrum composition in dairy cows

Josef J Gross, Evelyne C Kessler and Rupert M Bruckmaier*

Veterinary Physiology, Vetsuisse Faculty, University of Bern, CH-3001 Bern, Switzerland

Received 15 April 2014; accepted for publication 26 June 2014; first published online 16 September 2014

Instruments for on-farm determination of colostrum quality such as refractometers and densimeters are increasingly used in dairy farms. The colour of colostrum is also supposed to reflect its quality. A paler or mature milk-like colour is associated with a lower colostrum value in terms of its general composition compared with a more yellowish and darker colour. The objective of this study was to investigate the relationships between colour measurement of colostrum using the CIELAB colour space (CIE L^* =from white to black, a^* =from red to green, b^* =from yellow to blue, chroma value G =visual perceived colourfulness) and its composition. Dairy cow colostrum samples ($n=117$) obtained at 4.7 ± 1.5 h after parturition were analysed for immunoglobulin G (IgG) by ELISA and for fat, protein and lactose by infrared spectroscopy. For colour measurements, a calibrated spectrophotometer was used. At a cut-off value of 50 mg IgG/ml, colour measurement had a sensitivity of 50.0%, a specificity of 49.5%, and a negative predictive value of 87.9%. Colostral IgG concentration was not correlated with the chroma value G , but with relative lightness L^* . While milk fat content showed a relationship to the parameters L^* , a^* , b^* and G from the colour measurement, milk protein content was not correlated with a^* , but with L^* , b^* , and G . Lactose concentration in colostrum showed only a relationship with b^* and G . In conclusion, parameters of the colour measurement showed clear relationships to colostral IgG, fat, protein and lactose concentration in dairy cows. Implementation of colour measuring devices in automatic milking systems and milking parlours might be a potential instrument to access colostrum quality as well as detecting abnormal milk.

Keywords: Colostrum, colour, immunoglobulin G, dairy cow.

Colostrum, defined as the first secretion of the mammary gland after parturition (Jaster, 2005), is characterised by its high content of immunoglobulins (Ig), predominantly IgG (Butler, 1981). Calves are born agammaglobulinaemic since the placenta of dairy cows is impermeable to Ig and consequently depend on timely administration of high-quality colostrum to acquire immunisation. Determination of colostrum quality, i.e. in a narrower sense Ig concentration, on farm must be rapid and give reliable results. Currently, instruments to measure specific gravity (hydrometer/densimeter/colostrometer) or the refractive index of colostrum (Brix refractometer) are used in practice. However, the accuracy in estimation of Ig concentration by using these methods is limited. Relationships between gravity and refractive measurements and colostral protein content were reported to be higher than the correlation with

IgG₁ concentration that represents most of the Ig in colostrum (Fleenor & Stott, 1980; Quigley et al. 1994; Morin et al. 2001; Gross et al. 2013). Furthermore, temperature of colostrum (Mechor et al. 1992; Chigerwe et al. 2008) and season of calving (Morin et al. 2001) affect the validity of gravity measurements.

Colostrum colour can have a wide spectrum, from dark brown/red over yellow to pale white (Fig. 1). Brandt et al. (2010) gave an overview on technical solutions including colour measurement to detect and separate abnormal milk. In the study of Madsen et al. (2004), colour measurement was identified as a potential tool to follow changes from colostrum to marketable milk with respect to identification of unsuitable milk for the dairy industry. Argüello et al. (2005) confirmed the high sensitivity and suitability of colour measurement for predicting colostral IgG concentration in goats. However, Argüello et al. (2005) based their findings on a pool of first and second milking samples. Furthermore, owing to the perceptual differences of colour distances and chroma values, especially in yellow shades, the CIELAB

*For correspondence; e-mail: rupert.bruckmaier@vetsuisse.unibe.ch



Fig. 1. Spectrum of colostrum colours in dairy cows compared with mature milk (right).

colour system was modified according to the DIN99 formula in order to harmonise sensation of colours (ASTM D2244, 2011).

Up to now, no results showing the association between colostrum composition and its colour have been reported for dairy cows. However, the colour measurement was successfully tested to detect elevated milk cell contents in automatic milking systems (Brandt et al. 2010) and for determination of milk quality in dairy industries (Solah et al. 2007). The objective of this study was to investigate the relationships between colostrum IgG, fat, protein, and lactose concentration and colour measurement (including latest colour system adaptations to DIN99) in first colostrum of dairy cows. Furthermore, sensitivity, specificity, and negative predictive value (NPV) of the colour method were evaluated for suitability to predict IgG concentration in dairy cow colostrum.

Materials and methods

From September to December 2012, 117 colostrum samples of primiparous ($n=81$) and multiparous ($n=36$) Holstein dairy cows milked at $4\text{ h }42\text{ min} \pm 1\text{ h }30\text{ min}$ post partum (mean \pm SD) were obtained by machine milking at the Agroscope research farm Posieux, Switzerland. Cows were fed the same lot of hay for ad-libitum intake during the dry period until parturition. Samples were frozen at -20°C until analysis of IgG by ELISA and content of fat, protein, and lactose by infrared spectroscopy as described recently (Gross et al. 2014). Colostrum colour [CIE 1976 (L^* , a^* , b^*) colour space – CIELAB] was measured in thawed and homogenised samples in triplicate at 25°C using a calibrated Microflash 200 d spectrophotometer (Datacolor International, Dietikon, Switzerland), with the coordinates L^* representing relative lightness (black to white), a^* depicting the relative position between green and red, and b^* indicating the relative position between blue and yellow. According to the DIN99 formula (ASTM D2244, 2011), measured a^* and b^* values were transformed using

the following equations:

$$\text{Redness } e = a^* \times \cos(16^\circ) + b^* \times \sin(16^\circ)$$

$$\text{Yellowness } f = 0.7 \times (-a^* \times \sin(16^\circ) + b^* \times \cos(16^\circ))$$

The parameters e and f were used to calculate the chroma value G (indicating visual perceived colourfulness):

$$G = \sqrt{e^2 + f^2}.$$

Data were analysed using the statistical program package SAS (Version 9.2, SAS Institute, Cary NC, USA). The UNIVARIATE procedure was used to check for normal distribution of the data and identification of outliers. Using the CORR procedure, Pearson correlation coefficients were calculated to describe the relationships between colour measurement coordinates (L^* , a^* , b^* , G) and colostrum IgG, fat, protein, and lactose concentration. P values < 0.05 were considered to be significant. Sensitivity, specificity, and the negative predictive value (NPV) of the colour measurement for testing IgG concentration were calculated using two-way contingency tables according to Argüello et al. (2005). Considering IgG values of the ELISA analysis as true values, the colour method was classified as the test value. For this study, different threshold values between high- and low-quality colostrum were chosen: 50, 75, and 100 mg IgG/ml. Sensitivity was calculated by the proportion of samples identified by the colour method to have an IgG concentration < 50 (75 and 100, respectively) mg/ml in all samples with a true IgG concentration < 50 (75 and 100, respectively) mg/ml. Specificity represents the probability of samples with an IgG concentration ≥ 50 (75 and 100, respectively) mg/ml identified by the colour method relative to all samples classified as high quality by the ELISA measurement. Calculation of the NPV was in agreement with Pritchett et al. (1994) and Argüello et al. (2005), and depicts the portion of truly high samples (estimated via ELISA) in all samples being identified as high-quality colostrum by the colour method. Since cows in the present study were fed the same diet, feed is unlikely to explain variation in the results of colour measurement.




Fig. 2. Relationship between colostral IgG concentration and Chroma G for colostrum of primiparous and multiparous dairy cows.

Table 1. Two-way contingency table [according to Argüello et al. (2005)] for calculation of sensitivity†, specificity‡, and negative predictive value (NPV)§ of colour measurement to estimate colostral IgG concentration at a threshold of 50 mg/ml. Figures in the table show the number of observations

	IgG concentration determined by ELISA		Total
	Low-quality (< 50 mg/ml)	High-quality (≥ 50 mg/ml)	
Colour measurement (Chroma G)			
Low-quality	7 ^a	52 ^b	59
High-quality	7 ^c	51 ^d	58
Total	14	103	117

† Sensitivity = $a/(a+c) \times 100 = 7/14 \times 100 = 50.0\%$

‡ Specificity = $d/(b+d) \times 100 = 51/103 \times 100 = 49.5\%$

§ NPV = $d/(c+d) \times 100 = 51/58 \times 100 = 87.9\%$

Results and discussion

Colour of milk in dairy cows was shown to be clearly dominated by carotenoids and their transfer from blood to milk (Nozière et al. 2006; Calderón et al. 2007). Though carotenoids were not analysed in milk samples in the present study, hay feeding provided only a small amount of β-carotene in the feed. All cows were of the same breed, and the different parities might contribute to only a small degree on variation in plasma and milk carotenoid content (Nozière et al. 2006). Further factors such as metabolism and transfer rate of carotenoids might lead to considerable individual variation between cows despite a constant feeding management (Nozière et al. 2006; Calderón et al. 2007) and might have an impact on the observations.

No significant relationship between the Chroma G and IgG concentration in first colostrum of dairy cows was

Table 2. Two-way contingency table [according to Argüello et al. (2005)] for calculation of sensitivity†, specificity‡, and negative predictive value (NPV)§ of colour measurement to estimate colostral IgG concentration at a threshold of 75 mg/ml. Figures in the table show the number of observations

	IgG concentration determined by ELISA		Total
	Low-quality (< 75 mg/ml)	High-quality (≥ 75 mg/ml)	
Colour measurement (Chroma G)			
Low-quality	17 ^a	42 ^b	59
High-quality	25 ^c	33 ^d	58
Total	42	75	117

† Sensitivity = $a/(a+c) \times 100 = 17/42 \times 100 = 40.5\%$

‡ Specificity = $d/(b+d) \times 100 = 33/75 \times 100 = 44.0\%$

§ NPV = $d/(c+d) \times 100 = 33/58 \times 100 = 56.9\%$

Table 3. Two-way contingency table [according to Argüello et al. (2005)] for calculation of sensitivity†, specificity‡, and negative predictive value (NPV)§ of colour measurement to estimate colostral IgG concentration at a threshold of 100 mg/ml. Figures in the table show the number of observations

	IgG concentration determined by ELISA		Total
	Low-quality (< 100 mg/ml)	High-quality (≥ 100 mg/ml)	
Colour measurement (Chroma G)			
Low-quality	30 ^a	29 ^b	59
High-quality	37 ^c	21 ^d	58
Total	67	50	117

† Sensitivity = $a/(a+c) \times 100 = 30/67 \times 100 = 44.8\%$

‡ Specificity = $d/(b+d) \times 100 = 21/50 \times 100 = 42.0\%$

§ NPV = $d/(c+d) \times 100 = 21/58 \times 100 = 36.2\%$

found in the present study ($r^2=0.0061$; $P=0.40$; Fig. 2) contrary to the findings in goat colostrum in the study of Argüello et al. (2005). Parity did not have an effect on the relationship between IgG concentration and the Chroma G ($P=0.81$; Fig. 2). Values for sensitivity, specificity and NPV were highest at the threshold of 50 mg IgG/ml compared to 75 and 100 mg/ml in colostrum (Tables 1–3). The threshold of 50 mg IgG/ml is commonly accepted to classify good colostrum in dairy cows (Tyler et al. 1996, 1999). Similar values for sensitivity and NPV were reported for the usage of a hydrometer to predict IgG₁ concentration in dairy cow colostrum by Pritchett et al. (1994), though values for sensitivity, specificity and NPV were markedly lower compared with the findings for colour measurements in goat colostrum reported by Argüello et al. (2005) and Brix refractometry in dairy cow colostrum by Quigley et al. (2013). As already shown above, IgG concentration was not correlated with Chroma G ($P=0.40$), but a significant

Table 4. Pearson correlation coefficients (*r*) between colostrum IgG, fat, protein, and lactose concentrations and the CIE coordinates *L**, *a**, *b** and Chroma *G*

Concentration of:	<i>L*</i>		<i>a*</i>		<i>b*</i>		<i>G</i>	
	<i>R</i>	<i>P</i> -value	<i>r</i>	<i>P</i> -value	<i>r</i>	<i>P</i> -value	<i>r</i>	<i>P</i> -value
IgG, mg/ml	-0.45	<0.0001	0.03	0.77	-0.10	0.29	-0.08	0.40
Fat, %	0.50	<0.0001	0.40	<0.0001	0.51	<0.0001	0.51	<0.0001
Protein, %	-0.29	<0.01	0.02	0.87	0.27	<0.01	0.24	<0.05
Lactose, %	0.16	0.11	-0.14	0.15	-0.44	<0.0001	-0.41	<0.0001

correlation was found with the relative lightness *L** ($P < 0.0001$, Table 4). Besides IgG concentration, colostrum constituents (fat, protein, and lactose) contribute to the nutritive value of colostrum. Milk fat content was significantly correlated with *L**, *a**, *b** and *G* ($P < 0.0001$, Table 4). Milk protein content did not show a significant correlation with *a**, but with *L**, *b** and *G* (Table 4). Lactose content in colostrum was not correlated with *L** and *a** ($P > 0.05$), but with *b** and *G* ($P < 0.0001$, Table 4). Low correlations between parameters of colour measurements (*a**, *b**, *G*) and colostrum constituents (IgG and protein) might be explained by their high variation compared with the relative concentrations of fat and lactose.

The predictive value to estimate colostrum IgG concentration by colour measurement for dairy cows showed comparable values to sensitivity and NPV obtained by specific gravity (Pritchett et al. 1994) and refractive measurements (Quigley et al. 2013). Though not being as accurate as reported for goats by Argüello et al. (2005), colour measurements of dairy cow colostrum might be a new tool to assess colostrum quality. Implementation of this technique on dairy farms is already state of the art in automatic milking systems to separate abnormal milk, e.g. during mastitis (Brandt et al. 2010). Adjustments for colostrum would be the next step. Argüello et al. (2005) suggested a colour fan for visual estimation of IgG concentration by farm personnel.

The present study confirmed the relationship between milk fat content and lightness *L** as well as yellowness *b** in agreement with Solah et al. (2007). Protein in colostrum contributed to lightness *L** in samples of the present study in agreement with findings for mature milk (Quinones et al. 1998; Solah et al. 2007). Thus, colostrum with a more yellow and darker colour is very likely to contain more milk fat and protein as well as immunoglobulins compared with a more pale colostrum. However, it needs to be considered that also blood components and dietary composition (e.g. β -carotene) might affect colostrum colour (Madsen et al. 2004; Solah et al. 2007).

In conclusion, colour measurement of colostrum in dairy cows can be an additional instrument besides colostrometer and refractometer to assess colostrum quality. Sufficient sensitivity, specificity, and NPV of this method anticipate a reliable estimation of colostrum quality though the chroma value alone does not reflect IgG concentration in a satisfactory manner. Though colour measurement

does not improve estimation of IgG content in colostrum samples compared with currently available instruments and implies the further development of complementary and more accurate methods, other purposes (e.g. detection of non-deliverable milk) might be addressed in a very sensitive way.

The authors thank the Milchprüfing Baden-Württemberg e.V. (Ravensburg, Germany) and the Milchprüfing Bayern e.V. (Wolnzach, Germany) for coordination, validation, and performance of colostrum analyses.

References

- Argüello A, Castro N & Capote J 2005 Short Communication: evaluation of a color method for testing immunoglobulin G concentration in goat colostrum. *Journal of Dairy Science* **88** 1752–1754
- ASTM D2244-11 2011 Standard practice for calculation of color tolerances and color differences from instrumentally measured color coordinates. *ASTM International* DOI: 10.1520/D2244-11
- Brandt M, Haeussermann A & Hartung E 2010 Invited review: technical solutions for analysis of milk constituents and abnormal milk. *Journal of Dairy Science* **93** 427–436
- Butler JE 1981 A concept of humoral immunity among ruminants and an approach to its investigation. In *The Ruminant Immune System*, pp. 3–55 (Ed. JE Butler). New York: Plenum (Advances in experimental medicine and biology, Vol. 137).
- Calderón F, Chauveau-Duriot B, Martin B, Graulet B, Doreau M & Nozière P 2007 Variations in carotenoids, vitamins A and E, and color in cow's plasma and milk during late pregnancy and the first three months of lactation. *Journal of Dairy Science* **90** 2335–2346
- Chigerwe M, Tyler JW, Middleton JR, Spain JN, Dill JS & Stevens BJ 2008 Comparison of four methods to assess colostrum IgG concentration in dairy cows. *Journal of the American Veterinary Medical Association* **233** 761–766
- Fleener WA & Stott GH 1980 Hydrometer test for estimation of immunoglobulin concentration in bovine colostrum. *Journal of Dairy Science* **63** 973–977
- Gross JJ, Kessler EC & Bruckmaier RM 2013 Suitability of refractometer and densimeter for on-farm determination of colostrum quality in dairy cows and heifers. *Journal of Animal Science* **91**(E-Suppl. 2)/*Journal of Dairy Science* **96**(E-Suppl. 1) 325, W296. Abstract
- Gross JJ, Kessler EC, Bjerre-Harpoth V, Dechow C, Baumrucker CR & Bruckmaier RM 2014 Peripartur progesterone and prolactin have little effect on the rapid transport of immunoglobulin G into colostrum of dairy cows. *Journal of Dairy Science* **97** 2923–2931
- Jaster EH 2005 Evaluation of quality, quantity, and timing of colostrum feeding on immunoglobulin G1 absorption in Jersey calves. *Journal of Dairy Science* **88** 296–302
- Madsen BD, Rasmussen MD, Nielsen MO, Wiking L & Larsen LB 2004 Physical properties of mammary secretions in relation to chemical

- changes during transition from colostrum to milk. *Journal of Dairy Research* **71** 263–272
- Mechor GD, Gröhn YT, McDowell LR & Van Saun RJ** 1992 Specific gravity of bovine colostrum immunoglobulins as affected by temperature and colostrum components. *Journal of Dairy Science* **75** 3131–3135
- Morin DE, Constable PD, Maunsell FP & McCoy GC** 2001 Factors associated with colostrum specific gravity in dairy cows. *Journal of Dairy Science* **84** 937–943
- Nozière P, Graulet B, Lucas A, Martin B, Grolier P & Doreau M** 2006 Carotenoids for ruminants: from forages to dairy products. *Animal Feed Science and Technology* **131** 418–450
- Pritchett LC, Gay CC, Hancock DD & Besser TE** 1994 Evaluation of the hydrometer for testing immunoglobulin G1 concentrations in Holstein colostrum. *Journal of Dairy Science* **77** 1761–1767
- Quigley JD, Martin KR, Dowlen HH, Wallis LB & Lamar K** 1994 Immunoglobulin concentration, specific gravity, and nitrogen fractions of colostrum from Jersey cattle. *Journal of Dairy Science* **77** 264–269
- Quigley JD, Lago A, Chapman C, Erickson P & Polo J** 2013 Evaluation of the Brix refractometer to estimate immunoglobulin G concentration in bovine colostrum. *Journal of Dairy Science* **96** 1148–1155
- Quinones HJ, Barbano DM & Philips LG** 1998 Influence of protein standardization by ultrafiltration on the viscosity, color, and sensory properties of 2 and 3·3% milks. *Journal of Dairy Science* **81** 884–894
- Solah VA, Staines V, Honda S & Limley HA** 2007 Measurement of milk color and composition: effect of dietary intervention on Western Australian Holstein-Friesian cow's milk quality. *Journal of Food Science* **72** S560–S566
- Tyler JW, Hancock DD, Parish SM, Rea DE, Besser TE, Sanders SG & Wilson LK** 1996 Evaluation of 3 assays for failure of passive transfer in calves. *Journal of Veterinary Internal Medicine* **10** 304–307
- Tyler JW, Hancock DD, Thorne JG, Gay CC & Gay JM** 1999 Partitioning the mortality risk associated with inadequate passive transfer of colostrum immunoglobulins in dairy calves. *Journal of Veterinary Internal Medicine* **13** 335–337