

# Infection levels of proteocephalidean cestodes in *Cichla piquiti* (Osteichthyes: Cichlidae) of the Volta Grande Reservoir, Minas Gerais, Brazil, relative to host body weight and gender

M.L. Martins<sup>1\*</sup>, J. Pereira Jr<sup>2</sup>, A. de Chambrier<sup>3</sup>  
and J.L.P. Mouriño<sup>4</sup>

<sup>1</sup>AQUOS – Laboratory of Aquatic Organisms Health, Aquaculture Department, Federal University of Santa Catarina-UFSC, Rod. SC 404, km 3, 88040-900, Florianópolis, SC, Brazil: <sup>2</sup>Laboratory of Aquatic Organisms Parasitology, ICB, Federal University of Rio Grande – FURG, 96650-900, Rio Grande, RS, Brazil: <sup>3</sup>Department of Invertebrates, Natural History Museum Geneva, PO Box 6434, 1211 Geneva 6, Switzerland: <sup>4</sup>Laboratory of Marine Shrimp, UFSC, Rua Beco dos Coroas s/n, 88062-601, Barra da Lagoa, Florianópolis, SC, Brazil

(Accepted 23 November 2010; First Published Online 6 January 2011)

## Abstract

We evaluated the relationship between infection by proteocephalid cestodes and the sex and weight classes of tucunaré (*Cichla piquiti*) captured between August 1999 and June 2001 in the Volta Grande Reservoir, Minas Gerais, Brazil. A total of 96 fish, 75.9 ± 9.3% males and 88.9 ± 6.4% females, were parasitized by *Proteocephalus macrophallus* and *P. microscopicus*, with total mean intensities of 76.6 ± 23.9 and 145.2 ± 36.7, respectively, during this period. In the majority of the months analysed, males showed 71.4–100% prevalence of parasitism and females 80–100%. Although there was no significant difference, females showed a higher mean intensity of infection (145.2 ± 36.7) than males (76.6 ± 23.9). Fish weighing 300–800 g showed a higher mean abundance of parasites ( $P < 0.05$ ) compared with the biggest specimens weighing 801–2750 g. Analysing both males and females together, the greatest mean intensities of infection were found in October and December ( $P < 0.05$ ) independent of the year, which coincides with the months of highest rainfall. These results show that fish living in reservoirs may be more susceptible to intermediate hosts than those that live in rivers.

## Introduction

Proteocephalidean tapeworms (Cestoda: Proteocephalidea) are cosmopolitan parasites of freshwater fish, reptiles and amphibians (Schmidt, 1986), and have a worldwide distribution (Freze, 1965). They constitute the

most numerous group of South American cestodes and, until 2008, 109 nominal proteocephalidean species in freshwater fish were known (Rego *et al.*, 1999; de Chambrier, pers. comm.).

Machado *et al.* (1994) did not find a relationship between the sex of the host and proteocephalidean parasites in *Pseudoplatystoma corruscans* Agassiz, 1829 and *Schizodon borelli* Boulenger 1900. On the other hand, Takemoto & Pavanelli (2000) found that the host sex

\*Fax: + 55 48 33343441  
E-mail: mlaterca@cca.ufsc.br

influenced the intensity of infection by *Spatulifer maringaensis* and *Nupelia portoricensis* in *Sorubim lima* Bloch & Schneider 1801, and that the prevalence was correlated with the host length.

Machado *et al.* (2000), studying the proteocephalidean cestodes in *Cichla monoculus* Spix, 1831 from the Paraná River, found a positive correlation in the host length and prevalence of *Proteocephalus macrophallus* (Diesing, 1850) Scholz, de Chambrier, Prouza & Royero, 1996 and *Proteocephalus microscopicus* Woodland, 1935. Moreover, there were significant differences in the prevalence of these cestode species between male and female hosts. The sex of the host also influenced the intensity of infection by *S. maringaensis* and *N. portoricensis* in *Sorubim lima* Bloch & Schneider, 1801 (Takemoto & Pavanelli, 2000).

Martins *et al.* (2009) showed that *P. macrophallus* Diesing, 1850 and *P. microscopicus* Woodland, 1935 are parasites of *Cichla piquiti* Kullander & Ferreira, 2006 (tucunaré) (Cichlidae), a fish from the Amazon Basin introduced into the Volta Grande Reservoir, Minas Gerais, Brazil. These authors pointed out abiotic factors, except rainfall, as possible indirect causes for parasite dissemination. As tucunaré usually eats small fish (Gomiero & Braga, 2004a, b), cannibalism can be suggested as a possible explanation for the high prevalence and intensity of infection. Indeed, this situation is favoured by the exotic host in the reservoir (Martins *et al.*, 2009). According to Gomiero & Braga (2004c), who studied the fish population in the Volta Reservoir, Minas Gerais, Brazil, species of *Cichla* have a high frequency of immature individuals during summer and autumn (December to May).

This study aims to examine whether host sex and weight classes influence parasitism by proteocephalidean cestodes in *C. piquiti* Kullander & Ferreira, 2006 (*sin. part.* *C. ocellaris* 'sensu' Kullander & Ferreira, 2006) in the Volta Grande Reservoir, Minas Gerais, Brazil. In the studies of Martins *et al.* (2009), general aspects of parasitism were approached. Here, therefore, we analyse the same data from another point of view, where fish are separated by sex and weight classes.

## Materials and methods

### Collection and examination of fish

From August 1999 through to June 2001, a total of 96 tucunaré (50 males and 46 females) were captured with

net and hook and brought to the laboratory to be measured and weighed prior to parasitological analysis. The collection was conducted at the hydroelectric power station of the Volta Grande Reservoir (CEMIG), Grande River, Minas Gerais, Brazil (21°46'14"S, 42°32'20"W). In each sample, water pH, electrical conductivity, dissolved oxygen, chlorophyll, transparency and temperature were measured (table 1). The rainfall during the period of collection was measured by the CEMIG Hydrology Department. It was not possible to obtain water-quality data in February 2000 and June 2001. Worms collected from the intestine were relaxed in distilled water at 4°C, fixed in AFA (alcohol-formaldehyde-acetic acid) for 24 h for later storage in 70% alcohol and identification. The prevalence of infection, mean intensity (MI) and mean abundance were calculated according to Bush *et al.* (1997).

### Data analysis

Differences between the parasitological indexes of groups (host sex and weight classes) were analysed using the Kruskal–Wallis test. When statistically different by the Kruskal–Wallis test, a non-parametric multiple-comparisons test was applied to examine all possible comparisons. A *P* value of less than 0.05/[ $k \times (k - 1)/2$ ] was considered statistically significant, with *k* corresponding to the number of animals in the group. When statistically different, the mean intensity (MI) and the weight classes were compared by Fisher's exact test, to determine whether there were non-random associations between two categorical variables. The Pearson correlation analysis was used to verify the significance between the values of rainfall and water quality, and the prevalence and MI of infection and mean abundance of parasites.

## Results

### Water quality and rainfall

No significant correlation ( $P > 0.05$ ) was observed between prevalence rates, intensity of infection and mean abundance and the rainfall indexes and water quality. In spite of the low transparency in April 2000 (2.1 m), no variation in electrical conductivity and water pH was observed (table 1). The months with the highest

Table 1. Mean values of the rainfall (RF) and water quality measured in the Volta Grande Reservoir, Minas Gerais State, Brazil, between August 1999 and April 2001 during the collection of fish.

Months	RF (mm)	Transparency (m)	Chlorophyll ( $\mu\text{g/l}$ )	pH	Conductivity ( $\mu\text{S/cm}$ )	Temperature ( $^{\circ}\text{C}$ )	Oxygen (mg/l)
August 1999	6	2.85	4.70	7.16	34.67	21.10	7.50
October 1999	7	3.60	3.90	6.40	34.33	25.33	7.10
December 1999	241	3.00	–	6.66	34.00	26.20	7.63
April 2000	53	2.15	11.3	6.41	35.00	26.87	8.17
June 2000	0	3.85	1.70	6.23	35.00	21.83	8.90
August 2000	249	5.35	5.30	7.08	29.00	22.93	8.50
October 2000	669	3.25	3.00	6.87	35.00	26.93	7.80
December 2000	1617	3.90	3.00	7.52	35.33	27.83	7.47
February 2001	1582	7.30	2.60	7.90	34.33	28.40	6.73
April 2001	747	5.30	4.50	6.92	29.00	28.00	8.43

temperatures were December 2000 (27.8°C), February (28.4°C) and April 2001 (28.0°C). Dissolved oxygen did not vary more than 2.17 mg/l, while the chlorophyll analysis showed the highest value (11.30 µg/l) in April 2000. The rainfall means were high in December 1999, October 2000, December 2000 and April 2000 (table 1).

*Infection levels and host body weight*

The parasites were identified as *P. macrophallus* and *P. microscopicus*. When the fish were separated by classes of weight, the mean abundance of parasites of fish weighing 301–800 g was higher than in the biggest fish (fig. 1A). The mean abundance was 64.33 in those of 13–300 g weight; 150.96 in those of 301–800 g and 47.24 in those of 801–2750 g (fig. 1A). Fish weighing between 301 and 800 g showed a high range of intensity, varying from 3 to 1031 parasites per host. Consequently, fish weighing 301–800 g were more parasitized ( $P < 0.05$ ) than the largest ones (fig. 1A).

The months with the highest levels of parasitism were October and December, as shown in fig. 1B. Fish

examined in April (except for females), June and August showed lower MI of infection than those examined in October and December (fig. 1D). The highest distribution frequencies were: 1–10, 11–30, 31–70 and 101–200 parasites per host.

*Infection levels and host gender*

When analysing the infection related to the sex of the hosts, during the whole period, in males, the total prevalence rate was  $75.9 \pm 9.3\%$  with a MI of infection of  $76.6 \pm 23.9$  and, in females  $88.9 \pm 6.4\%$  prevalence and  $145.2 \pm 36.7$  MI. We observed high values of MI of infection in females captured in February 2000 (MI = 223), April 2000 (MI =  $404 \pm 314.2$ ), October 2000 (MI =  $259.0 \pm 96.6$ ), December 2000 (MI =  $204 \pm 74.8$ ) and April 2001 (MI =  $234 \pm 118.1$ ). Consequently, a 100% prevalence commonly occurred in females (table 2). In general, no difference was found between males and females during the whole period independently of the month (fig. 1C). Females captured in April and December




Fig. 1. Parasitological indexes in *Cichla piquiti* from Volta Grande Reservoir, Minas Gerais, Brazil, infected by proteocephalidean cestodes. (A) Relationship between mean abundance and weight classes of the host. (B) Variation in the mean intensity in the months analysed. (C) Mean intensity in males (M) and females (F). (D) Monthly changes in the mean intensity in males and females. \*All different lowercase letters indicate significant difference among bars ( $P < 0.05$ ).

Table 2. Parasitological indexes in *Cichla piquiti* parasitized by proteocephalidean cestodes from the Volta Grande Reservoir, Minas Gerais, Brazil. IM/CM, infected males/collected males; IFE/CFE, infected females/collected females; P, prevalence; MI, mean intensity followed by standard error and amplitude of variation in parentheses. \*Values obtained from the averages in the months. \*\*Total number of fish.

Months	Fish weight (g)	Fish length (cm)	IM/CM	P (%)	MI	IFE/CFE	P (%)	MI
August 1999	201.5 ± 10.4 (160–241)	24.5 ± 0.1 (23–25)	1/1	100.0	56.0	6/6	100.0	75.8 ± 24.1 (2–142)
October 1999	275.5 ± 20.8 (203–391)	27.2 ± 0.4 (25–29)	5/7	71.4	171.0 ± 45.8 (63–288)	3/3	100.0	42.3 ± 32.9 (8–108)
December 1999	704.9 ± 28.1 (275–2500)	32.3 ± 2.7 (27–50)	2/2	100.0	11.0 ± 1.0 (10–12)	2/2	100.0	221.5 ± 26.8 (195–248)
February 2000	212.0 ± 315.6 (13–638)	18.9 ± 13.1 (10–36)	1/1	100.0	231.0	1/1	100.0	223.0
April 2000	384.3 ± 158.5 (180–765)	27.7 ± 3.1 (22–35)	0/0	0	0	3/3	100.0	404.0 ± 314.2 (65–1031)
June 2000	697.0 ± 91.3 (30–1034)	34.3 ± 2.5 (14–43)	1/4	25.0	25.0 ± 5.6	4/5	80.0	13.7 ± 6.0 (3–32)
August 2000	1781.3 ± 222.8 (165–2750)	44.9 ± 2.5 (23–54)	1/5	20.0	5.0 ± 1.0	3/8	37.5	9.0 ± 2.8 (2–23)
October 2000	768.0 ± 194.5 (168–1503)	38.9 ± 3.5 (23–49)	2/4	50.0	169.5 ± 59.0 (26–313)	4/4	100.0	259.0 ± 96.6 (4–629)
December 2000	978.5 ± 194.1 (470–2500)	37.9 ± 2.3 (30–53)	7/7	100.0	97.7 ± 28.5 (16–213)	4/4	100.0	204.0 ± 74.8 (87–422)
February 2001	1,174.2 ± 187.1 (205–2200)	40.0 ± 2.2 (26–52)	5/6	83.3	22.1 ± 6.3 (3–39)	4/4	100.0	18.0 ± 4.1 (11–30)
April 2001	226.3 ± 57.8 (94–710)	25.0 ± 1.7 (19–37)	6/7	85.7	18.8 ± 3.2 (12–25)	1/2	50.0	234.0 ± 118.1
June 2001	716.3 (290–1015)	34.7 (27–41)	6/6	100.0	35.8 ± 13.4 (7–98)	4/4	100.0	38.5 ± 11.4 (11–65)
Total mean*	676.6 ± 135.1	32.2 ± 2.2	37/50**	75.9 ± 9.3	76.6 ± 23.9	39/46**	88.9 ± 6.4	145.2 ± 36.7

showed the highest MI ( $P < 0.05$ ) when compared with males examined in the same months (fig. 1D).

## Discussion

### *Infection levels and host body weight*

Species of *Cichla* are native to the Amazonian Rivers (Gomiero & Braga, 2003a, 2004a, b), but have been introduced in several localities (Hoeinghaus *et al.*, 2003). In the Volta Grande Reservoir, Minas Gerais, species of *Cichla* are alien and their strong voracity affects the other small fish including the same fish species (Gomiero & Braga, 2004a, b). For this reason, predation might be the main cause of elevated parasitological indexes (Martins *et al.*, 2009). Among the factors that contribute to disease dissemination in fish are host susceptibility, fish stocking densities, the duration of the infectious process, the introductions and inadequate handling, as mentioned by Reno (1998). Consequently, the trophic relationship of this alien fish in the reservoir should be better understood to explain the elevated parasitological indexes found by Martins *et al.* (2009); they are discussed below regarding the sex and weight of the host and parasitism.

The diet variations among *Cichla* species are related to food availability, their length (ontogeny) and seasonality (Rabelo & Araújo-Lima, 2002; Gomiero & Braga, 2004b). No difference was related to feeding and condition factor between the sexes in *Cichla* spp. (Gomiero & Braga, 2003b). However, there are no studies on the relationship between parasitological indexes in males and females in that region. In the studies of Martins *et al.* (2009), general parasitological indexes in *C. piquiti* were discussed.

This new information on the relationship between the sex and weight of the hosts and proteocephalidean infection in tucunaré from this reservoir contributes to the knowledge of the host-parasite-environment relationships. The intensity of infection associated with increased body size shown by Moravec *et al.* (2002) differs from our results. In the present case, *C. piquiti* weighing 301–800 g showed the highest MI of infection, probably related to the fact that fish of lower weight might eat invertebrates (Gomiero & Braga, 2004a) that are intermediate hosts of proteocephalideans (Scholz, 1999). In contrast to the present study, Moravec *et al.* (2002) did not observe seasonal variation. A 100% prevalence in *C. piquiti* was also reported in fish parasitized by *Proteocephalus chamelensis* (Pérez Ponce de León *et al.*, 1995) and by *S. maringaensis* (De Chambrier & Vaucher, 1999; Arredondo & Gil de Pertierra, 2008). According to Scholz (1987), larger perch were the most parasitized by *Proteocephalus percae* Müller, 1780. Also studying the parasitic fauna of fish from lakes, Molloy *et al.* (1995) reported low infection by cestodes in trout, in contrast to our results. Contrary to the present results, Zelman & Arai (1998) reported a significant increase in the plerocercoid abundance in female perch.

When analysing the length of the hosts parasitized by *S. maringaensis*, a positive correlation between intensity of infection and standard length was observed (Takemoto & Pavanelli, 2000). A correlation between total length and



prevalence was also verified in *C. monoculus* weighing 53–803 g parasitized by proteocephalideans (Machado *et al.*, 2000). These results differ from ours in that fish from 301–800 g showed the highest MI of infection.

The cumulative effect of parasitism with regard to host length, suggested by Machado *et al.* (2000) in fish from the same genus, was not clear in our material. The frequency of distribution of proteocephalideans in *C. piquiti* was more important between 1 and 200 parasites. A case of elevated intensity of cestode infection (1031 helminths) was reported in April 2000, but this phenomenon was not common, as supported by Martins *et al.* (2009). On the other hand, Valtonen & Koskivaara (1994) reported that the MI of infection was higher for wild than for farmed salmon and trout.

It could be possible that the highest MIs observed in October and December (fig. 1B) are related to the greater age of the fish. In these months, fish presented total lengths between 23 and 53 cm, which would correspond to the ages of 2–6 years, according to Gomiero & Braga (2007), in *Cichla* spp. from the Volta Grande reservoir. As discussed by Gomiero & Braga (2007), the study of *Sagitta* can offer only preliminary information to characterize fish length. In this study, it is not possible to know the exact age of the fish, providing only length data as supported by Gomiero & Braga (2007). Apart from the possible interference of reproductive stage (Gomiero & Braga, 2003a), fish weight is a variable parameter when measured in fish in which differences may be present between the sexes.

#### *Infection levels and host gender*

Despite the large sample of males and females in the present study, there was no significant association between sex and prevalence of parasitism, as also supported by Takemoto & Pavanelli (2000). These results are supported by Gomiero & Braga (2003b), who did not find significant differences in the condition of males and females of *Cichla* spp. in the same reservoir. They argued that fish diet is not influenced by the sex of the host.

The greatest MIs of infection in October and December in both sexes of *C. piquiti* when analysed together (fig. 1B) may be explained by the highest MI in males in October and females in December (fig. 1D). These numbers coincide with the highest rainfalls (669–1617 mm) and possibly water temperature (25.33–27.83°C), as discussed by Martins *et al.* (2003). It can be clearly observed in table 2 that from December to April 1999 and December to April 2001 (except February 2001) both fish sexes showed a high parasite load. As mentioned by Gomiero & Braga (2004a, b), this could be explained by the immature population of *Cichla* occurring mainly in this period.

Moreira *et al.* (2005) observed parasites only in females of *Iheringichthys labrosus* from the upper Paraná floodplain, in contrast to our results in that no significant difference was found between parasitized males and females. Moreira *et al.* (2005) suggested that the number of fish captured must be larger. The quantity of fish used in the present study was sufficient for statistical analysis and the number of males and females was very similar.

However, we cannot reject the possibility of individual variation.

In contrast to our observations on *C. piquiti* in the Volta Grande Reservoir, Takemoto & Pavanelli (2000) reported that male *Sorubim lima* hosts were the most parasitized sex. However, the evaluation of Machado *et al.* (1994) indicated no relationship between host sex and parasitism. Our results show that females have higher prevalence rates than males, but without a significant difference between the sexes.

In marine fish, Luque *et al.* (1996) observed that seven parasite species had a positive correlation between intensity of infection and total length of the host, but only digeneans showed a relationship between intensity and sex of the host. In the present study, fish with weight of 301–800 g showed the highest MI.

Supporting this view, Martins *et al.* (2003, 2009) emphasized the influence of aquatic characteristics and rainfall on prevalence and MI of infection in fish from the Volta Grande Reservoir. It must be emphasized that the studied environment constitutes a reservoir where fish may be more susceptible to intermediate hosts than fish from the rivers.

#### Acknowledgements

We thank the authorities of the Hydroelectric Power Station of Minas Gerais (CEMIG), Brazil for field support and the National Council for Scientific and Technological Development (CNPq) for a grant to the first author.

#### References

- Arredondo, N. & Gil de Pertierra, A.A. (2008) The taxonomic status of *Spatulifer* cf. *maringaensis* Pavanelli & Rego, 1989 (Eucestoda: Proteocephalidea) from *Sorubim lima* (Bloch & Schneider) (Pisces: Siluriformes), and the use of the microthrix pattern in the discrimination of *Spatulifer* spp. *Systematic Parasitology* **70**, 223–236.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. & Shostak, W. (1997) Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* **83**, 575–583.
- De Chambrier, A. & Vaucher, C. (1999) Proteocephalidae et Monticelliidae (Eucestoda: Proteocephalidea) parasites de poissons d'eau douce du Paraguay avec descriptions d'un genre nouveau et de dix espèces nouvelles. *Revue Suisse de Zoologie* **106**, 165–240.
- Freze, V.I. (1965) *Proteocephalata in Fish, Amphibians and Reptiles. Essentials of Cestodology*, vol. V. 538 pp. Moscow, Nauka (in Russian; English translation, Israel Program of Scientific Translation (1969) Cat. No. 1853. v + 597 pp.).
- Gomiero, L.M. & Braga, F.M.S. (2003a) Pesca experimental do Tucunaré, gênero *Cichla* (Osteichthyes, Cichlidae), no reservatório da UHE de Volta Grande, Rio Grande (48°25'47"35' W, 19°57'20"10' S). *Boletim do Instituto de Pesca* **29**, 29–37.
- Gomiero, L.M. & Braga, F.M.S. (2003b) Relação peso-comprimento e fator de condição para *Cichla* cf. *ocellaris* e *Cichla monoculus* (Perciformes, Cichlidae)

- no reservatório de Volta Grande, rio Grande – MG/SP. *Acta Scientiarum* **25**, 79–86.
- Gomiero, L.M. & Braga, F.M.S.** (2004a) Cannibalism as the feeding behaviour of Tucunarés introduced in Southeast Brazil. *Brazilian Journal of Biology* **64**, 625–632.
- Gomiero, L.M. & Braga, F.M.S.** (2004b) Feeding of introduced species of *Cichla* (Perciformes, Cichlidae) in Volta Grande reservoir, River Grande (MG/SP). *Brazilian Journal of Biology* **64**, 787–795.
- Gomiero, L.M. & Braga, F.M.S.** (2004c) Reproduction of species of the genus *Cichla* in a reservoir in South-eastern Brazil. *Brazilian Journal of Biology* **64**, 613–624.
- Gomiero, L.M. & Braga, F.M.S.** (2007) Description of the otoliths of peacock bass (*Cichla* sp. and *Cichla monoculus*) in the reservoir of Volta Grande dam (SP–MG). *Ciencia Animal Brasileira* **8**, 119–126.
- Hoinghaus, D.J., Layman, C.A., Arrington, D.A. & Winemiller, K.O.** (2003) Movement of *Cichla* species (Cichlidae) in a Venezuelan floodplain river. *Neotropical Ichthyology* **1**, 121–126.
- Luque, J.L., Amato, J.F.R. & Takemoto, R.M.** (1996) Comparative analysis of the communities of metazoan parasites of *Orthopristis ruber* and *Haemulon stendachneri* (Osteichthyes: Haemulidae) from the Southeastern Brazilian Littoral: I. Structure and influence of the size and sex of hosts. *Revista Brasileira de Biologia (Brazilian Journal of Biology)* **56**, 279–292.
- Machado, M.H., Pavanelli, G.C. & Takemoto, R.M.** (1994) Influence of host's sex and size on endoparasitic infrapopulations of *Pseudoplatystoma corruscans* and *Schizodon borelli* (Osteichthyes) of the High Paraná River, Brazil. *Revista Brasileira de Parasitologia Veterinaria* **3**, 143–148.
- Machado, P.M., Almeida, S.C., Pavanelli, G.C. & Takemoto, R.M.** (2000) Ecological aspects of endohelminths parasitizing *Cichla monoculus* Spix, 1831 (Perciformes: Cichlidae) in the Paraná River near Porto Rico, State of Paraná, Brazil. *Comparative Parasitology* **67**, 210–217.
- Martins, M.L., Fenerick, J. Jr, Nomura, D.T. & Braga, F.M.S.** (2003) Prevalence, intensity and seasonality of *Ichthyouris voltagrandsensis* (Nematoda: Pharyngodonidae) in *Myleus tiete* (Osteichthyes: Characidae) from Volta Grande Reservoir, Minas Gerais, Brazil. *Boletim do Instituto de Pesca* **29**, 39–46.
- Martins, M.L., Pereira, J. Jr, de Chambrier, A. & Yamashita, M.M.** (2009) Proteocephalid cestode infection in alien fish, *Cichla piquiti* (Osteichthyes: Cichlidae), from Volta Grande Reservoir, Minas Gerais, Brazil. *Brazilian Journal of Biology* **69**, 631–637.
- Molloy, S., Holland, C. & Poole, R.** (1995) Metazoan parasite community structure in brown trout from two Lakes in Western Ireland. *Journal of Helminthology* **69**, 237–242.
- Moravec, F., Mendoza-Franco, E., Vivas-Rodríguez, C., Vargas-Vázquez, J. & González-Solís, D.** (2002) Observations on seasonal changes in the occurrence and maturation of five helminth species in the pimelodid catfish, *Rhamdia guatemalensis*, in the cenote (=sinkhole) Ixin-há, Yucatán, Mexico. *Acta Societatis Zoologicae Bohemicae* **66**, 121–140.
- Moreira, S.T., Ito, K.F., Takemoto, R.M. & Pavanelli, G.C.** (2005) Ecological aspects of the parasites of *Iheringichthys labrosus* (Lütken, 1874) (Siluriformes: Pimelodidae) in reservoirs of Paraná basin and upper Paraná floodplain, Brazil. *Acta Scientiarum Biological Sciences* **27**, 317–322.
- Pérez-Ponce de León, G., Brooks, D.R. & Berman, R.** (1995) *Proteocephalus chamelensis* n. sp. (Cestoda: Proteocephalidae) from the 'Guavina' *Gobiomorus maculatus* (Osteichthyes: Eleotrididae) in Chamela bay, Jalisco, México. *Journal of Parasitology* **81**, 773–776.
- Rabelo, H. & Araújo-Lima, C.A.R.M.** (2002) A dieta e o consumo diário de alimento de *Cichla monoculus* na amazônia central. *Acta Amazonica* **32**, 707–724.
- Rego, A.A., Chubb, J.C. & Pavanelli, G.C.** (1999) Cestodes in South American freshwater teleost fishes: keys to genera and brief description of species. *Revista Brasileira de Zoologia* **16**, 299–367.
- Reno, P.W.** (1998) Factors involved in the dissemination of disease in fish populations. *Journal of Aquatic Animal Health* **10**, 160–171.
- Schmidt, G.D.** (1986) *CRC Handbook of tapeworm identification*. 675 pp. Boca Raton, Florida, CRC Press.
- Scholz, T.** (1987) On the occurrence of endoparasitic helminthes in perch (*Perca fluviatilis* L.) from the Mácha lake fishpond system, Czechoslovakia. *Acta Universitatis Carolinae – Biologica* **11**, 301–318.
- Scholz, T.** (1999) Life cycles of species of *Proteocephalus*, parasites of fishes in the Palearctic Region: a review. *Journal of Helminthology* **73**, 1–19.
- Takemoto, R.M. & Pavanelli, G.C.** (2000) Aspects of the ecology of proteocephalid cestodes parasites of *Sorubim lima* (Pimelodidae) of the upper Paraná river, Brazil. I. Structure and influence of host's size and sex. *Revista Brasileira de Biologia (Brazilian Journal of Biology)* **60**, 577–584.
- Valtonen, E.T. & Koskivaara, M.** (1994) Relationship between the parasites of some wild and cultured fishes in two lakes and a fish farm in Central Finland. *International Journal for Parasitology* **24**, 109–118.
- Zelmer, D.A. & Arai, H.P.** (1998) The contributions of host age and size to the aggregated distribution of parasites in yellow perch, *Perca flavescens*, from Garner Lake, Alberta, Canada. *Journal of Parasitology* **84**, 24–28.