

Significance of Dopamine Transmission in the Rat Medial Prefrontal Cortex for Conditioned Fear

Marie A. Pezze^{1*}, Tobias Bast^{2*} and Joram Feldon

Laboratory of Behavioral Neurobiology, Swiss Federal Institute of Technology Zurich (ETH), Switzerland

*Both authors contributed equally to the present study

Present address: ¹Department of Experimental Psychology, University of Cambridge, Downing Street, Cambridge CB2 2EB, UK

Present address: ²Department of Neuroscience, University of Edinburgh, 1 George Square, Edinburgh EH8 9JZ, UK

Previous studies have demonstrated activation of dopamine transmission in the medial prefrontal cortex (mPFC) by conditioned fear stimuli. Therefore, the present study investigated the functional significance of mPFC dopamine for a conditioned fear response to a tone. We examined the effects of inhibition or stimulation of mPFC dopamine transmission by local microinfusion of the D1/D2-receptor antagonist *cis*-flupenthixol or the indirect dopamine receptor agonist *D*-amphetamine, respectively, in a classical fear-conditioning paradigm in Wistar rats. Rats received tone–shock pairings and were later exposed to the tone alone. Freezing was used as measure of conditioned fear. Presence of the drugs in the mPFC during the tone–shock pairings did not affect freezing during subsequent presentation of the tone alone. However, when *cis*-flupenthixol and *D*-amphetamine were present in the mPFC during presentation of the tone alone, freezing to the tone was reduced. We demonstrated that the decreased freezing could be explained neither by state dependency nor infusion-induced alterations in activity. Our data indicate that mPFC dopamine transmission is important for the retrieval/expression, but not the formation, of conditioned fear. The reduction of conditioned fear by prefrontal infusion of both *cis*-flupenthixol and *D*-amphetamine may reflect normal expression of conditioned fear requires an optimal level of mPFC dopamine activity.

Introduction

In rats and humans, the medial prefrontal cortex (mPFC) is characterized by similar connections and has been suggested to play a pivotal role in emotional responses (Nauta, 1971; Davidson and Irwin, 1999; Bechara *et al.*, 2000; Öngür and Price, 2000; Simpson *et al.*, 2000). Research on emotion in laboratory rodents has focused on fear, the neural substrates of which can readily be studied in classical fear-conditioning experiments (LeDoux, 2000). In classical fear conditioning, an emotionally neutral conditioned stimulus (CS), such as a tone, is paired with an aversive unconditioned stimulus (US), usually a foot shock, so that subsequent presentation of the CS elicits conditioned fear responses.

Conditioned fear is accompanied by marked alterations in mPFC neurotransmission. In particular, neurochemical studies in rats have consistently suggested activation of mPFC dopamine transmission by presentation of a CS previously paired with foot shocks [(Goldstein *et al.*, 1994, 1996; Inoue *et al.*, 1994; Yoshioka *et al.*, 1996; Morrow *et al.*, 1999b,c; Feenstra *et al.*, 1999, 2001; Feenstra, 2000), but see Wilkinson *et al.* (Wilkinson *et al.*, 1998)]. This activation was proposed to be necessary for the normal expression (Morrow *et al.*, 1999c) or extinction of fear responses to a CS (Morrow *et al.*, 1999b; Feenstra *et al.*, 2001). Moreover, specific patterns of electrical activity (García *et al.*, 1999; Baeg *et al.*, 2001) and increased extracellular norepinephrine (Feenstra *et al.*, 1999, 2001) were evoked in the mPFC by fear CS.

The functional significance of the reported alterations in

mPFC neurotransmission for a normal conditioned fear response is not clear. Currently, a popular view is that processes in the mPFC are important for inhibitory control of conditioned fear (Maren, 2001). This view is based on the findings that lesions to the dorsal (mainly anterior cingulate) mPFC enhanced acquisition of conditioned fear (Morgan and LeDoux, 1995; Vouimba *et al.*, 2000) and lesions to the ventral mPFC impaired fear extinction (Morgan *et al.*, 1993; Quirk *et al.*, 2000). Importantly, destruction of the most caudal and ventral infralimbic mPFC was demonstrated to be critical for the extinction deficit (Quirk *et al.*, 2000). Based on the finding that 6-hydroxydopamine lesions of the mPFC, depleting the local catecholamine, in particular dopamine, innervation, reduced extinction of very strong, but not moderate, conditioned fear, mPFC dopamine was proposed to be important for fear extinction under certain conditions (Morrow *et al.*, 1999b). There is also strong evidence that some subregions of the mPFC are important for the normal expression of conditioned fear. Thus, in line with Nauta's (Nauta, 1971) earlier proposition that frontal cortex lesions result in an 'interoceptive agnosia', reflected by an absence of normal emotional responses, lesions destroying the central mPFC, including prelimbic and infralimbic parts, but sparing the most caudal and ventral infralimbic portions, reduced the expression of conditioned fear (Fryszak and Neafsey, 1991, 1994; Lacroix *et al.*, 2000b; McLaughlin *et al.*, 2001). Finally, there are several studies (Holson, 1986; Rosen *et al.*, 1992; Gewirtz *et al.*, 1997) in which mPFC lesions, destroying considerable parts of the prelimbic and infralimbic subregions, did not affect conditioned fear responses. This may be related to compensatory processes induced by permanent lesions (Bast *et al.*, 2001a). Altogether, the specific functional significance of prefrontal processes accompanying conditioned fear, the best documented of which is activation of dopamine transmission, awaits clarification.

In contrast to permanent lesions, local microinfusions of neuroactive substances allow the role of different transmitter systems in the mPFC during different stages of a fear-conditioning experiment to be addressed specifically. In order to clarify the functional significance of alterations in mPFC dopamine transmission for aversively conditioned emotional responses, the present study examined the effects of specifically manipulating mPFC dopamine transmission by local microinfusion of the dopamine D1/D2-receptor antagonist *cis*-flupenthixol or the indirect dopamine agonist *D*-amphetamine in a classical fear-conditioning paradigm in Wistar rats. Freezing, an immobile posture in response to an inescapable aversive CS (Fanselow, 1984), was used as measure of conditioned fear.

Materials and Methods

Animals

One hundred male adult Wistar rats (Zur:Wist[HanIbm]; Research Unit Schwerzenbach, Schwerzenbach, Switzerland), ~2 months old and

weighing ~250–300 g at the time of surgery, were used for the experiments of the present study (experiment I, 24 rats; experiment II, 39; experiment III, 37). The animals were housed under a reversed light–dark cycle (lights on 19:00–07:00) in a temperature ($21 \pm 1^\circ\text{C}$) and humidity ($55 \pm 5\%$) controlled room and were provided with food and water *ad libitum*. All rats received bilateral implantation of guide cannulae aiming at the mPFC. Before surgery, rats were housed in groups of four per cage; after surgery, they were individually caged. Beginning 3 days before surgery and thereafter until the beginning of the behavioral experiments, all rats were handled daily. All experimental procedures were carried out in the dark phase of the cycle. Principles of laboratory animal care (NIH publication No. 86-23, revised 1985) and Swiss regulations for animal experimentation were followed.

Implantation of Guide Cannulae for Intracerebral Infusion

Rats were anesthetized with 1 ml of Nembutal (sodium pentobarbital, 50 mg/ml; Abbott Labs, North Chicago, IL) per kg body wt and their heads were placed in stereotaxic frames (Kopf Instruments, Tujunga, CA). After application of a local anesthetic (lidocaine), the scalp was incised to expose the skull, and bregma and lambda were aligned in the same horizontal plane. A pair of guide cannulae (9 mm, 26 gauge, stainless steel) in a Perspex holder (custom made) were implanted through small holes (1.5 mm diameter) drilled on each side of the skull. The tips of the guide cannulae aimed at the following coordinates (in mm): 3.0 anterior and ± 0.5 lateral to bregma and 3.1 ventral to skull surface. The guide cannulae were fixed with dental cement for which three small stainless screws, previously screwed into the skull, served as anchors. Stainless steel stylets (34 gauge) which extended 0.5 mm beyond the tips of the guide cannulae were placed inside the guide cannulae to prevent occlusion. After surgery, the experimenters gave the rats daily health checks and gentle handling, and replaced missing stylets. The behavioral experiments commenced 5 days after surgery.

Intracerebral Microinfusion and Drugs

For microinfusions into the mPFC, rats were manually restrained and the stylets removed from the guide cannulae. Infusion cannulae (34 gauge, stainless steel, custom made) were then inserted into the guide cannulae. The infusion cannulae were fixed to 26 gauge cuffs at one end to prevent them from slipping through the guide cannulae and connected to 10 μl Hamilton microsyringes mounted on a microinfusion pump (KD Scientific or WPI sp200i). The tips of the infusion cannulae protruded into the mPFC 1.5 mm beyond the tip of the guide cannulae, thus aiming at a final coordinate of 4.6 mm below the skull surface. The rats were bilaterally infused with *cis*-flupenthixol dihydrochloride (FLU, 20 or 25 μg , $\text{C}_{23}\text{H}_{25}\text{F}_3\text{N}_2\text{O}_5 \times 2\text{HCl}$; Sigma, Switzerland) or *D*-amphetamine sulfate [AMPH, 13.6 μg (10 μg free base), $\text{C}_{18}\text{H}_{26}\text{N}_2 \times \text{H}_2\text{SO}_4$; Sigma, Switzerland] in 0.5 μl vehicle (0.9% saline) or with 0.5 μl vehicle (VEH) only. Drug solutions were freshly prepared on the day of infusion. The infusion speed was 0.5 $\mu\text{l}/\text{min}$. After infusion, the infusion cannulae were left in the brain for 60 s to allow for absorption of the infusion bolus by the brain tissue and then replaced by the stylets. Rats were then immediately subjected to the behavioral testing.

AMPH and FLU were chosen for this first examination of the functional significance of mPFC dopamine transmission for conditioned fear because these compounds affect both D1- and D2-receptor-mediated processes. Furthermore, we could draw on the experience from previous experiments performed in our laboratory (Lacroix *et al.*, 2000a) in which these drugs had been infused into the mPFC. While the indirect dopamine agonist AMPH may also stimulate the release of other monoamines (Azzaro and Rutledge, 1973), the direct nonselective dopamine agonist apomorphine also affects transmission by other monoamines; for example, apomorphine has similar affinities to dopamine and α_1 -adrenoceptors (Baldessarini *et al.*, 1994). AMPH was infused into the mPFC at a dose of 13.6 μg (corresponding to 10 μg of the free base)/0.5 $\mu\text{l}/\text{side}$ in the present experiments because this dose produced a behavioral effect in a previous study from our laboratory (Lacroix *et al.*, 2000a). With regard to FLU, 12 $\mu\text{g}/0.5 \mu\text{l}/\text{side}$ infused into the mPFC did not yield any behavioral effect in the aforementioned study (Lacroix *et al.*, 2000a) and so a dose of 20 $\mu\text{g}/0.5 \mu\text{l}/\text{side}$ was used in the first fear-conditioning experiment (experiment I). Since this dose yielded a behavioral effect without any adverse side effect (e.g. disrupted motor

coordination, brain damage), a slightly higher dose of 25 $\mu\text{g}/0.5 \mu\text{l}/\text{side}$ was used in the final fear conditioning experiment (experiment III) in order to obtain the behavioral effect reliably.

Two experiments were conducted to control for the low pH (2–3) of the FLU solutions. Subjects were tested for the expression of conditioned fear and for activity following infusion of acidified (adjusted to pH 2 by concentrated hydrochloric acid, $n = 5$), as detailed below (experiment I, activity experiment). No significant behavioral effects were observed relative to rats infused into the mPFC with normal saline (pH 6–7, $n = 5$, data not shown). Therefore, subsequent comparisons were made between rats infused into the mPFC with FLU solutions (pH 2–3) and subjects receiving mPFC infusion of normal saline (pH 6–7).

Apparatus for Behavioral Testing

Eight operant test boxes, four shock boxes and four no-shock boxes (Habitest; Coulbourn Instruments, Allentown, PA) were used for the fear-conditioning experiments (experiments I–III). Shock boxes were used for conditioning and context-test sessions, while the no-shock boxes were used to assess fear to the tone CS in an environment distinct from that during conditioning. Shock boxes were fitted with a parallel grid shock floor (16 parallel bars, E10-10RF; Coulbourn Instruments), through which scrambled shocks could be delivered, and placed in light- and sound-attenuating chambers measuring $55 \times 40 \times 55$ cm (wide \times long \times high). These chambers had two side walls of aluminium and rear and front walls of clear Perspex. A white waste tray was situated below the grid floor. The four no-shock boxes were fitted with a lattice grid (E10-18NS; Coulbourn Instruments) and placed in light- and sound-attenuating chambers measuring $72 \times 45 \times 45$ cm. They had three black walls and a front wall of clear Perspex. A brown waste tray was situated below the lattice grid. The four shock and the four no-shock boxes were placed in two different rooms. Presentation of the tone CS and delivery of electric foot shock were controlled by a PC with dedicated software (S. Frank, Psychology Department, University of Tel Aviv, Israel) connected to a Coulbourn Universal Environment Interface (E91-12) with Coulbourn Universal Environment Port (L91-12). The tone CS [85 dB(A)] was produced by a 2.9 kHz tone module (E12-02) fixed at one wall of the operant chamber. Shocks were delivered with a Coulbourn Precision Animal Shocker (E13-12), which generated bipolar rectangular 10 ms current pulses with a frequency of 10 Hz. Background noise was provided by a ventilation fan affixed to the light- and sound-attenuating chambers during all sessions. A monochrome minivideo camera with a wide angle (100°) 2.5 mm lens (VPC-465B; CES AG, Zurich, Switzerland) was attached to the center of the ceiling of each operant chamber. Four infrared (875 nm) light-emitting diodes (HSDL-4220; Hewlett Packard) positioned in the ceiling of each operant chamber provided light sufficient for camera function. Throughout all sessions, images from the test boxes were provided by these cameras, integrated into a four-quarter single image (100 000 pixels) by a multiplexer (DX216CE; Sony) and recorded by a video-recorder (SVT1000; Sony).

Sixteen cubicles ($25 \times 40 \times 40$ cm), each contained within an individual sound-attenuating wooden cabinet, were used for the activity experiment examining the effects of FLU and AMPH infusion into the mPFC on general activity. The cubicles had three walls of clear plastic and one side wall of wooden panels. The floor of each cubicle was a black removable pan holding a thin layer of dark, absorbent, autoclaved earth. The ceiling was open. A fan mounted on the wall of each cabinet provided ventilation. A monochrome minivideo camera with a wide angle (100°) 2.5 mm lens (VPC-465B; CES AG, Zurich, Switzerland), centered ~49 cm above the compartment floor, was mounted in the ceiling of each cabinet. Six infrared (875 nm) light-emitting diodes (HSDL-4220; Hewlett Packard) positioned in the ceiling of each cabinet provided light sufficient for camera function. Images provided from each of the 16 cubicles by these cameras were integrated into a sixteen-part single image (400 000 pixels) by a 16-channel multiplexer (YS-DX216CE; Sony) and the image was recorded by a video-recorder (SVT1000; Sony).

Automated Measurement of Activity and Freezing

The video images were transferred to a computer (fear-conditioning experiments, 7600/120 Power Macintosh; control experiment, Dell OptiPlex GXpro with a Pentium Pro Processor) equipped with an analysis program (Image, <http://rsb.info.nih.gov/nih-image>) and a macroprogram

(P. Schmid, Behavioral Neurobiology Laboratory, Swiss Federal Institute of Technology Zurich). The percentage of changed pixels between two adjacent 1 s images from one of the test boxes was used as a measure of activity (Richmond *et al.*, 1998). Freezing is commonly identified as cessation of any movement except for respiratory movements (Anagnostaras *et al.*, 1999). If the percentage of changed pixels between two adjacent 1 s images was <0.05%, this corresponded well to such immobility and the behavior of the rat was scored as 'freezing' for the respective second. Validation and principle of the automated analysis of freezing behavior have been described in detail in previous publications (Richmond *et al.*, 1998; Pryce *et al.*, 1999).

Fear-conditioning Experiments

Three fear-conditioning experiments (experiments I-III) were conducted testing the effects of FLU or AMPH infusion into the mPFC on the formation, retrieval/expression and extinction of fear conditioning to a tone. In all three experiments there were three types of sessions: conditioning, context-test and tone-test sessions. Conditioning sessions consisted of two (experiment I) or six (experiments II and III) tone-shock pairings (30 s tone coterminating with a 1 s, 0.5 mA foot shock) preceded and followed by 2 min (experiment I) or 1.5 min blocks (experiments II and III) in the shock boxes. Context-test sessions consisted of 8 min during which the rats were placed in the shock boxes without receiving any stimulation. Context-test sessions were conducted to assess long-term conditioned fear (i.e. conditioned fear persisting beyond the conditioning session) to the context in which the foot shocks were experienced, as well as for extinction of fear to the context, in order to obtain a pure measure of fear to the tone during the subsequent tone test. Tone tests, to assess long-term conditioned fear to the tone, were conducted in the no-shock boxes – i.e. an environment distinct from the conditioning context – and consisted of a continuous 8 min tone presentation, preceded by 2 min without stimulation. During all sessions, freezing was assessed by the automated analysis system as a measure of conditioned fear. In those experiments involving drug infusions before conditioning (experiments II and III), the effects of the drug infusions on unconditioned behaviors were assessed by observing some unconditioned behaviors in the different infusion groups during the conditioning session. The video images taken during conditioning were observed in order to determine whether all infusion groups reacted to the shock by similar vigorous twitching and jumping – marked components of the unconditioned immediate shock response (Anagnostaras *et al.*, 1999). Activity before the first tone or shock was assessed using the percentage of changed pixels between adjacent 1 s video images given by the automated system. The different infusion groups, as well as conditioning and test procedures, in experiments I, II and III are described in detail below and summarized in Table 1.

Experiment I

In experiment I, rats received bilateral infusions of VEH ($n = 8$), AMPH (13.6 μg , $n = 8$), or FLU (20 μg , $n = 8$) into the mPFC immediately before the tone-test session. The three infusion groups were matched for freezing levels during conditioning and the context test. The conditioning session consisted of two tone-shock pairings. Using a low number of pairings, the aim was to achieve a relatively low level of conditioned freezing to the tone in order to facilitate detection of an extinction deficit. Context-test sessions were conducted 1 day after conditioning, tone-test sessions 2 days after conditioning.

Experiments II and III

Experiment I indicated that infusions of FLU as well as AMPH before tone test resulted in reduced conditioned freezing to the tone, suggesting that both treatments interfere with the retrieval/expression of conditioned fear. Experiments II and III were aimed at confirming this suggestion. In particular, we tested whether the reduced conditioned freezing during tone test might merely have reflected state dependency, i.e. that associations formed in an altered brain-state may subsequently be retrieved only in a similar brain-state (Overton, 1964), rather than an impairment of the specific processes underlying retrieval/expression of conditioned fear. For that purpose, FLU (experiment II) and AMPH (experiment III) were infused not only immediately before the tone test,

Table 1

Summary of the procedures in the three fear-conditioning experiments (experiment I-III)

Experiment	Infusion	Conditioning (shock box)	Context test (shock box, 8 min)	Tone test (no-shock box, 10 min, tone on after 2 min)
Experiment I	Immediately before tone test: VEH, FLU (20 $\mu\text{g}/\text{side}$), AMPH (13.6 $\mu\text{g}/\text{side}$)	Two tone-shock pairings each preceded and followed by 2 min blocks (7 min)	One day after conditioning	Two days after conditioning
Experiment II	Immediately before conditioning and tone test: VEH, FLU (25 $\mu\text{g}/\text{side}$)	Six tone-shock pairings each preceded and followed by 1.5 min blocks (13.5 min)	Six days after conditioning	Seven days after conditioning
Experiment III	Immediately before conditioning and tone test: VEH, AMPH (13.6 $\mu\text{g}/\text{side}$)	As in experiment II	As in experiment II	As in experiment II

AMPH, *D*-amphetamine sulfate; FLU, *cis*-flupentixol dihydrochloride; VEH, vehicle. Tone: 30 s, 2.9 kHz, 85 dB(A). Shock: 1 s, 0.5 mA foot shock, concurrent with the last second of the tone. Conditioned fear, in the form of freezing, during conditioning and testing was automatically measured in the form of freezing.

but also before conditioning sessions. Thus, the results of experiments II and III are also relevant regarding the role of mPFC dopamine transmission in the formation of conditioned fear. In both experiments all rats received a first infusion of either VEH or drug immediately before conditioning and a second infusion immediately before the tone test. In experiment II, all rats received either bilateral infusions of VEH ($n = 19$) or FLU (25 μg ; $n = 20$) into the mPFC before conditioning. Immediately before the tone test all rats received a second infusion of VEH or FLU. Half of the rats received the same infusion as before conditioning and the other half received a different infusion, resulting in four groups differing with respect to the combinations of infusions (before conditioning-before tone test): VEH-VEH ($n = 9$); VEH-FLU ($n = 10$); FLU-VEH ($n = 10$); and FLU-FLU ($n = 10$). In experiment III, all rats received either bilateral infusions of VEH ($n = 18$) or AMPH (13.6 μg ; $n = 19$) into the mPFC before conditioning. Immediately before the tone test, all rats received a second infusion of VEH or AMPH. Half of the rats received the same infusion as before conditioning and the other half received a different infusion, resulting in four groups differing with respect to the combinations of infusions (before conditioning-before tone test): VEH-VEH ($n = 8$); VEH-AMPH ($n = 10$); AMPH-VEH ($n = 9$); and AMPH-AMPH ($n = 10$). In both experiment II and III, the groups that received the same infusion before conditioning, but different infusions before the tone test, were matched for freezing levels during the conditioning and context-test sessions.

For conditioning, rats were put in the shock boxes for a total of 13.5 min and received six tone-shock pairings. A higher number of pairings than in experiment I was used in order to facilitate the detection of infusion-induced reductions in conditioned freezing. Six days after conditioning, a context test was conducted to minimize residual contextual freezing during the subsequent tone test. The tone test was conducted 7 days after conditioning. The time span between conditioning and test sessions was chosen so that repeated infusions were 1 week apart in order to allow for a recovery of the brain tissue from disturbances that may result from one intracerebral infusion before applying the next one.

Activity Experiment

Twenty-three rats used in experiment I were subjected to an additional experiment, 2 weeks after termination of experiment I, in order to test whether FLU or AMPH in the mPFC induced alterations in general activity that might have influenced the freezing response (Bast *et al.*, 2001b). (In experiments II and III, unconditioned activity was monitored before the first shock during the conditioning sessions, i.e. during the first 1.5 min

following the infusions, but possible infusion-induced alterations in unconditioned general activity after these first 1.5 min could not be assessed.) On day 1, basal activity was measured for 30 min. The animals were then allocated to one of three infusion groups, counterbalanced with respect to the infusions received during experiment I and matched according to their baseline activity. On day 2, rats received infusions of VEH ($n = 9$), FLU (20 $\mu\text{g}/\text{side}$, $n = 7$), or AMPH (13.6 $\mu\text{g}/\text{side}$, $n = 7$) immediately before their activity was monitored for 2 h. On day 3, activity was measured for 30 min without a preceding infusion.

Histology

After completion of the behavioral experiments, the cannulated rats were deeply anesthetized with an overdose of 2.5 ml/kg Nembutal (sodium pentobarbital, 50 mg/ml, i.p.) and transcardially perfused with 0.9% NaCl solution to rinse out the blood, followed by 250 ml of 4% formalin (4°C) to fix the brain tissue. After extraction from the skull, the brains were post-fixed in 4% formalin solution and subsequently cut into 40 μm coronal sections on a freezing microtome. For the verification of the infusion sites, every fifth section through the mPFC was mounted on a gelatine-treated slide and stained with cresyl violet. After staining, the sections were dehydrated and coverslipped. Subsequently, they were examined with a light microscope under $\times 10$ magnification. It was verified that the tips of the infusion cannulae were placed in the mPFC and their approximate locations were drawn onto plates from the atlas of Paxinos and Watson (Paxinos and Watson, 1998). Moreover, brain damage around the infusion sites was compared between the infusion groups.

Data Analysis

Statistical analysis was conducted using the Statview software system. From the freezing scores obtained for each second ('freezing' or 'not freezing'), the percentage of time spent freezing in a given time block was calculated. By averaging the percentage of pixels changed between adjacent 1 s video images in a given time block, a relative activity measure for this time block was calculated. Data were first subjected to analysis of variance (ANOVA), using the different infusions as between-subjects factor and the different time blocks of testing as repeated-measures factors. *Post hoc* comparisons were conducted using Fisher's protected least significant difference test. Level of significance was set at $P < 0.05$. Freezing during the context-test sessions is not reported, since it was negligible in all three experiments (on average $< 15\%$), reflecting that the conditioning procedures used in the present study did not result in considerable conditioned fear to the conditioning context. Data during the first four 1 min blocks of tone presentation served as a reliable measure of tone freezing that was not confounded by within-session habituation (Corcoran and Maren, 2001).

Results

Histology

In all rats used in experiments I-III, the tips of the infusion cannulae were located within or around the borders of the mPFC (Fig. 1). Therefore, behavioral data of all rats were included in the analysis. Damage was restricted to the area immediately surrounding the guide and infusion cannulae and did not differ between the infusion groups.

Experiment I: FLU and AMPH in the mPFC during Tone Test

In experiment I, in which rats received only one infusion of FLU, AMPH, or VEH immediately before the tone test, the three groups did not differ in their freezing levels during conditioning and context test, reflecting successful matching (data not shown). Freezing exhibited by the three different infusion groups during the tone test of experiment I is depicted in Figure 2.

Conditioned freezing to the tone previously paired with foot




Figure 1. Infusion sites in the medial prefrontal cortex. (A) Photomicrograph of a coronal brain section with the tracks of the guide cannulae and beneath them the infusion sites visible in both hemispheres. (B) Approximate location of the tips of the infusion cannulae depicted on plates of coronal sections through the rat brain (Paxinos and Watson, 1998); numbers on the right indicate distance from bregma in millimeters.




Figure 2. Freezing during tone-test sessions of experiment I. Infusions of VEH, FLU and AMPH were given immediately before the tone test. Mean percentage of time spent freezing during the tone test is depicted for the two 1 min blocks preceding the tone, as well as for the first four 1 min blocks of tone presentation. Bars represent two standard errors of the differences of means (2 SED), derived from the appropriate mean square of the ANOVA according to the published formula (Cochran and Cox, 1950). The 2 SED provide an estimate of population variance for between-groups comparisons and are used in line graphs for the sake of clarity.

shocks was lower in the AMPH and FLU groups than in the VEH rats. During the two 1 min blocks before tone onset, rats exhibited virtually no conditioned fear, as evidenced by low levels of immobility ($< 15\%$). ANOVA of freezing levels during the first four 1 min blocks of the tone presentation revealed a significant effect of infusion [$F(2,22) = 5.19$, $P < 0.02$]. *Post hoc* comparisons indicated that average freezing levels were lower in the AMPH ($P < 0.01$) and FLU ($P < 0.02$) rats than in the VEH group. AMPH and FLU rats did not differ ($P > 0.96$). Thus, the results of experiment I indicated that both AMPH




Figure 3. Effects of the different infusions on unconditioned activity during the 1.5 min before the first tone–shock pairing in experiments II and III. Infusions of VEH, FLU and AMPH were applied immediately before behavioral testing. Presented values are means + SEM.

and FLU in the mPFC reduced expression of conditioned fear. To corroborate this suggestion, we conducted experiments II and III.

Experiments II and III: FLU and AMPH in the mPFC during Conditioning and Tone Test

Unconditioned Behaviors during Conditioning Session

Figure 3 depicts unconditioned activity during the 1.5 min block preceding the first tone–shock pairing for the different infusion groups in experiments II and III.

FLU (experiment II) and AMPH (experiment III) rats did not differ significantly from VEH rats in the unconditioned activity exhibited throughout the 1.5 min block preceding the first tone–shock pairing at the beginning of the conditioning session. ANOVA of average activity throughout the 1.5 min preceding the first tone–shock pairing did not yield significant differences between FLU and VEH rats in experiment II [$F(1,37) = 1.75, P > 0.19$] or AMPH and VEH rats in experiment III [$F(1,35) = 0.65, P > 0.43$].

Inspection of the video images from the conditioning sessions of experiments II and III revealed that all infusion groups exhibited similar vigorous twitching and marked jumps as immediate response to the foot shock, indicating that FLU or AMPH infusions did not affect shock sensitivity.

Conditioned Freezing during Conditioning and Tone Test

The freezing data for conditioning and tone test of experiment II, in which rats received infusions of either FLU or VEH before conditioning, as well as before the tone test, are depicted in Figure 4.

During the conditioning session, the development of conditioned freezing in response to the inescapable foot shocks did not differ between rats that received FLU or VEH infusion before conditioning. ANOVA of average freezing levels during the seven 1.5 min blocks preceding and following the tone–shock pairings, as well as of the six 30 s blocks of tone presentations, did not yield an effect of the infusion received before conditioning [$F(1,37) = 0.84, P > 0.36$; $F(1,37) = 2.29, P > 0.13$] or an interaction of infusion \times time block [$F(6,222) = 0.47, P > 0.83$; $F(5,185) = 1.84, P > 0.10$]. There was only a significant effect of 1.5 min block [$F(6,222) = 56.31, P < 0.0001$] and 30 s tone presentation [$F(5,185) = 44.53, P < 0.0001$], reflecting the gradual increase of freezing with repeated shock administration.

During the tone test, the two groups that received FLU immediately before the tone test (VEH–FLU, FLU–FLU) exhibited less conditioned freezing to the tone than the groups that received VEH immediately before the tone test (VEH–VEH, FLU–VEH), while the infusion received before conditioning had




Figure 4. Freezing during conditioning and tone-test sessions of experiment II. A first infusion of VEH or FLU was given immediately before conditioning, resulting in two infusion groups for conditioning. A second infusion of VEH or FLU was given immediately before the tone test, resulting in four infusion groups (FLU–FLU; FLU–VEH; VEH–FLU; VEH–VEH), differing with respect to the combination of infusions received before conditioning and tone test. Mean percentage of time spent freezing during conditioning is depicted for the seven 1.5 min blocks preceding and following the six tone–shock pairings (A) as well as for the six 30 s blocks preceding and following the pairings (B). Mean percentage of time spent freezing during the tone test is depicted for the two 1 min blocks preceding the tone, as well as for the first four 1 min blocks of tone presentation (C). Bars represent two standard errors of the differences of means (2 SED).

no effect. Rats did not exhibit conditioned fear before tone onset, as indicated by the negligible freezing levels during the two 1 min blocks preceding tone onset ($<10\%$). Freezing strongly increased with tone onset. However, peak levels of freezing were considerably higher in the rats that received VEH before the tone test than in those that received FLU. This was reflected by the results of a three-way ANOVA (infusion before conditioning \times infusion before tone test \times 1 min block) of freezing during the first four 1 min blocks of tone presentation. Thus, there was an interaction of infusion before tone test \times 1 min block [$F(3,105) = 2.79, P < 0.05$] along with an effect of 1 min block [$F(3,105) = 8.22, P < 0.0001$], but neither a main effect of infusion before conditioning [$F(1,35) = 0.33, P > 0.56$] or of infusion before tone test [$F(1,35) = 2.57, P > 0.11$], nor interactions of infusion before conditioning \times 1 min block [$F(3,105) = 2.18, P > 0.09$] or of infusion before conditioning \times infusion before tone test \times 1 min block [$F(3,105) = 0.73, P > 0.53$]. Further analysis, based on the significant interaction of infusion before tone test \times 1 min block, revealed that groups receiving FLU before the tone test (VEH–FLU, FLU–FLU) exhibited less freezing during the second and third 1 min blocks




Figure 5. Freezing during conditioning and tone-test sessions of experiment III. A first infusion of VEH or AMPH was given immediately before conditioning, resulting in two infusion groups for conditioning. A second infusion of VEH or AMPH was given immediately before the tone test, resulting in four infusion groups (AMPH-AMPH; AMPH-VEH; VEH-AMPH; VEH-VEH), differing with respect to the combination of infusions received before conditioning and tone test. Mean percentage of time spent freezing during conditioning is depicted for the seven 1.5 min blocks preceding and following the six tone-shock pairings (A) as well as for the six 30 s tone presentations of the pairings (B). Mean percentage of time spent freezing during the tone test is depicted for the two 1 min blocks preceding the tone, as well as for the first four 1 min blocks of tone presentation (C). Bars represent two standard errors of the differences of means (2 SED).

of tone presentation than rats receiving VEH before the tone test (VEH-VEH, FLU-VEH). Thus, separate ANOVAs of freezing levels for each single 1 min block yielded a significant main effect of infusion before tone test for the second 1 min block [$F(1,35) = 5.05, P < 0.04$] and a strong tendency towards an effect for the third 1 min block [$F(1,35) = 3.67, P = 0.06$], while not indicating differences for the first [$F(1,35) = 0.06, P > 0.81$] and the fourth [$F(1,35) = 0.10, P > 0.75$] 1 min blocks.

The freezing data for conditioning and tone test of experiment III, in which rats received infusions of either AMPH or VEH before conditioning as well as before the tone test, are depicted in Figure 5.

During conditioning, freezing that developed in response to the foot shocks was lower in rats that received AMPH before conditioning than in those that received VEH. ANOVA of freezing during the seven 1.5 min blocks preceding and following the tone-shock pairings, as well as during the six 30 s blocks of tone presentations, yielded a main effect of infusion before conditioning [$F(1,35) = 6.09, P < 0.02$; $F(1,35) = 5.63, P < 0.03$] and 1.5 min block [$F(6,210) = 47.96, P < 0.0001$] or 30 s block




Figure 6. General activity after VEH, FLU and AMPH infusion in the activity experiment. Mean activity + SEM is depicted averaged over the complete 2 h (inset bar graph), as well as for each single of the first 15 1 min blocks (main line graph) following infusion. The bar in the main graph represents two standard errors of the differences of means (2 SED).

[$F(5,175) = 37.77, P < 0.0001$], respectively. There was also a strong tendency towards an interaction of infusion before conditioning \times 30 s block of tone presentation [$F(5,175) = 2.24, P = 0.0523$].

During the tone test, the two groups that received AMPH immediately before the tone test (VEH-AMPH, AMPH-AMPH) exhibited less conditioned freezing to the tone than the groups that received VEH immediately before the tone test (VEH-VEH, AMPH-VEH). The infusion received before conditioning had virtually no effect on conditioned freezing during the tone test. Rats did not exhibit conditioned fear before tone onset, as indicated by the negligible freezing levels during the two 1 min blocks preceding tone onset ($<5\%$). In all groups, freezing increased with tone onset. However, freezing reached considerably higher peak levels in the rats that received VEH before the tone test than in those that received AMPH. Three-way ANOVA (infusion before conditioning \times infusion before tone test \times 1 min block) of freezing during the first four 1 min blocks of tone presentation yielded a main effect of infusion before tone test [$F(1,33) = 4.75, P < 0.04$] and a significant main effect of 1 min block [$F(3,99) = 6.00, P < 0.001$] without an interaction of infusion before tone test \times 1 min block [$F(3,99) = 2.28, P > 0.08$]. Although it appeared as if the AMPH-AMPH group exhibited slightly higher levels of freezing than the VEH-AMPH rats, ANOVA did not yield a main effect of infusion before conditioning [$F(1,33) = 0.01, P > 0.94$] or any significant interaction involving infusion before conditioning [infusion before conditioning \times 1 min block, $F(3,99) = 0.50, P > 0.68$; infusion before conditioning \times infusion before tone test, $F(1,33) = 1.58, P > 0.21$; infusion before conditioning \times infusion before tone test \times 1 min block, $F(3,99) = 0.75, P > 0.52$]. Thus, only the infusion received immediately before the tone test significantly affected levels of freezing to the tone.

Activity Experiment: Effects of FLU or AMPH in the mPFC on General Activity

The effects of FLU or AMPH infusion into the mPFC on general activity during 2 h following infusion are depicted in Figure 6.

FLU decreased the general activity as compared to VEH infusions, while AMPH did not have any effect. ANOVA of the average activity during the 2 h of testing yielded a significant group effect [$F(2,20) = 14.23, P < 0.0001$] and *post hoc* comparisons indicated that activity was lower in the FLU group than in both the AMPH ($P < 0.0015$) and VEH ($P < 0.0001$) groups, which did not differ ($P > 0.2$). A more detailed analysis was done for the activity data collected during the first 15 1 min blocks following infusion, which corresponded to the period of

conditioning and tone-test sessions in the fear conditioning experiments. ANOVA yielded a significant effect of group [$F(2,20) = 8.37, P < 0.003$] and 1 min blocks [$F(14,580) = 22.71, P < 0.0001$], with the latter reflecting habituation. *Post hoc* comparisons indicated that average activity throughout the 15 min following infusion was lower in the FLU group than in both the AMPH ($P < 0.0015$) and VEH ($P < 0.003$) groups, which did not differ ($P > 0.58$). Although ANOVA did not yield an interaction of group \times time block [$F(28,580) = 1.25, P > 0.18$], it can be seen from Figure 6 that, consistent with the measurement of unconditioned activity during the 1.5 min preceding the first tone-shock pairing in experiment II (Fig. 3), activity in the FLU group was not considerably lower than in the VEH and AMPH rats during the first 1 min block. Indeed, a separate ANOVA on activity during the first 1 min block did not yield an effect of group [$F(2,20) = 1.40, P > 0.27$]. On day 3, groups did not differ in their activity any more (data not shown).

Discussion

The main finding of the present study is that both the dopamine receptor antagonist *cis*-flupenthixol (experiments I and II) and the dopamine receptor agonist *D*-amphetamine (experiments I and III) in the mPFC reduced the expression of conditioned fear, measured as freezing, to a tone previously paired with foot shocks. While *D*-amphetamine also reduced immediate conditioned freezing during conditioning, presence of neither drug during the tone-shock pairings affected conditioned fear during subsequent presentation of the tone alone. These data indicate that dopamine-receptor-mediated processes within the mPFC are important for the retrieval/expression, but not the formation, of conditioned fear.

Specificity of the Infusion Effects

Infusion sites were located in the central mPFC, mainly in the prelimbic and the dorsal infralimbic subdivisions. With the small infusion volume (0.5 μ l/side) and fine infusion cannulae (34 gauge) used in the present study, the estimated spread of the infused substances was <1 mm (Myers, 1966; Myers *et al.*, 1971; Routtenberg, 1972). It can therefore be assumed that differences between rats infused with vehicle compared to rats infused with FLU or AMPH reflected a temporary alteration of neuronal activity by blockade of dopamine receptors or increased release of dopamine and possibly other monoamines, respectively, within the central mPFC.

The similar immediate shock responses observed in all infusion groups in the present study, as well as previous findings that neither FLU nor AMPH in the mPFC, at the doses used in the present study, affected the acoustic startle reaction or its prepulse inhibition (Lacroix *et al.*, 2000a; Pezze *et al.*, 2001a; Bast *et al.*, 2002b), indicate that these infusions do not considerably affect sensory processing.

The activity experiment demonstrated that, at the doses used in the present study, FLU in the mPFC decreased activity (Beninger *et al.*, 1990; Pezze *et al.*, 2001a; Bast *et al.*, 2002b), while AMPH did not alter activity. Consistent with the similar unconditioned activity of VEH and FLU rats during the 1.5 min block preceding the first tone-shock pairing in experiment II, the decrease in activity induced by FLU in the control experiment was not evident before the second 1 min block. Infusion-induced hypoactivity cannot, however, account for reduced freezing. Rather, hypoactivity facilitates freezing (Bast *et al.*, 2001b) and may have somewhat concealed reduced conditioned fear in rats infused with FLU.

Finally, the reduced conditioned freezing observed during

the tone-test sessions in rats that received drug infusions immediately before the sessions cannot be explained by state dependency, i.e. that information learned in a particular brain state can in some cases only be retrieved with the same brain state prevailing (Overton, 1964). This was demonstrated by the results of experiments II and III, in which the drug-vehicle groups expressed normal freezing, while the drug-drug groups demonstrated a pronounced deficit.

Role of the mPFC and Local Dopamine Transmission in Fear Conditioning

Previous neurochemical studies indicated mPFC dopamine activation by a fear CS [(Goldstein *et al.*, 1994, 1996; Inoue *et al.*, 1994; Yoshioka *et al.*, 1996; Morrow *et al.*, 1999b,c; Feenstra, 2000; Feenstra *et al.*, 2001), but see Wilkinson *et al.* (Wilkinson *et al.*, 1998)] without clarifying the functional relation between dopamine in the mPFC and conditioned fear. The present finding that both the dopamine receptor antagonist FLU and the dopamine receptor agonist AMPH in the mPFC reduced freezing to a tone previously paired with foot shocks suggests that retrieval/expression of conditioned fear depends on normal mPFC dopamine transmission. The fact that the expression of conditioned fear was not completely blocked, but only reduced, by interference with prefrontal dopamine transmission is consistent with the widely held notion that the mPFC has a modulatory, rather than primary, role in the expression of fear (Morrow *et al.*, 1999a). Moreover, AMPH, but not FLU, in the mPFC reduced the expression of conditioned freezing during conditioning. This may be because dopamine receptor-mediated processes important for the retrieval/expression of immediate conditioned fear during conditioning differ from those relevant for the retrieval/expression of long-term conditioned fear assessed in later test sessions. Finally, manipulating mPFC dopamine transmission during the tone-shock pairings did not affect conditioned freezing expressed during a subsequent presentation of the aversive CS, suggesting that mPFC dopamine transmission does not play a role in the formation of a stable fear memory.

As described in the Introduction, the functional relation between neural activity in the mPFC and conditioned fear may depend on the subregion within the mPFC. In several previous studies, lesions centering on the prelimbic and dorsal infralimbic mPFC resulted in decreased conditioned fear, suggesting that processes in these regions may be essential for normal conditioned fear (Fryszak and Neafsey, 1991, 1994; Lacroix *et al.*, 2000b; McLaughlin *et al.*, 2001). Consistent with this suggestion, the results of the present study indicate that dopamine transmission in the prelimbic and dorsal infralimbic mPFC, which was the center of the infusion sites, is critical for the normal expression of conditioned fear. Dopamine transmission in other mPFC subregions may play a different role, given that lesions affecting the anterior cingulate (Morgan and LeDoux, 1995; Vouimba *et al.*, 2000) and the most caudal and ventral infralimbic mPFC (Morgan *et al.*, 1993; Quirk *et al.*, 2000) were reported to have facilitating effects on conditioned fear. This issue may be addressed in future studies.

In some studies, lesions destroying considerable parts of the prelimbic and dorsal infralimbic mPFC did not alter conditioned fear (Holson, 1986; Rosen *et al.*, 1992; Gewirtz *et al.*, 1997; Quirk *et al.*, 2000). Moreover, 6-hydroxydopamine lesions to the mPFC, depleting dopamine (by 86%), norepinephrine (by 51%), which has been implicated in extinction of conditioned responses (Dalley *et al.*, 2001) and serotonin (by 14%) in the rostral cingulate, prelimbic and rostral infralimbic mPFC, did

not affect moderate conditioned freezing to a tone (peak levels ~40–50%), while extinction of strong conditioned freezing (peak levels ~80%) was strongly reduced (Morrow *et al.*, 1999b). The latter suggests that the significance of catecholamine transmission in the mPFC for the extinction of conditioned fear may depend on the intensity of fear (Morrow *et al.*, 1999b; Quirk *et al.*, 2000). The fact that permanent damage to the prelimbic and dorsal infralimbic mPFC, in contrast to our temporary manipulation of local dopamine transmission, sometimes failed to reduce conditioned fear may reflect compensatory processes induced by permanent lesions (Bast *et al.*, 2001a).

Interactions of mPFC Dopamine and Fear Circuits

While efferents to the nucleus accumbens, amygdala, periaqueductal gray, or lateral hypothalamus (Sesack *et al.*, 1989) link the mPFC to structures that have been implicated in the expression of conditioned fear (Fendt and Fanselow, 1999; LeDoux, 2000; Murphy *et al.*, 2000; Pezze *et al.*, 2001b, 2002a), the mPFC may receive information about CS-US associations through afferents from the amygdala (McDonald, 1991) and ventral hippocampus (Jay and Witter, 1991). Both the ventral hippocampus (Bast *et al.*, 2001a,b, 2003; Zhang *et al.*, 2001) and amygdala (Cahill *et al.*, 1999; Fanselow and LeDoux, 1999; LeDoux, 2000) have been implicated in the associative processes underlying formation of conditioned fear and the amygdala may even be the site where the CS-US association is permanently stored [(Fanselow and LeDoux, 1999; Fendt and Fanselow, 1999; LeDoux, 2000), but see Cahill *et al.* (Cahill *et al.*, 1999)]. Interestingly, electrical stimulation of the amygdala (Jackson and Moghaddam, 2001) or the ventral hippocampus (Gurden *et al.*, 2000) has been demonstrated to result in measurable effects on neural processes in the mPFC, including an increase of extracellular dopamine. Thus, mPFC dopamine transmission may contain information about CS-US associations transmitted from the amygdala or the ventral hippocampus and the reduced expression of conditioned fear observed after interference with mPFC dopamine transmission may indicate that the transfer of this information was disrupted.

Relation between mPFC Dopamine and Conditioned Fear

In the present study, both the dopamine receptor antagonist FLU and the agonist AMPH reduced expression of conditioned fear when infused into the mPFC. This may be because normal retrieval/expression of conditioned fear requires time and synapse specific dopamine-receptor stimulation in the mPFC, but this is unlikely in view of the evidence for dopaminergic volume transmission in the mPFC (Bast *et al.*, 2002a). Alternatively, it may reflect an inverted U-shaped function relating the retrieval/expression of conditioned fear to dopamine-receptor stimulation in the mPFC, i.e. that normal retrieval/expression of conditioned fear requires an optimal level of dopamine activity in the mPFC. Similarly, working memory has been suggested to require an optimal level of prefrontal dopamine-receptor stimulation, because PFC infusions of both dopamine-receptor antagonists and agonists impaired working memory (Sawaguchi and Goldman-Rakic, 1991; Williams and Goldman-Rakic, 1995; Zahrt *et al.*, 1997; Arnsten, 1998; Seamans *et al.*, 1998; Goldman-Rakic *et al.*, 2000; Robbins, 2000). These effects have been found with D1-receptor-specific agents and some studies have directly demonstrated that D2-receptor antagonists are without an effect [(Sawaguchi and Goldman-Rakic, 1991; Williams and Goldman-Rakic, 1995; Seamans *et al.*, 1998), but see Arnsten *et al.* and Druzin *et al.* (Arnsten *et al.*, 1995; Druzin *et al.*, 2000)]. Attentional performance also appears to depend on an optimal

level of mPFC D1-receptor stimulation (Granon *et al.*, 2000; Robbins, 2000). Electrophysiological findings indicate that behavioral processes involving dopamine-mediated mechanisms in the mPFC may require an optimal level of local dopamine activity, because an optimal level of local D1-receptor stimulation is essential for the 'sharpening' of prefrontal signal transmission (Yang and Seamans, 1996). A final consideration is that AMPH may not only directly stimulate dopamine, but also norepinephrine and serotonin transmission (Azzaro and Rutledge, 1973) and that this might contribute to the reduction of conditioned fear by AMPH in the mPFC. Thus, future studies should examine the effects of mPFC infusion of specific D1- or D2-receptor antagonists or agonists on conditioned fear in order to clarify whether local D1 and D2 receptors are differently involved in conditioned fear and to substantiate the suggestion that excess prefrontal dopamine receptor stimulation may reduce conditioned fear.

Conclusion

The present study suggested that inhibition as well as stimulation of dopamine receptors in the (prelimbic and dorsal infralimbic) mPFC reduced conditioned fear to a tone previously paired with a foot shock. Our findings corroborate the view that the mPFC and local dopamine transmission are essential for normal emotional responses. Furthermore, they suggest the retrieval/expression of conditioned fear, as with working memory and attentional performance, may be related to mPFC dopamine transmission by an inverted U-shaped function, indicating an intriguing relation between the substrates of emotion, working memory and attention.

Notes

This research was supported by grants from the Swiss National Science Foundation Zurich (grants Nos 3160-051657 and 3100-042009.94) and the Swiss Federal Institute of Technology Zurich. The authors thank L. Lacroix for helpful advice to start this study; J. Kupper, P. Guela, O. Aspiron and J. Torlucci for animal care; P. Schmid for set-up and maintenance of the computerized systems for behavioral analysis; L. Weber for histological preparations; C. Schlatter for preparation of the guide cannulae; and J. Fotheringham for her editorial help. W. White and P. Schmid set up the system for measurement of general activity. Part of the results presented in this paper have previously been published in preliminary form [Pezze *et al.*, 2001a, 2002b].

Address correspondence to Joram Feldon, Laboratory of Behavioral Neurobiology, Swiss Federal Institute of Technology Zurich (ETH), Schorenstrasse 16, 8603 Schwerzenbach, Switzerland. Email: feldon@behav.biol.ethz.ch.

References

- Anagnostaras SG, Maren S, Sage JR, Goodrich S, Fanselow MS (1999) Scopolamine and Pavlovian fear conditioning in rats: dose-effect analysis. *Neuropsychopharmacology* 21:731–744.
- Arnsten A (1998) Catecholamine modulation of prefrontal cortical cognitive function. *Trends Cogn Sci* 2:336–447.
- Arnsten AFT, Cai JX, Steere JC, Goldman-Rakic PS (1995) Dopamine D2 receptor mechanisms contribute to age related cognitive decline: the effects of quinpirole on memory and motor performance in monkeys. *J Neurosci* 15:3429–3439.
- Azzaro AJ, Rutledge CO (1973) Selectivity of release of norepinephrine, dopamine, and 5-hydroxytryptamine by amphetamine in various regions of rat brain. *Biochem Pharmacol* 22:2801–2813.
- Baeg EH, Kim YB, Jang J, Kim HT, Mook-Jung I, Jung MW (2001) Fast spiking and regular spiking neural correlates of fear conditioning in the medial prefrontal cortex of the rat. *Cereb Cortex* 11:441–451.
- Baldessarini RJ, Kula NS, Zong R, Neumeyer JL (1994) Receptor affinities of aporphine enantiomers in rat brain tissue. *Eur J Pharmacol* 254:199–203.

- Bast T, Zhang W-N, Feldon J (2001a) Hippocampus and classical fear conditioning. *Hippocampus* 11:828–831.
- Bast T, Zhang W-N, Feldon J (2001b) The ventral hippocampus and fear conditioning in rats. Different anterograde amnesias of fear after tetrodotoxin inactivation and infusion of the GABA(A) agonist muscimol. *Exp Brain Res* 139:39–52.
- Bast T, Diekamp B, Thiel C, Schwarting RKW, Güntürkün O (2002a) Functional aspects of dopamine metabolism in the putative prefrontal cortex analogue and striatum of pigeons (*Columba livia*). *J Comp Neurol* 445:58–67.
- Bast T, Pezze MA, Feldon J (2002b) Dopamine receptor blockade in the rat medial prefrontal cortex reduces spontaneous and amphetamine-induced activity and does not affect prepulse inhibition. *Behav Pharmacol* 13:669–673.
- Bast T, Zhang W-N, Feldon J (2003) Dorsal hippocampus and classical fear conditioning to tone and context in rats: the effects of local NMDA-receptor blockade and stimulation. *Hippocampus* 13: in press.
- Bechara A, Damasio H, Damasio AR (2000) Emotion, decision making and the orbitofrontal cortex. *Cereb Cortex* 10:295–307.
- Beninger RJ, Musgrave MA, Dickson PR (1990) Unilateral injections of a D2 but not D1 agonist into the frontal cortex of rats produce a contralateral directional bias. *Pharmacol Biochem Behav* 37:387–392.
- Cahill L, Weinberger NM, Roozendaal B, McGaugh JL (1999) Is the amygdala a locus of 'conditioned fear'? Some questions and caveats. *Neuron* 23:227–228.
- Cochran WG, Cox GM (1950). *Experimental designs*. New York: John Wiley.
- Corcoran KA, Maren S (2001) Hippocampal inactivation disrupts contextual retrieval of fear memory after extinction. *J Neurosci* 21:1720–1726.
- Dalley JW, McGaughy J, O'Connell MTO, Cardinal RN, Levita L, Robbins TW (2001) Distinct changes in cortical acetylcholine and noradrenaline efflux during contingent and noncontingent performance of a visual attentional task. *J Neurosci* 21:4908–4914.
- Davidson RJ, Irwin W (1999) The functional neuroanatomy of emotion and affective style. *Trends Cogn Sci* 3:11–21.
- Druzin MY, Kurzina NP, Malinina EP, Kozlov AP (2000) The effects of local application of D2 selective dopaminergic drugs into the medial prefrontal cortex of rats in a delayed spatial choice task. *Behav Brain Res* 109:99–111.
- Fanselow MS (1984) What is conditioned fear? *Trends Neurosci* 7:460–462.
- Fanselow MS, LeDoux JE (1999) Why we think plasticity underlying Pavlovian fear conditioning occurs in the basolateral amygdala. *Neuron* 23:229–232.
- Feenstra MG (2000) Dopamine and noradrenaline release in the prefrontal cortex in relation to unconditioned and conditioned stress and reward. *Prog Brain Res* 126:133–163.
- Feenstra MG, Teske G, Botterblom MH, De Bruin JP (1999) Dopamine and noradrenaline release in the prefrontal cortex of rats during classical aversive and appetitive conditioning to a contextual stimulus: interference by novelty effects. *Neurosci Lett* 272:179–182.
- Feenstra MG, Vogel M, Botterblom MH, Joosten RN, de Bruin JP (2001) Dopamine and noradrenaline efflux in the rat prefrontal cortex after classical aversive conditioning to an auditory cue. *Eur J Neurosci* 13:1051–1054.
- Fendt M, Fanselow MS (1999) The neuroanatomical and neurochemical basis of conditioned fear. *Neurosci Biobehav Rev* 23:743–760.
- Fryszak RJ, Neafsey EJ (1991) The effect of medial frontal cortex lesions on respiration, 'freezing,' and ultrasonic vocalizations during conditioned emotional responses in rats. *Cereb Cortex* 1:418–425.
- Fryszak RJ, Neafsey EJ (1994) The effect of medial frontal cortex lesions on cardiovascular conditioned emotional responses in the rat. *Brain Res* 643:181–193.
- Garcia R, Vouimba RM, Baudry M, Thompson RF (1999) The amygdala modulates prefrontal cortex activity relative to conditioned fear. *Nature* 402:294–296.
- Gewirtz JC, Falls WA, Davis M (1997) Normal conditioned inhibition and extinction of freezing and fear-potentiated startle following electrolytic lesions of medial prefrontal cortex in rats. *Behav Neurosci* 111:712–726.
- Goldman-Rakic PS, Muly EC III, Williams GV (2000) D₁ receptors in prefrontal cells and circuits. *Brain Res Rev* 31:295–301.
- Goldstein LE, Rasmusson AM, Bunney BS, Roth RH (1994) The NMDA glycine site antagonist (+)-HA-966 selectively regulates conditioned stress-induced metabolic activation of the mesoprefrontal cortical dopamine but not serotonin systems: a behavioral, neuroendocrine, and neurochemical study in the rat. *J Neurosci* 14:4937–4950.
- Goldstein LE, Rasmusson AM, Bunney BS, Roth RH (1996) Role of the amygdala in the coordination of behavioral, neuroendocrine, and prefrontal cortical monoamine responses to psychological stress in the rat. *J Neurosci* 16:4787–4798.
- Granon S, Passetti F, Thomas KL, Dalley JW, Everitt BJ, Robbins TW (2000) Enhanced and impaired attentional performance after infusion of D1 dopaminergic receptor agents into rat prefrontal cortex. *J Neurosci* 20:1208–1215.
- Gurden H, Takita M, Jay TM (2000) Essential role of D1 but not D2 receptors in the NMDA receptor-dependent long term potentiation at hippocampal-prefrontal cortex synapses *in vivo*. *J Neurosci* 20:RC106.
- Holston RR (1986) Mesial prefrontal cortical lesions and timidity in rats. I. Reactivity to aversive stimuli. *Physiol Behav* 37:221–230.
- Inoue T, Tsuchiya K, Koyama T (1994) Regional changes in dopamine and serotonin activation with various intensity of physical and psychological stress in the rat brain. *Pharmacol Biochem Behav* 49:911–920.
- Jackson ME, Moghaddam B (2001) Amygdala regulation of nucleus accumbens dopamine output is governed by the prefrontal cortex. *J Neurosci* 21:676–681.
- Jay TM, Witter MP (1991) Distribution of hippocampal CA1 and subicular efferents in the prefrontal cortex of the rat studied by means of anterograde transport of Phaseolus vulgaris-leucoagglutinin. *J Comp Neurol* 313:574–586.
- Lacroix L, Broersen LM, Feldon J, Weiner I (2000a) Effects of local infusions of dopaminergic drugs into the medial prefrontal cortex of rats on latent inhibition, prepulse inhibition and amphetamine induced activity. *Behav Brain Res* 107:111–121.
- Lacroix L, Spinelli S, Heidbreder CA, Feldon J (2000b) Differential role of the medial and lateral prefrontal cortices in fear and anxiety. *Behav Neurosci* 114:1119–1130.
- LeDoux JE (2000) Emotion circuits in the brain. *Annu Rev Neurosci* 23:155–184.
- McDonald AJ (1991) Organization of amygdaloid projections to the prefrontal cortex and associated striatum in the rat. *Neuroscience* 44:1–14.
- McLaughlin J, Flaten MA, Chachich M, Powell DA (2001) Medial prefrontal lesions attenuate conditioned reflex facilitation but do not affect prepulse modification of the eyeblink reflex in rabbits. *Exp Brain Res* 140:318–325.
- Maren S (2001) Neurobiology of Pavlovian fear conditioning. *Annu Rev Neurosci* 24:897–931.
- Morgan MA, LeDoux JE (1995) Differential contribution of dorsal and ventral medial prefrontal cortex to the acquisition and extinction of conditioned fear in rats. *Behav Neurosci* 109:681–688.
- Morgan MA, Romanski LM, LeDoux JE (1993) Extinction of emotional learning: contribution of medial prefrontal cortex. *Neurosci Lett* 163:109–113.
- Morrow BA, Elsworth JD, Inglis FM, Roth RH (1999a) An antisense oligonucleotide reverses the footshock-induced expression of fos in the rat medial prefrontal cortex and the subsequent expression of conditioned fear-induced immobility. *J Neurosci* 19:5666–5673.
- Morrow BA, Elsworth JD, Rasmusson AM, Roth RH (1999b) The role of mesoprefrontal dopamine neurons in the acquisition and expression of conditioned fear in the rat. *Neuroscience* 92:553–564.
- Morrow BA, Elsworth JD, Zito C, Roth RH (1999c) Biochemical and behavioral anxiolytic-like effects of R(+)-HA-966 at the level of the ventral tegmental area in rats. *Psychopharmacology (Berl)* 143:227–234.
- Murphy CA, Pezze M, Feldon J, Heidbreder C (2000) Differential involvement of dopamine in the shell and core of the nucleus accumbens in the expression of latent inhibition to an aversively conditioned stimulus. *Neuroscience* 97:469–477.
- Myers RD (1966) Injection of solutions into cerebral tissue: relation between volume and diffusion. *Physiol Behav* 1:171–174.
- Myers RD, Tytell M, Kawa A, Rudy T (1971) Micro-injection of 3H-acetylcholine, 14C-serotonin and 3H-norepinephrine into the hypothalamus of the rat: diffusions into tissue and ventricles. *Physiol Behav* 7:743–751.
- Nauta WJH (1971) The problem of the frontal lobe: a reinterpretation. *J Psychiatr Res* 8:167–187.

- Öngür D, Price JL (2000) The organization of networks within the orbital and medial prefrontal cortex of rats, monkeys and humans. *Cereb Cortex* 10:206–219.
- Overton DA (1964) State-dependent or 'dissociated' learning produced with pentobarbital. *J Comp Physiol Psychol* 57:3–12.
- Paxinos G, Watson C (1998) *The rat brain in stereotaxic coordinates*. San Diego, CA: Academic Press.
- Pezze M, Bast T, Feldon J (2001a) Effects of dopamine stimulation or blockade in the medial prefrontal cortex on activity, sensory motor gating, and conditioned fear. *Behav Pharmacol* 12:S77.
- Pezze MA, Heidbreder CA, Feldon J, Murphy CA (2001b) Selective responding of nucleus accumbens core and shell dopamine to aversively conditioned contextual and discrete stimuli. *Neuroscience* 108:91–102.
- Pezze M, Feldon J, Murphy C (2002a) Increased conditioned fear response and altered balance of dopamine in the shell and core of the nucleus accumbens during amphetamine withdrawal. *Neuropharmacology* 42:633–643.
- Pezze M, Bast T, Feldon J (2002) Significance of dopamine transmission in the rat medial prefrontal cortex for conditioned fear. *FENS Abstr* 1:A042.21.
- Pryce CR, Lehmann J, Feldon J (1999) Effect of sex on fear conditioning is similar for context and discrete CS in Wistar, Lewis and Fischer rat strains. *Pharmacol Biochem Behav* 64:753–759.
- Quirk GJ, Russo GK, Barron JL, Lebron K (2000) The role of ventromedial prefrontal cortex in the recovery of extinguished fear. *J Neurosci* 20:6225–6231.
- Richmond MA, Murphy CA, Pouzet B, Schmid P, Rawlins JN, Feldon J (1998) A computer controlled analysis of freezing behaviour. *J Neurosci Methods* 86:91–99.
- Robbins TW (2000) Chemical neuromodulation of frontal-executive functions in humans and other animals. *Exp Brain Res* 133:130–138.
- Rosen JB, Hitchcock JM, Miserendino MJ, Falls WA, Campeau S, Davis M (1992) Lesions of the perirhinal cortex but not of the frontal, medial prefrontal, visual, or insular cortex block fear-potentiated startle using a visual conditioned stimulus. *J Neurosci* 12:4624–4633.
- Routtenberg A (1972) Intracranial chemical injection and behavior: a critical review. *Behav Biol* 7:601–641.
- Sawaguchi T, Goldman-Rakic PS (1991) D1 dopamine receptors in prefrontal cortex: involvement in working memory. *Science* 251:947–950.
- Seamans JK, Floresco SB, Phillips AG (1998) D₁ receptor modulation of hippocampal-prefrontal cortical circuits integrating spatial memory with executive functions in the rat. *J Neurosci* 18:1613–1621.
- Sesack SR, Deutch AY, Roth RH, Bunney BS (1989) Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with Phaseolus vulgaris leucoagglutinin. *J Comp Neurol* 290:213–242.
- Simpson JR, Öngür D, Akbudak E, Conturo TE, Ollinger JM, Snyder AZ, Gusnard DA, Raichle ME (2000) The emotional modulation of cognitive processing: an fMRI study. *J Cogn Neurosci* 12:157–170.
- Vouimba RM, Garcia R, Baudry M, Thompson RF (2000) Potentiation of conditioned freezing following dorsomedial prefrontal cortex lesions does not interfere with fear reduction in mice. *Behav Neurosci* 114:720–724.
- Wilkinson LS, Humby T, Killcross AS, Torres EM, Everitt BJ, Robbins TW (1998) Dissociations in dopamine release in medial prefrontal cortex and ventral striatum during the acquisition and extinction of classical aversive conditioning in the rat. *Eur J Neurosci* 10:1019–1026.
- Williams GV, Goldman-Rakic PS (1995) Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature* 376:572–575.
- Yang CR, Seamans JK (1996) Dopamine D1 receptor actions in layers V–VI rat prefrontal cortex neurons *in vitro*: modulation of dendritic-somatic signal integration. *J Neurosci* 16:1922–1935.
- Yoshioka M, Matsumoto M, Togashi H, Saito H (1996) Effect of conditioned fear stress on dopamine release in the rat prefrontal cortex. *Neurosci Lett* 209:201–203.
- Zahrt J, Taylor JR, Mathew RG, Arnsten AF (1997) Supranormal stimulation of D1 dopamine receptors in the rodent prefrontal cortex impairs spatial working memory performance. *J Neurosci* 17:8528–8535.
- Zhang WN, Bast T, Feldon J (2001) The ventral hippocampus and fear conditioning in rats: different anterograde amnesias of fear after infusion of *N*-methyl-D-aspartate or its noncompetitive antagonist MK-801 into the ventral hippocampus. *Behav Brain Res* 126:159–174.