

In vitro cultivation of pleomorphic *Trypanosoma brucei* stocks: a possible source of variable antigens for immunization studies

L. JENNI AND R. BRUN

Swiss Tropical Institute, CH-4051 Basle, Switzerland

The number and pattern of variable antigen types (VAT) in metacyclic forms of *Trypanosoma brucei* and *T. rhodesiense* has been investigated by several workers (CUNNINGHAM, 1966; BARRY & HAJDUK, 1979; JENNI, 1979). However, the degree of antigenic heterogeneity of metacyclic populations is still far from being clear. Further experiments have been carried out recently in order to analyse the pattern of metacyclic VAT which appeared in *Glossina m. morsitans* during cyclical transmission of one stock of *T. brucei*.

Material and Methods

Trypanosoma brucei STIB 247 was isolated in 1971 in the Serengeti National Park (Tanzania) from a hartebeest (*Alcelaphus buselaphus*) and cryopreserved after one rat passage. For the last 13 months this primary isolate has been cyclically passaged through *Glossina m. morsitans* and Swiss-ICR mice. Metacyclic populations were harvested and used as starting inoculum for continuous *in vitro* cultivation of vertebrate-infective forms according to BRUN *et al.* (1979). The feeder layer consisted of rabbit embryo fibroblast-like cells.

The neutralization of infectivity test was carried out as described by SCHLÄPPI & JENNI (1977) by incubating 10^4 trypanosomes per test. Rabbit antiserum against metacyclic VAT was prepared according to JENNI (1979).

In vitro cultured trypanosomes were harvested, centrifuged at 1000g and the pellet resuspended in PSG buffer at pH 8.0. This suspension was then passed through a DEAE-cellulose column

(LANGHAM & GODFREY, 1970) in order to separate the trypanosomes from detached feeder layer cells. The trypanosomes were irradiated with 50 krad at 2 to 4 °C by a ^{60}Co source in PSG and fresh rabbit serum. After irradiation, the trypanosomes were counted and injected intraperitoneally into 10 female Swiss-ICR mice (25g) in equal amounts three times with no additional adjuvant (Table 1). Control animals were injected with the same amount of PSG and fresh rabbit serum as the test animals. One week after the last immunizing injection the mice were weekly challenged with one infective tsetse fly bite (Table I).

Results and Discussion

In vitro cultures were initiated with 10^4 metacyclic forms per ml culture medium (1 ml per well). After six hours the metacyclic forms transformed to slender-like bloodstream forms and after another six hours started to divide. After 50 hours a total trypanosome density (including parasites between feeder layer cells and in the supernatant medium) of 2.7×10^5 trypanosomes per well was normally reached.

The infectivity of metacyclic forms was completely neutralized by the rabbit antiserum. Daily tests with *in vitro* cultured parasites showed that the neutralizing activity of the antiserum was lost when the parasites had been in culture for more than 50 hours. After this growth period initial antigenic variation had occurred and new VAT were present within the parasite population which were not recognized by the antiserum. Thus for the *in vitro* culture trypanosomes were

Table I—Immunization of mice against the infective tsetse fly bite (*T. brucei* STIB 257)

	IMMUNIZATION			TSETSE FLY CHALLENGE				
	No. of irradiated trypanosomes injected per mouse at weekly intervals			No. of mice showing parasitaemia				
10 Test mice	6.5×10^4 Tryps in 0.1 ml PSG + 0.5% rs	1.2×10^4 Tryps in 0.1 ml PSG + 2.5% rs	5.0×10^4 Tryps in 0.1 ml PSG + 1.0% rs	↓	↓	↓	↓	↓
				0/10	0/10	0/10	0/10	0/10
				↓				
10 Control mice	0.1 ml PSG + 0.5% rs	0.1 ml PSG + 2.5% rs	0.1 ml PSG + 1.0% rs	10/10	10/10	10/10	7/10 3 mice dead	— all mice dead
Days	0	7	14	21	28	35	42	49

↓ Challenge with one infective tsetse fly bite

harvested 50 hours after initiation with metacyclic forms and used for immunization.

The mice were challenged each time with single infective flies which were taken at random from the infective fly pool in the laboratory. These flies derived from different transmission series with STIB 247 and were different from those flies from which the metacyclic forms were harvested for *in vitro* cultivation.

The tail blood of the challenged mice was examined by the haematocrit method. While the 10 test mice showed no parasitaemia after five infective bites all control mice showed parasites in the blood within 72 hours after the first challenge and died within four weeks (Table I).

DOYLE *et al.*, (1979). reported antigenic variation in *T. brucei* *in vitro* for the first time. Our results showed that this also occurred *in vitro* where trypanosomes directly derived from metacyclic forms were grown in the culture system described by BRUN *et al.* (1979). In addition, this system also allows the growth of clones from single metacyclic forms. These clones showed new VAT after eight days. However, new VAT appeared earlier in mice which have been cyclically infected. In these mice new VAT could already be detected 36 hours after the infective fly bite.

The above results show that initial antigenic variation *in vitro* is independent of the first cell divisions after the transformation from metacyclic to the slender-like bloodstream form. The results also show that metacyclic VAT can be quantitatively increased *in vitro* during a limited initial growth period. It is interesting to note that STIB 247 reverted during each fly passage to the same metacyclic VAT, for the immunized mice were completely protected against any metacyclic population of STIB 247. On the other hand, metacyclic VAT of other trypanosome stocks did infect mice immunized against metacyclic VAT of STIB 247. This is indicated by current experiments where mice immunized against STIB 247 are being

challenged with flies infected with different *T. brucei* and *T. congolense* stocks.

Acknowledgements

This study was supported by the Swiss National Science Foundation, grant No. 3.346-0.78. We wish to thank K. F. Schell, M. Schönenberger and M. Kauffmann for skilful technical assistance. We are grateful to Drs. A. M. Jordan and L. W. Roberts for the supply of *Glossina* pupae.

References

- Barry, J. D. & Hajduk, S. L. (1979). Antigenic heterogeneity of bloodstream and metacyclic forms of *T. brucei*. In: *Pathogenicity of Trypanosomes*. Losos, G. & Chouinard, A. (Editors). IDRC-132e, Ottawa, Canada, pp. 51-56.
- Brun, R., Jenni, L., Tanner, M., Schönenberger, M. & Schell, K.-F. (1979). Cultivation of vertebrate infective forms derived from metacyclic forms of pleomorphic *Trypanosoma brucei* stocks. *Acta Tropica*, **36**, 387-390.
- Cunningham, M. P. (1966). The preservation of viable metacyclic forms of *Trypanosoma rhodesiense*. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **60**, 126.
- Doyle, J. J., Hirumi, H. & de Gee, A. L. W. (1979). Mechanisms of antigenic variation in salivarian trypanosomes. In: *Pathogenicity of Trypanosomes*. Losos, G. & Chouinard, A. (Editors). IDRC-132e, Ottawa, Canada, pp. 44-45.
- Jenni, L. (1979). Cyclical transmission and antigenic variation. In: *Pathogenicity of Trypanosomes*. Losos, G. & Chouinard, A. (Editors). IDRC-132e, Ottawa, Canada, pp. 49-50.
- Lanham, S. M. & Godfrey, D. G. (1970). Isolation of salivarian trypanosomes from man and other mammals using DEAE-cellulose. *Experimental Parasitology*, **28**, 521-534.
- Schläppi, B. & Jenni, L. (1977). Studies on antigenic variation of cyclically transmitted *Trypanosoma congolense*. *Acta Tropica*, **34**, 43-51.