

# *Corynebacterium lipophiloflavum* sp. nov. isolated from a patient with bacterial vaginosis

Guido Funke <sup>a,\*</sup>, Roger A. Hutson <sup>b</sup>, Markus Hilleringmann <sup>c</sup>,  
Wolfgang R. Heizmann <sup>d</sup>, Matthew D. Collins <sup>b</sup>

<sup>a</sup> Department of Medical Microbiology, University of Zürich, 8028 Zürich, Switzerland

<sup>b</sup> Department of Microbiology, BBSRC Institute of Food Research, Reading RG6 6BZ, UK

<sup>c</sup> Max-Planck-Institute of Developmental Biology, 72076 Tübingen, Germany

<sup>d</sup> Medical Microbiology Laboratory Nordhorn, 48529 Nordhorn, Germany

Received 20 February 1997; revised 10 March 1997; accepted 11 March 1997

## Abstract

A unique coryneform bacterium was isolated from a patient with bacterial vaginosis. Chemotaxonomical investigations demonstrated that the unknown bacterium belonged to the genus *Corynebacterium*. The yellow-pigmented, slightly lipophilic, oxidative, urea-hydrolyzing bacterium could be phenotypically readily differentiated from the other members of the genus *Corynebacterium*. Comparative 16S rRNA gene analysis revealed that the bacterium represented a new subline within the genus *Corynebacterium* for which the name *Corynebacterium lipophiloflavum* sp. nov. is proposed. The type strain is CCUG 37336 (DSM 44291).

**Keywords:** *Corynebacterium lipophiloflavum* sp. nov.; *Corynebacterium* taxonomy; Bacterial vaginosis

## 1. Introduction

Bacterial vaginosis (BV) is a polymicrobial process involving *Gardnerella vaginalis* and anaerobic bacteria such as *Prevotella* spp., *Mobiluncus* spp., and *Peptostreptococcus* spp. as possible etiologic agents [1]. During the search for further bacterial pathogens involved in BV we isolated a unique coryneform bacterium which was further characterized. Investigations applying both phenotypic and molecular genetic methods revealed that the isolate represented a new *Corynebacterium* species.

Within the genera of coryneform bacteria the genus *Corynebacterium* comprises at present the largest number of single species, with eleven species having been defined within the last ten years [2,3]. This report adds further evidence to the enormous diversity within the genus *Corynebacterium*.

## 2. Materials and methods

### 2.1. Microorganism and cultivation

Strain DMMZ 1944, now named *C. lipophiloflavum*, was initially isolated from a vaginal swab which, apart from the other media used, was plated

\* Corresponding author. Tel.: +41 (1) 257-2700;  
fax: +41 (1) 252-8107; e-mail: [funke@immv.unizh.ch](mailto:funke@immv.unizh.ch)

Table 1  
Characteristics differentiating *C. lipophiloflavum* from other lipophilic *Corynebacterium* spp.<sup>a</sup>

Species	Type of metabolism <sup>b</sup>	Nitrate reduction	Urea hydrolysis	Pyrazinamidase	Alkaline phosphatase	Acid production <sup>c</sup>				TBSA <sup>d</sup>	Pigment
						Glu	Mal	Suc	Man		
<i>C. lipophiloflavum</i>	O	–	+	+	+	–	–	–	–	–	Yellow
<i>C. accolens</i>	F	+	–	V <sup>e</sup>	–	+	–	V	V	–	–
<i>C. afermentans</i> subsp. <i>lipophilum</i>	O	–	–	+	+	–	–	–	–	–	–
<i>C. bovis</i>	O	–	–	V	+	+	–	–	–	+	–
<i>C. diphtheriae</i> biotype <i>intermedius</i>	F	+	–	–	–	+	+	–	–	–	–
<i>C. jeikeium</i>	O	–	–	+	+	+	V	–	–	–	–
<i>C. macginleyi</i>	F	+	–	–	+	+	–	+	V	–	–
<i>C. urealyticum</i>	O	–	+	+	V	–	–	–	–	+	–
CDC group F-1	F	V	+	+	–	+	+	+	–	–	–
CDC group G	F	V	–	+	+	+	V	V	–	V	–

<sup>a</sup>Data from reference [2].

<sup>b</sup>O, oxidative; F, fermentative.

<sup>c</sup>Glu, glucose; Mal, maltose; Suc, sucrose; Man, mannitol.

<sup>d</sup>TBSA, tuberculostearic acid.

<sup>e</sup>V, variable.

on sheep blood agar (SBA) and incubated for 24 h in a 5% CO<sub>2</sub>-enriched atmosphere. The vaginal swab had been taken from a 32-year-old female with the clinical diagnosis of BV. *G. vaginalis* and *Peptostreptococcus* spp. were cultured from the same specimen in addition to strain DMMZ 1944.

## 2.2. Phenotypical analysis

The biochemical profile (enzymatic reactions, acid production from carbohydrates) of strain DMMZ 1944 was determined as described previously [4]. Chemotaxonomic investigations included determination of the cellular fatty acids (CFA), the diamino acid of the peptidoglycan, the presence of mycolic acids, and the G+C content [3]. The antimicrobial susceptibility pattern was determined by using the MCN system (Merlin Diagnostics, Bornheim, Germany) as outlined before [5].

## 2.3. Phylogenetic analysis

Determination of the 16S rRNA gene sequence of strain DMMZ 1944 (EMBL accession number Y09045) was by PCR-mediated amplification of the gene followed by cycle sequencing of the PCR product, and reading the reaction products on a model

373A automatic sequencer (Applied Biosystems, Foster City, USA) as described previously [6]. Phylogenetic analysis included alignment, calculation of percentage sequence similarity, construction of a phylogenetic tree, and an assessment of tree topology by bootstrap analysis as has also been outlined previously [6].

## 3. Results and discussion

Strain DMMZ 1944 formed tiny, non-hemolytic colonies of less than 0.2 mm in diameter after 24 h incubation on SBA at 37°C. Colonies were larger (up to 1 mm in diameter) when strain DMMZ 1944 was grown on SBA supplemented with 1% Tween 80, i.e., the strain was lipophilic [2]. The colonies showed an intense yellow pigment. The Gram stain showed typical club-shaped gram-positive rods of 1–3 µm in length which were arranged as single cells, as pairs, or in clusters. Strain DMMZ 1944 was not partially acid-fast.

The basic phenotypical screening reactions [7] of strain DMMZ 1944 were as follows: catalase positive (which separated the strain from the catalase-negative *G. vaginalis*); non-motile; oxidative metabolism; nitrate reduction negative; urea hydrolysis



Table 2

Levels of 16S rRNA sequence similarity between *C. lipophiloflavum* sp. nov. and other *Corynebacterium* species as well as *Turicella otitidis*

Species	% 16S rRNA sequence similarity with <i>C. lipophiloflavum</i> sp. nov.	Species	% 16S rRNA sequence similarity with <i>C. lipophiloflavum</i> sp. nov.
<i>C. accolens</i> (X80500) <sup>a</sup>	94.6	<i>C. matruchotii</i> (X84443)	92.2
<i>C. 'acetoacidophilum'</i> (X84240)	93.0	<i>C. minutissimum</i> (X84678)	94.5
<i>C. afermentans</i> (X81874)	96.6	<i>C. mycetoides</i> (X84241)	97.2
<i>C. ammoniagenes</i> (X84440)	94.4	<i>C. pilosum</i> (X84246)	95.8
<i>C. amycolatium</i> (X84244)	94.2	<i>C. propinquum</i> (X84438)	93.7
<i>C. auris</i> (X82493)	97.1	<i>C. pseudodiphtheriticum</i> (X84258)	93.6
<i>C. bovis</i> (X84444)	93.7	<i>C. 'pseudogenitalium'</i> (X81874)	96.2
<i>C. callunae</i> (X84251)	93.4	<i>C. pseudotuberculosis</i> (X84255)	94.1
<i>C. coyleae</i> (X96497)	96.7	<i>C. renale</i> (X84249)	94.6
<i>C. cystitidis</i> (X84252)	94.7	<i>C. 'segmentosum'</i> (X84437)	95.2
<i>C. diphtheriae</i> (X84248)	94.9	<i>C. striatum</i> (X84442)	94.8
<i>C. 'fastidiosum'</i> (X84245)	94.9	<i>C. 'tuberculostearicum'</i> (X84247)	95.2
<i>C. flavescens</i> (X84441)	95.2	<i>C. ulcerans</i> (X84256)	94.3
<i>C. 'genitalium'</i> (X84253)	95.7	<i>C. urealyticum</i> (X84439)	93.5
<i>C. glucuronolyticum</i> (X86688)	93.3	<i>C. variabilis</i> (X53185)	93.4
<i>C. glutamicum</i> (X84257)	93.4	<i>C. vitarumen</i> (X84680)	93.8
<i>C. jeikeium</i> (X84250)	94.0	<i>C. xerosis</i> (X84446)	93.9
<i>C. kutscheri</i> (X81871)	94.0	<i>T. otitidis</i> (X73976)	91.5
<i>C. macginleyi</i> (X80499)	94.8		

<sup>a</sup>The numbers in parentheses are EMBL 16S rRNA nucleotide sequence accession numbers.

droyzed urea more slowly (after 12 h in Christensen's urea broth [13]) than *C. urealyticum* strains (usually positive after only 5 min in Christensen's urea broth). Further enzymatic activities of strain DMMZ 1944 included esterase, esterase lipase, lipase, leucine arylamidase, acid phosphatase, and phosphoamidase whereas valine arylamidase, cystine arylamidase, trypsin, chymotrypsin,  $\alpha$ -galactosidase,  $\beta$ -galactosidase,  $\beta$ -glucuronidase,  $\alpha$ -glucosidase,  $\beta$ -glucosidase, *n*-acetyl- $\beta$ -glucosaminidase,  $\alpha$ -mannosidase, and  $\alpha$ -fucosidase could not be detected. In our experience, the presence of lipase activity is very rarely observed in other true *Corynebacterium* spp. (G. Funke, unpublished data).

Antimicrobial susceptibility testing demonstrated that the strains were susceptible to (minimal inhibitory concentrations given in  $\mu$ g/ml) amikacin (1.0), amoxicillin (0.25), amoxicillin/clavulanic acid ( $\leq 0.0625$ ), azithromycin (0.0625), cefazolin ( $< 0.125$ ), ceftriaxone (0.5), cefuroxime (0.5), chloramphenicol (2.0), ciprofloxacin (0.25), clarithromycin ( $< 0.03$ ), clindamycin (0.125), erythromycin ( $\leq 0.03$ ), fusidic acid (0.125), gentamicin (0.125), imipenem (0.0625), meropenem (0.25), netilmicin (0.125), nor-

floxacin (2.0), oxacillin (1.0), penicillin (0.125), piperacillin (4.0), rifampin (0.0625), teicoplanin (0.5), tetracycline (2.0), tobramycin (0.125), sparfloxacin (0.25), and vancomycin (0.5) but resistant to aztreonam ( $> 64$ ), cefetamet (64), and fosfomicin ( $> 256$ ) [14]. In contrast, *C. urealyticum* is almost ever multi-resistant including resistance to  $\beta$ -lactams, penems, aminoglycosides, macrolides, clindamycin, rifampin, and quinolones [15]. Due to the fact that the unknown *Corynebacterium* exhibited very low MICs for nearly all clinically applied antimicrobial agents it is not unlikely that it, like other fastidious lipophilic corynebacteria, might not be recovered and, therefore, underestimated in clinical specimens after administration of antibiotics. However, the etiologic role of the unknown *Corynebacterium* in our patient with BV remained unclear.

In order establish the precise phylogenetic position and the genealogical distinctiveness of the unknown *Corynebacterium* the gene encoding the 16S rRNA was amplified by PCR and subjected to sequence analysis. The almost complete 16S rRNA gene sequence (ca. 1400 bp) was determined. The unknown bacterium clustered within the genus *Corynebacte-*

*rium sensu stricto*, where it formed a distinct subline (Fig. 1). The non-lipophilic, oxidative *C. auris* and the non-lipophilic, fermentative *C. mycetoides* were the closest phylogenetic neighbors of the unknown *Corynebacterium* showing ca. 3% 16S rRNA sequence divergence (Table 2). Based on the phenotypic and molecular genetic distinctiveness of the unknown bacterium we formally propose a new *Corynebacterium* species, *Corynebacterium lipophiloflavum* sp. nov., for the strain studied.

### 3.1. Description of *Corynebacterium lipophiloflavum* sp. nov.

*Corynebacterium lipophiloflavum* (li'po.phi.lo fla.vum. Gr. n. *lipos*, fat; Gr. adj. *philos*, loving; L. adj. *flavus*, yellow; M.L. adj. *lipophiloflavum*, fat loving and yellow) cells are gram-positive, asporogenous, club-shaped, 1 to 3  $\mu\text{m}$  in length, and non-motile. Colonies are yellowish, circular, convex, slightly dry, and less than 0.2 mm in diameter after 24 h of incubation on SBA but colonies are up to 1 mm in diameter when cells are grown on SBA supplemented with 1% Tween 80. Catalase positive. Acid is not produced from glucose, maltose, sucrose, mannitol, xylose, lactose, and glycogen. Nitrate is not reduced. Urea is slowly hydrolyzed but esculin is not hydrolyzed. The CAMP reaction is negative. Pyrazinamidase, alkaline phosphatase, esterase, esterase lipase, lipase, leucine arylamidase, acid phosphatase, and phosphoamidase activities are detected but pyrrolidonyl arylamidase, valine arylamidase, cystine arylamidase, trypsin, chymotrypsin,  $\alpha$ -galactosidase,  $\beta$ -galactosidase,  $\beta$ -glucuronidase,  $\alpha$ -glucosidase,  $\beta$ -glucosidase, *n*-acetyl- $\beta$ -glucosaminidase,  $\alpha$ -mannosidase, and  $\alpha$ -fucosidase activities are not detected.

The cell wall contains *meso*-diaminopimelic acid. Mycolic acids are present. The main straight-chain saturated fatty acids are palmitic and stearic acids; oleic acid is the predominant unsaturated fatty acid. The DNA base composition is 65 mol% G+C. Isolated from a patient with bacterial vaginosis. The type strain has been deposited in the Culture Collection of the University of Göteborg, Sweden, as CCUG 37336, and in the German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany, as DSM 44291.

### Acknowledgments

The authors thank A. von Graevenitz for a careful review of the manuscript. This study was made possible due to the financial support by the European Community (BIO2-CT943098), the Hochschulverein and the Jubiläumsspende der Universität Zürich.

### References

- [1] Spiegel, C.A. (1991) Bacterial vaginosis. *Clin. Microbiol. Rev.* 4, 485–502.
- [2] Funke, G., von Graevenitz, A., Clarridge III, J.E., and Bernard, K.A. (1997) Clinical microbiology of coryneform bacteria. *Clin. Microbiol. Rev.* 10, 125–159.
- [3] Funke, G., Pascual Ramos, C., and Collins M.D. (1997) *Corynebacterium coyleae* sp. nov., isolated from human clinical specimens. *Int. J. Syst. Bacteriol.* 47, 92–96.
- [4] Funke, G., Martinetti Lucchini, G., Pfyffer, G.E., Marchiani, M., and von Graevenitz, A. (1993) Characteristics of CDC group 1 and group-1-like coryneform bacteria isolated from clinical specimens. *J. Clin. Microbiol.* 31, 2907–2912.
- [5] Funke, G., Alvarez, N., Pascual, C., Falsen, E., Akervall, E., Sabbe, L., Schouls, L., Weiss, N., and Collins, M.D. (1997) *Actinomyces europaeus* sp. nov. isolated from human clinical specimens. *Int. J. Syst. Bacteriol.*, in press.
- [6] Pascual, C., Lawson, P.A., Farrow, J.A.E., Navarro Gimenez, M., and Collins, M.D. (1995) Phylogenetic analysis of the genus *Corynebacterium* based on 16S rRNA gene sequences. *Int. J. Syst. Bacteriol.* 45, 724–728.
- [7] von Graevenitz, A., and Funke, G. (1996) An identification scheme for rapidly and aerobically growing gram-positive rods. *Zentralbl. Bakteriologie* 284, 246–254.
- [8] Bernard, K.A., Bellefeuille, M., and Ewan, E.P. (1991) Cellular fatty acid composition as an adjunct to the identification of asporogenous, aerobic gram-positive rods. *J. Clin. Microbiol.* 29, 83–89.
- [9] von Graevenitz, A., Osterhout, G., and Dick, J. (1991) Grouping of some clinically relevant gram-positive rods by automated fatty acid analysis. *APMIS* 99, 147–154.
- [10] Collins, M.D., and Cummins, C.S. (1986) *Corynebacterium*. In: *Bergey's Manual of Systematic Bacteriology* (Sneath, P.H.A., Mair, N.S., Sharpe, M.E., and Holt, J.G., Eds.), Vol.2, pp. 1266–1276. Williams and Wilkins, Baltimore, MD.
- [11] Pitcher, D., Soto, A., Soriano, F., and Valero-Guillén, P. (1992) Classification of coryneform bacteria associated with human urinary tract infection (group D2) as *Corynebacterium urealyticum* sp. nov. *Int. J. Syst. Bacteriol.* 42, 178–181.
- [12] Soriano, F., Aguado, J.M., Ponte, C., Fernandez-Roblas, R., and Rodriguez-Tudela, J.L. (1990) Urinary tract infection caused by *Corynebacterium* group D2. Report of 82 cases and review. *Rev. Infect. Dis.* 12, 1019–1034.
- [13] Nash, P., and Krenz, M.M. (1991) Culture media. In: *Manual of Clinical Microbiology* (Balows, A., Hausler, Jr., W.J., Her-

- mann, K.L., Isenberg, H.D., and Shadomy, H.J., Eds.), 5th edition, pp. 1226–1288. American Society for Microbiology, Washington, D.C.
- [14] National Committee for Clinical Laboratory Standards (1997) Minimum inhibitory concentration (MIC) interpretive standards ( $\mu\text{g/ml}$ ) for organisms other than *Haemophilus* spp., *Neisseria gonorrhoeae*, and *Streptococcus* spp. NCCLS document M7-A4. National Committee for Clinical Laboratory Standards, Wayne, PA.
- [15] Soriano, F., Zapardiel, J., and Nieto, E. (1995) Antimicrobial susceptibilities of *Corynebacterium* species and other non-spore-forming gram-positive bacilli to 18 antimicrobial agents. *Antimicrob. Agents Chemother.* 39, 208–214.