

*The Astrophysics of Planetary Systems: Formation, Structure, and Dynamical Evolution*

*Proceedings IAU Symposium No. 276, 2010*

A. Sozzetti, M. G. Lattanzi & A. P. Boss, eds.

© International Astronomical Union 2011

doi:10.1017/S1743921311021090

# Stellar noise and planet detection.

## II. Radial-velocity noise induced by magnetic cycles

Xavier Dumusque<sup>1,2</sup>, Christophe Lovis<sup>1</sup>, Stephane Udry<sup>1</sup> and Nuno C. Santos<sup>2</sup>

<sup>1</sup>Observatoire de Genève, 51 ch. des Maillettes, 1290 Sauverny, Switzerland  
email: [xavier.dumusque@unige.ch](mailto:xavier.dumusque@unige.ch)

<sup>2</sup>Centro de Astrofísica, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal

**Abstract.** For the 451 stars of the HARPS high precision program, we study correlations between the radial-velocity (RV) variation and other parameters of the Cross Correlated Function (CCF). After a careful target selection, we found a very good correlation between the slope of the RV-activity index ( $\log(R'_{HK})$ ) correlation and the  $T_{\text{eff}}$  for dwarf stars. This correlation allow us to correct RV from magnetic cycles given the activity index and the  $T_{\text{eff}}$ .

**Keywords.** planetary systems, stars: activity, techniques: radial velocities

### Introduction

The Sun has a 11-years magnetic cycle, during which the activity level (Noyes *et al.* 1984) varies between  $\log(R'_{HK})=-5$  at minimum and  $\log(R'_{HK})=-4.75$  at maximum. This cycle can also be seen looking at the total number of sunspots or the Sun luminosity variation. In a recent paper, Meunier *et al.* (2010) show a correlation between the Sun RV variation and its magnetic cycle. Amplitudes of tenths of meter per second could be induced by such cycles, hiding signals of long period small-mass planets.


Solar type stars share the property of having an external convective envelope. Since upward flows of convection (granules) have a total surface more important than downward flows (intergranules), the stellar spectrum will be bluishifted (Dravins 1982; Gray 2009). When the activity level increases, the number of magnetic features (spots and plages) raise up, and since these regions are known to inhibit the convection due to strong magnetic fields, the stellar spectrum will be shifted to the red. Since the activity level is correlated to the number of magnetic features, a long-term variation of the activity level, as it is the case in magnetic cycles, will induce a long-term RV variation.

### Target selection


To study only long-term activity noise, we select stars that are measured more than 3 years, with a minimum of 5 bins of 3 months (each bin must contains at least 3 measurements). In addition, only stars with  $\log(R'_{HK}) < -4.75$  was selected to address solar-like activity. After this selection, we are left with 91 stars out of 451.

To study potential correlations between the activity index and RV variations (see Dumusque *et al.* (2010), in prep.), we first remove all the planets present in the sample. Then, we select stars in the following way (always for the 3 months binned data):

- Correlation between the CCF Full Width at Half Maximum (FWHM) and  $\log(R'_{HK})$ ,  $R_{FWHM} > 0.5$  or between the CCF BISsector span (BIS) and  $\log(R'_{HK})$ ,  $R_{BIS} > 0.5$ ,
- $\text{Max}(\log(R'_{HK})) > -5$ ,
- Peak to peak variation in  $\log(R'_{HK}) > 0.05$ .



**Figure 1.** Comparison of the long-term RV and activity index ( $\text{Log}(R'_{HK})$ ) variation. The Pearson correlation between the 2 parameters is very good, for 3 month bins,  $R = 0.85$ .



**Figure 2.** Slope of the correlation  $\text{RV}-\log(R'_{HK})$  as a function of the  $T_{\text{eff}}$ . The size of the circle surrounding each star scales with  $R_{RV}$ , the correlation coefficient between RV and  $\log(R'_{HK})$ . Small size if  $R_{RV} > 0.5$  and large size if  $R_{RV} > 0.75$ . Green numbers correspond to stars which have  $R_{FWHM} > 0.5$  or  $R_{BIS} > 0.5$  and red numbers (71 and 23), to stars with no well defined correlation.

After this second selection, only 31 stars are left. However, a nice correlation appears when we plot these stars in a graph representing the slope of the correlation  $\text{RV}-\log(R'_{HK})$  as a function of the  $T_{\text{eff}}$  (see Fig 2). The higher the slope, the more the RV will be affected by magnetic activity. Therefore, RVs of early-Kdwarfs are less affected by magnetic cycles than RVs of early-G dwarfs, making early-K dwarfs optimal targets for very small mass planets surveys. In addition, this relation allow us to correct RVs from the magnetic cycle given the activity index and the  $T_{\text{eff}}$  of the star.

**References**

- Noyes, R., *et al.* 1984, *ApJ*, 279, 763  
Meunier, N., Desort, M., & Lagrange, A.-M. 2010, *A&A*, 512, A39  
Dravins, D. 1982, *ARAA*, 20, 61  
Gray, D. F. 2009, *ApJ*, 697, 1032