

Determination of digestible energy values and fermentabilities of dietary fibre supplements: a European interlaboratory study *in vivo*

BY G. LIVESEY¹, T. SMITH¹, B. O. EGGUM², I. H. TETENS^{2*}, M. NYMAN³,
M. ROBERFROID⁴, N. DELZENNE⁴, T. F. SCHWEIZER⁵ AND J. DECOMBAZ⁵

¹ Institute of Food Research, Norwich Laboratory, Norwich Research Park, Colney,
Norwich NR4 7UA

² National Institute of Animal Science, Research Centre Foulum, PO Box 39, 8830 Tjele, Denmark

³ The Chemical Centre, Lund University, PO Box 124, 221 00 Lund, Sweden

⁴ Department of Pharmaceutical Science, Catholic University of Louvain, UCL 7369, 1200
Brussels, Belgium

⁵ Nestlé Research Centre, Nestec Ltd., Av. Nestlé 55, 1800 Vevey, Switzerland

(Received 23 March 1994 – Revised 6 June 1994 – Accepted 7 June 1994)

The performance of methods to determine energy conversion factors for dietary fibre (DF) supplements and fermentability (*D*) values of their non-starch polysaccharides (NSP) was investigated. Heats of combustion, digestible energy (*DE*) and *D* values were determined on five DF supplements in five European laboratories on five separate occasions. In each instance the DF supplements were fed to juvenile male Wistar rats at two doses, 50 and 100 g/kg basal diet, for 3 weeks with food and faeces collected in the 3rd week. Among-laboratory variations in heats of combustion (ΔH_c) were < 2%. *DE* values (kJ/g dry weight) at the upper and lower doses respectively were: 10.4 and 9.9 for a high-methoxyl apple pectin, 9.5 and 9.4 for a sugar-beet DF supplement, 12.2 and 12.7 for soyabean DF supplement, 3.8 and 4.0 for maize bran, and 0.3 and 0.3 for Solka-floc cellulose. Variations among laboratories, among occasions and among animals were < 1, < 2 and < 2.5 kJ/g respectively. The among-occasion: among-laboratory variance ratio for *DE* was 0.5, suggesting the method performed equally well in all laboratories. There was no evidence of learning or fatigue in the performance of the method. *D* values were also independent of dose and at the high and lower doses were: pectin 0.92 and 0.95, sugar-beet NSP 0.68 and 0.68, soyabean NSP 0.86 and 0.88, maize bran 0.17 and 0.18, cellulose 0.07 and 0.06. Among-laboratory variance tended to increase with decreasing fermentability and ranged from 0.03 to 0.18. The *DE* and *D* data were not significantly different from a previously proposed relationship $DE = 0.7 \times \Delta H_c \times D$, where ΔH_c is the heat of combustion of the supplement. We conclude that while the among-laboratory variation in the *D* of difficult-to-ferment NSP is too large for the reliable prediction of energy value the method for the direct determination of *DE* is both reproducible and repeatable, that *DE* is independent of dosage of DF supplement up to 100 g/kg diet, and that it is safe to discriminate between energy values with a precision of 3 kJ/g. The conversion of both *DE* and *D* to net metabolizable energy for the purpose of food labelling, tables and databases is described.

Food energy value: Fermentation: Non-Starch polysaccharide: Dietary fibre

Non-starch polysaccharides (NSP) are the main constituents of dietary fibre (DF) or unavailable carbohydrate (British Nutrition Foundation, 1990). The contribution of these carbohydrates to whole-body energy metabolism in humans and rats is small but is measurable (Harley *et al.* 1989; Johnson *et al.* 1990; Livesey, 1990) and is supplied via

* Present address: Research Department of Human Nutrition, The Royal Veterinary and Agricultural University, Rolighedsvej 25, DK-1958 Fredriksberg C, Denmark.

fermentation in the large intestine. How well the determinations of fermentability (D) of NSP and energy value of DF supplements compare in different laboratories and whether there is a relationship between the two values are questions addressed by the present study.

Human studies performed in different laboratories with various amounts and sources of unavailable carbohydrates for which energy values have been derived have yielded varied results (Southgate & Durnin, 1970; Göransson *et al.* 1983; Göransson & Forsum, 1987; Wisker *et al.* 1988; Miles, 1990). Based on an analysis of the world literature on over thirty such mixed diets eaten by humans in the Western world, the unavailable carbohydrate contributes an average of about 8 kJ/g to digestible energy (DE ; Livesey, 1991). However, different forms or sources of unavailable carbohydrates were suggested to have values in the range 0–12 kJ/g (Livesey, 1992). The precise value for any one source was proposed to be related to its susceptibility to fermentation (Livesey, 1990, 1992; Roberfroid *et al.* 1993). Nevertheless, because energy is lost to faeces, predominantly as unfermented carbohydrate and as bacterial biomass, all these DE values (DEV_g) are less than the 17.5 kJ/g value for available starch. A further amount of energy is lost during fermentation as heat and combustible gases; so it is thought that only about 50% of the unavailable carbohydrate fermented is recoverable as net metabolizable energy (NE_m ; British Nutrition Foundation, 1990; Livesey, 1992). A further possible route of energy loss is to urine, but such losses are usually ignored as they are either negligible (Livesey, 1990) or may be slightly reduced by unavailable carbohydrate (Brown & Livesey, 1994).

While knowledge about the availability of energy from unavailable carbohydrates has become clearer in recent years, it is not reflected in European Food Labelling Regulations, which have not yet assigned an energy value to this class of dietary fuel (European Council Directive, 1990). This situation contrasts with undigested oligosaccharides and sugar alcohols that partly undergo fermentation in the large intestine. For example, sugar alcohols have been assigned an average energy value of 10 kJ/g (European Council Directive, 1990).

Acceptance of energy values for unavailable carbohydrate in food labelling regulations, tables and databases requires energy evaluation methodology that is recognized by academic and commercial organizations and by regulatory authorities. The performance of such methods, therefore, needs to be tested and its use standardized among different laboratories. The degree of agreement between laboratories on the energy values of different preparations provides the limits on the accuracy and precision with which energy values can be assigned. However, until now no such interlaboratory study of method performance has been reported.

The primary aim of the present study was to determine how reproducible the determinations of energy value and D would be in different laboratories throughout Europe and to determine the replicability of these determinations within laboratories. The second aim was to establish whether the extent of fermentation of NSP predicts the availability of energy from NSP-rich DF supplements. The third aim was to provide standard energy and D values for five DF products and their NSP content respectively, which would help in the development of an *in vitro* method that predicts *in vivo* D and energy values (see accompanying paper, Barry *et al.* (1995)).

The rat was chosen for these investigations since we concluded that observations with the rat corresponded reasonably closely to observations with humans when NSP fermentability was determined (Nyman *et al.* 1986; Livesey, 1992; Bach-Knudsen *et al.* 1994). Five European laboratories were invited to participate, from Belgium, Denmark, England, Sweden, and Switzerland. Each laboratory determined the D of NSP and DEV_g of each of five different DF supplements. The supplements had been selected for their minimal starch content and wide range of D from about 0 to about 100%. The sources chosen were a high-

methoxyl pectin from apple, sugar-beet DF supplement, soyabean DF supplement, maize bran and Solka-floc cellulose. Because there is little information in the literature on the dependency of energy values and *D* on dietary dosage with DF supplements, each laboratory made these determinations at two dose levels. Further, by making replicate observations on five consecutive occasions an assessment was made of whether increasing experience (learning or fatigue) in execution of the methods affected the results. This was important since the protocol and analytical methods used were new to most of the researchers and technical staff involved.

METHODS

Materials

Apple pectin was 'slow-set' high-methoxyl pectin kindly provided by Sanofi Bio-Industries (66 Avenue Marceau, 75008, Paris, France). Soyabean DF was 'FIBRIM 2000 (red)' from Protein Technologies International (16A, Princewood Road, Earstree Industrial Estate, Corby, Northants). Sugar-beet DF was Betafibre kindly provided by British Sugar plc, Peterborough, Cambs. Maize bran was from Honeyville Grain Inc., 4380W, 2100 South, Salt Lake City, Utah 84120, USA. Cellulose was Solka-floc, grade BW 2030, from Jurgenson and Wettre Ltd., Johnson House, Wellington Road, Wokingham, Berks.

Diets

The composition of the basal diet to which the DF supplements were added was as shown in Table 1. The basal diet was free of NSP and test diets were prepared by adding DF supplements at the rates of 50 and 100 g/kg basal diet. All the diets were prepared and distributed from one laboratory.

Animals

Male Wistar rats (70–130 g) obtained from local suppliers were used: for Belgium and Switzerland they were from Iffa Credo, Lyon, France; for Denmark from Mollgaards Breeding Centre, Lille Skensved, Denmark; for England from A. Tuck and Son, Battlebridge, Essex; and for Sweden from ALAB, Sollentuna, Sweden. Animals were allocated randomly to dietary treatments, kept at 20–24° and relative humidity 40–60%, had light for 12 h daily from approximately 06.00 hours and had water *ad lib*.

Experimental design

Each of the five laboratories determined the DEV_s (equations 1 and 2, p. 292) of the five DF preparations at two dietary doses on five occasions. On each occasion two rats per DF source and dose received a test diet and four rats the control diet. This produced an experimental design with 500 observations (five laboratories \times five DF supplements \times two doses \times five occasions \times two rats). Animals from the first three occasions were also used to determine *D* for NSP (equation 3, p. 293) and produced a total of 300 observations ($5 \times 5 \times 2 \times 3 \times 2$).

On each occasion each laboratory housed twenty-four rats singly in cages with wire-mesh floors. Feed (stock feed as usually used in each laboratory) was withdrawn from animals for 16 h before feeding the experimental diets. Experimental diets were provided at 10.00 hours each day for 21 d; basal diet alone was fed to four rats and each of the five DF-supplemented diets at each dosage was fed to two rats. Spilt food was collected daily for each animal and a fresh weight of the amount spilt was returned to the animal in the next day's ration. Each day the animals received spill-weight from the previous day plus 15 g basal diet or 15.8 g low-dose-DF diet or 16.5 g high-dose-DF diet. In some laboratories the

Table 1. *Composition of the experimental diets (g/kg)*

Basal diet		
Maize starch (moisture approximately 100 g/kg)*		330
Sucrose (moisture approximately 10 g/kg)		360
Casein (moisture approximately 50 g/kg)†		200
DL-Methionine		2
Maize oil‡		80
Vitamin mixture§		20
Mineral mixture		40
Supplemented diets		
Basal diet		1000
DF supplement	either	50
	or	100

* The starch was 'Snowflake' maize starch from Corn Products Ltd., Manchester; in addition there was approximately 18 g in the vitamin mix.

† Casein was edible casein, mesh 30, from G. Fiske and Co. Ltd., Richmond, Surrey.

‡ The maize oil was Mazola from CPC International, Esher, Surrey.

§ Provided the following amounts in the basal diet (mg/kg): nicotinic acid 60, cyanocobalamin in mannitol 50, calcium D-pantothenate 40, thiamine hydrochloride 10, riboflavin 10, folic acid 10, pyridoxine 10, D-biotin 1, vitamin K₁ 2, Rivomix E-50 (containing 7.5 mg vitamin E, Roche) 150, Rivomix A-500 (containing 3.75 mg vitamin A, Roche) 25, Rivomix D₃-500 (containing 0.19 mg vitamin D₃, Roche) 15, choline bitartrate 1800, maize-starch carrier 17800.

|| Provided the following amounts in the basal diet (g/kg): CaHPO₄ 13, CaCO₃ 8.2, KCl 7.04, Na₂HPO₄ 7.4, MnSO₄·H₂O 0.18, MgSO₄·H₂O 4, ZnCO₃ 0.1, FeSO₄·7H₂O 0.144, CuSO₄ 0.023, KIO₃ 0.001.

amount of diet spilt early after introduction of the DF supplements was partly discarded because the diet provided an amount that exceeded voluntary intakes, but during the period of faecal collections spillages were usually minimal.

Faeces were collected before feeding during the last 7 d of the 21 d of dietary treatment and were pooled for each animal. Faecal collections were frozen at -20° daily.

Analysis

The basal diet, DF supplements and faeces were analysed for gross energy and total NSP. Gross energy was the heat of combustion determined by adiabatic bomb calorimetry using benzoic acid thermochemical standard (for review, see Brown, 1993). Total NSP was the sum of the neutral and acid NSP. Neutral NSP was measured by the colorimetric method of Faulks & Timms (1985) with a glucose standard. Acidic NSP (uronic acid) was measured by the method of Scott (1979) with a galacturonic acid standard. The neutral NSP and uronic acids were measured without removal of the negligible amounts of available or resistant starches as in the apple pectin (Barry *et al.* 1995) by dimethyl sulphoxide treatment. Each value was expressed on a per g dry weight basis after freeze-drying. The supplements were also assayed for total dietary fibre (TDF) by the Association of Official Analytical Chemists' method (Prosky, 1985; Prosky *et al.* 1988).

Calculations

DEV_s were calculated using equations 1 and 2:

$$DEV_s = \Delta H_c - \Delta FE, \quad (1)$$

$$\Delta FE = [(E_{tf}/M_{td}) - (E_{cf}/M_{cd})]/(M_s/M_{td}), \quad (2)$$

where ΔH_c is the heat of combustion of the supplement, ΔFE is the increase in faecal energy per g DF supplement ingested, E is gross energy, M is mass, and the subscripts were: t , test;

c, control; *f*, faeces; *d*, basal diet; *s*, supplement. An example calculation and discussion has been given (Livesey, 1989).

D of the NSP was calculated as its 'digestibility', a method that assumes NSP was used only by fermentation:

$$D = (I - F)/I, \quad (3)$$

where *I* is intake of NSP (a product of supplement intake multiplied by the proportion of the supplement that was NSP) and *F* was NSP excretion in faeces during the balance period (after deduction of the NSP 'apparently' excreted from the basal diet alone).

DEV_s and *D* were converted to *NE_m* according to information from Livesey (1990, 1992) using equations 4 and 5 respectively.

$$NE_m = 0.7 DEV_s \quad (4)$$

$$NE_m = 0.5 \times \Delta H_c \times D. \quad (5)$$

In these equations, *DEV_s*, ΔH_c and *D* were as described for equations 1 and 3.

Statistics

For each DF supplement at each dose a factorial analysis was undertaken with occasions and laboratories represented by blocks in multiple regression. Each animal was the unit of measurement and the residual variation was ascribed to variation among animals (*SD_A*). In keeping with methods-performance assessment (Boyer *et al.* 1985), terms for interactions between factors were excluded and variation among occasions (*SD_O*, often referred to as replicability) and variation among laboratories (*SD_L*, often referred to as reproducibility) were obtained as the standard deviation about the mean determination for each occasion and laboratory respectively. LRD was the largest residual difference or deviation of any one laboratory mean value from the all-laboratory mean value.

Factorial analysis including all DF supplements at each dose in addition to each laboratory and each occasion was also undertaken by multiple regression to obtain inter-laboratory differences and among-laboratory variation and inter-occasional differences and among-occasion variation over all the energy determinations. Again terms for interactions were excluded from the analysis. For the energy values, homogeneity of variances within each factor (DF supplement, dose, laboratory and occasion) were evident from *F* ratios.

All data were retained in the statistical analyses unless they were laboratory means that either were impossible observations (e.g. highly negative fermentabilities) or were significant outliers in Dixon's *r*₁₂ test (Barnett & Lewis, 1978), which permitted data from one laboratory in five to be excluded from a statistical analysis (Boyer *et al.* 1985). The excluded data were few and are identified by parentheses in the tables.

Significance of difference between the estimates for each DF supplement of *NE_m* converted from *DEV_s* and *D* were based on the standard error of difference (SED). The SED was derived from the variance among animals (*SD_A*) according to $\sqrt{[(s_1^2/n_1) + (s_2^2/n_2)]}$, where *s*₁ was the *SD_A* for *D* with *n*₁ 60 determinations including both treatment doses and *s*₂ was the *SD_A* for *DEV_s* with *n*₂ 100 determinations including both doses.

RESULTS

Heats of combustion

Each of the five laboratories determined the heat of combustion of each of the five DF products (Table 2). The mean values ranged from 15.6 kJ/g for pectin to 19.0 kJ/g for the maize bran. Among-laboratory variation (*SD_L*) of the mean determinations for the

Table 2. *Heats of combustion (kJ/g dry weight) for five dietary fibre (DF) supplements determined in five laboratories participating in the study**

DF supplement	Laboratory					Mean	SD _L	LRD
	A	B	C	D	E			
Pectin	15.9	15.9	15.5	15.8	15.2	15.6	0.2	0.4
Sugar beet	17.7	17.5	17.7	17.6	17.4	17.6	0.1	0.2
Soyabean	17.3	17.3	16.6	16.8	16.8	16.9	0.3	0.4
Maize bran	19.5	19.1	18.6	18.9	18.9	19.0	0.3	0.4
Solka-floc cellulose	17.3	17.4	17.2	17.1	17.0	17.2	0.3	0.2

SD_L, standard deviation among laboratories; LRD, largest residual deviation of an individual laboratory mean value from the all-laboratory mean value.

* For details of supplements and procedures, see pp. 291–293.

Table 3. *Digestible energy values (kJ/g dry weight) for five dietary fibre (DF) supplements at two doses in five laboratories participating in the study**

DF supplement	Dose	Laboratory					Mean	SD _L	SD _O	SD _A	LRD
		A	B	C	D	E					
Pectin	L	12.0	11.1	9.3	10.3	9.4	10.4	1.1	0.5	1.9	1.6
	H	10.4	(8.4)†	10.2	10.7	9.9	10.3	0.4	0.7	1.8	0.4
Sugar beet	L	9.8	11.4	9.9	9.4	7.0	9.5	1.4	0.6	1.8	2.5
	H	10.0	11.8	9.2	9.3	6.6	9.4	1.9	0.2	1.3	2.8
Soyabean	L	13.3	11.9	11.3	13.6	11.1	12.2	1.0	0.9	1.7	1.4
	H	12.9	14.5	11.8	12.3	11.8	12.7	1.1	0.4	1.6	1.8
Maize bran	L	5.7	3.2	4.7	3.1	2.3	3.8	1.2	0.8	1.6	1.9
	H	5.5	3.4	3.8	4.7	2.7	4.0	1.0	0.8	1.8	1.5
Solka-floc cellulose	L	0.7	-0.4	0.6	0.8	-0.4	0.3	0.5	0.6	2.2	0.7
	H	0.7	0.4	-0.5	0.0	0.8	0.3	0.5	0.9	1.7	0.8

L, lower dose of 50 g DF supplement/kg basal diet; H, higher dose of 100 g/kg basal diet; SD_L, standard deviation among laboratories; SD_O, standard deviation among occasions; SD_A, standard deviation among animals; LRD, largest residual difference or deviation of an individual laboratory mean value from the all-laboratory mean value.

* For details of supplements and procedures, see pp. 291–293.

† The value is a statistical outlier.

individual DF supplements were obtained to within 0.3 kJ/g, which corresponds to a coefficient of variation of < 2%. All the laboratories obtained values within 0.4 kJ/g (3%) of the all-laboratory mean for each DF supplement (LRD, Table 2).

Digestible energy values

Each of the five laboratories determined the *DEV*_s of each of the five DF supplements at doses of 50 and 100 g/kg basal diet (Table 3). For all the supplements the mean values were independent of dosage and ranged from about 0 kJ/g for Solka-floc cellulose to 12 kJ/g for the soyabean-DF supplement. As examples of the data obtained, those observations for the DF supplements with the highest and lowest energy values at the lower of the two doses are shown in Fig. 1.

Variations in the *DEV*_s among laboratories (SD_L), among occasions (SD_O) and among animals (SD_A) were independent of the amount of DF supplement in the diet (Table 3).




Fig. 1. Examples of the digestible energy value data obtained for (a) Solka-floc cellulose supplement and (b) soyabean-DF supplement, both at 50 g/kg basal diet. Values are the mean observations for two rats determined on each occasion (nos. 1–5) in each laboratory (A–E). —, Mean values for each laboratory. Values are shown with the lowest to the left. For details of supplements and procedures, see pp. 291–293.

Limiting the precision of the determinations was variation among animals (SD_A), which tended to be higher than SD_L and SD_O . Replicability of the determinations among occasions (SD_O) was < 1 kJ/g ($< 6\%$ of the heat of combustion), whereas reproducibility of the determinations among laboratories (SD_L) was less than 2 kJ/g and variation among animals (SD_A) was less than 2.5 kJ/g.

All the laboratories obtained DEV_s for the individual DF supplements that were within 3 kJ/g (LRD, Table 3) of the all-laboratory mean values.

Factorial analysis incorporating all the DF supplements at both doses showed (Table 4) that overall there were no significant differences in DEV_s between occasions; the value for each occasion was within 0.5 kJ/g of the mean. There were significant ($P < 0.05$) differences between laboratories, which reached as high as 2 kJ/g or 12% of the heat of combustion (Table 4).

TDF and NSP contents of the DF supplements

The DF supplements consisted mainly of Prosky TDF (Prosky, 1985; Prosky *et al.* 1988) or Faulks & Timms (1985) total NSP (Table 5). There was agreement between the two

Table 4. *Inter-occasional and inter-laboratory differences in digestible energy value over all dietary fibre supplements and doses (kJ/g dry weight)**

Occasion	Occasional difference (kJ/g)	Laboratory	Laboratory difference†
First	0.25	A	1.00
Second	0.29	B	0.41
Third	0.22	C	-0.06
Fourth	-0.44	D	-0.20
Fifth	-0.36	E	-1.11
LSD	0.64	LSD	0.64
df	485	df	485
SD _O	0.4	SD _L	0.8
CV‡	2.3	CV‡	4.6

LSD, least significant difference at $P < 0.05$; SD_O, inter-occasional variance; SD_L, inter-laboratory variance; CV, coefficient of variation.

* For details of procedures, see pp. 291–293.

† Each difference (from the overall mean) included 100 determinations (i.e. not excluding the outlier shown in Table 3).

‡ Percentage of the heat of combustion.

Table 5. *Total dietary fibre (TDF) and total non-starch polysaccharide (NSP) content of the dietary fibre (DF) supplements (g/g dry weight)†*

Variable... Procedure...	TDF (g/g dry wt) Prosky (1985) and Prosky <i>et al.</i> (1988)		NSP (g/g dry wt) Faulks & Timms (1985)	
DF supplement				
Pectin	0.70	0.67	0.94	31
Sugar beet	0.78	0.77	0.67	13
Soyabean	0.82	0.81	0.69	9*
Maize bran	0.88	0.88	0.69	8**
Solka-floc cellulose	1.00	1.00	1.03	3

The total NSP values differed significantly from those for TDF: * $P < 0.05$, ** $P < 0.01$.

† TDF determinations are from two separate laboratories with previous experience of the methods. Total NSP are all-laboratory mean values and among-laboratory standard deviations with data from the five laboratories. For details of procedures, see pp. 291–293.

laboratories that determined TDF. The colorimetric method of Faulks & Timms (1985) showed values that differed from the TDF method and the difference was significant ($P < 0.05$) for both the soyabean supplement and the maize bran. Except for the Solka-floc cellulose, variances with the Faulks & Timms (1985) analysis were high and pectin gave particularly variable results.

Fermentability

Each of the five laboratories determined *D* for each of the five sources of NSP at two dietary dosages (Table 6). For all the sources of NSP the values were independent of dosage and mean values ranged from about zero for Solka-floc cellulose to about 0.9 for pectin.

Variations in *D* values among laboratories (SD_L), among occasions (SD_O) and among animals (SD_A) were independent of the amount of NSP source in the diet. Replicability of the determinations among occasions (SD_O) was < 0.1 , but reproducibility of the

Table 6. Fermentability (g/g dry weight) of non-starch polysaccharides in the five dietary fibre (DF) supplements determined at two doses in five laboratories participating in the study*

DF supplement	Dose	Laboratory					Mean	SD _L	SD _O	SD _A	LRD
		A	B	C	D	E					
Pectin	L	0.94	0.86	0.94	0.93	0.93	0.92	0.03	0.05	0.05	0.06
	H	0.94	(0.63)†	0.96	0.94	0.96	0.95	0.02	0.04	0.08	0.01
Sugar beet	L	0.72	0.70	0.80	0.56	0.64	0.68	0.09	0.02	0.07	0.12
	H	0.74	0.73	0.69	0.63	0.61	0.68	0.06	0.01	0.09	0.07
Soyabean	L	0.93	0.83	0.91	0.79	0.86	0.86	0.05	0.02	0.05	0.07
	H	0.94	0.91	0.91	0.75	0.88	0.88	0.07	0.03	0.10	0.13
Maize bran	L	0.30	0.02	0.23	0.30	0.04	0.17	0.17	0.09	0.11	0.15
	H	0.43	0.07	0.20	-0.02	0.06	0.15	0.18	0.05	0.11	0.28
Solka-floc cellulose	L	0.22	-0.13	0.19	(-0.50)†	0.00	0.07	0.16	0.05	0.12	0.20
	H	0.19	-0.10	0.11	(-0.26)†	0.05	0.06	0.13	0.08	0.12	0.16

L, lower dose of 50 g DF supplement/kg basal diet; H, higher dose of 100 g/kg basal diet; SD_L, standard deviation among laboratories; SD_O, standard deviation among occasions; SD_A, standard deviation among animals; LRD, largest residual deviation of an individual laboratory mean value from the all-laboratory mean value.

† Values did not contribute to the statistical analysis being excluded as either unreasonably negative (-0.50, -0.26) or as a statistical outlier (0.63).

* For details of supplements and procedures, see pp. 291-293.




Fig. 2. The relationship between the fermentability (D) of non-starch polysaccharide (NSP) and digestible energy value (DEV_s). The line shows the anticipated relationship $DEV_s = 0.7 \times D \times \Delta H_c$, where ΔH_c is the heat of combustion of the supplement (Livesey, 1990). (●, ○), Pectin; (●, ⊗), Sugar-beet NSP; (▲, △), Solka-floc cellulose; (■, ⊗), maize bran; (■, □), soyabean NSP; ○, ⊗, △, ⊗, □, 50 g/kg basal diet; ●, ▲, ■, 100 g/kg basal diet. Individual values and associated standard deviation (SD_A) are from Tables 3 and 6. For details of supplements and procedures, see pp. 291-293.

Table 7. Conversion of fermentabilities (D) and digestible energy values (DEV_s) to net metabolizable energy values for each dietary fibre (DF) supplement†

DF supplement	ΔH_c (kJ/g)	D	DEV_s (kJ/g)	Net metabolizable energy‡		SED
				$0.5 \times \Delta H_c \times D$ (kJ/g)	$0.7 DEV_s$ (kJ/g)	
Pectin	15.6	0.93	10.4	7.2	7.3	0.21 ^{NS}
Sugar beet	17.6	0.68	9.5	5.9	6.7	0.20 ^{**}
Soyabean	16.9	0.87	12.4	7.3	8.7	0.19 ^{**}
Maize bran	19.0	0.16	3.9	1.5	2.7	0.22 ^{**}
Solka-floc cellulose	17.2	0.06	0.3	0.5	0.2	0.24 ^{NS}

SED, standard error of difference with > 68 df; NS, not significant. The difference between net metabolizable energy values calculated by the two methods was significant: ** $P < 0.01$.

† For details of supplements and procedures, see pp. 291–293.

‡ The conversion procedures are based on Livesey (1990, 1992), where ΔH_c is the heat of combustion (see equations 4 and 5, p. 293).

determinations among laboratories (SD_L) was higher at < 0.2. Variation among animals (SD_A) tended to fall between SD_O and SD_L . These variances tended to be less for the NSP sources that were readily fermented than for those that were not.

Factorial analysis incorporating all the NSP sources at both doses showed that overall there were no inter-occasional or inter-laboratory differences in D values, nor was there any trend for a change in D with the number of occasions that the determinations were made (values not shown); however, variation in these values was sometimes large (Table 6).

Digestible energy value related to fermentability

The mean DEV_s of the DF preparations correlated with the mean D of their constituent NSP (r 0.92). All data fell within 3 kJ/g of the line expected (Fig. 2) when assuming (Livesey, 1990) the DEV_s to equal 0.7 multiplied by both the general heat of combustion and fermentability of the NSP.

Net metabolizable energy values

The conversion of mean DEV_s of the DF supplements and D of NSP to NE_m is shown in Table 7. Values obtained by the two approaches were within 1.5 kJ/g of each other (< 10% of their heat of combustion) and because of the large number of observations some differences were significantly different statistically, as indicated.

DISCUSSION

From an early view that unavailable carbohydrate (as Southgate dietary fibre) contributes no energy to the human diet (Southgate & Durnin, 1970) has evolved a current view that unavailable carbohydrate in mixed human diets (as determined by several different methods) contributes about 8 kJ or 2 kcal DE/g substrate (British Nutrition Foundation, 1990; Livesey, 1990, 1991, 1992). It should be emphasized that the assignment of a zero value was an empirical view as Southgate, Durnin and many other preceding authors were aware of fermentation in the large bowel and the possibility that short-chain fatty acids could provide energy. Southgate confirms the current view commenting (discussion following Wisker & Feldheim, 1992) that 'the 2 kcal/g [digestible energy] value is about right [for mixed diets] but for individual constituents the average value may either

overestimate or underestimate the energy [(Livesey, 1990, 1992; Roberfroid *et al.* 1993)] and that there is a need for methods to assess the energy values of ingredients so that specific conversion factors are available to food producers who wish to use the material as an ingredient'.

The revised thinking on the energy value of unavailable carbohydrates was reached for four main reasons which have been cited and discussed elsewhere (British Nutrition Foundation, 1990; Livesey, 1990, 1992). One is that provided an examination is made of a large enough data-set that contains varied sources and amounts of unavailable carbohydrate the energy values determined by experiment agree with theoretical expectations based on the extent and stoichiometry of fermentation. In keeping with this statement the present study incorporates a large data-set by running identical protocols in five different European laboratories and includes various sources and doses of DF supplements.

There are a number of differences between the present work and that in humans reviewed previously. The present study has focussed on DF supplements rich in NSP, with either no or minimal starch content, rather than on the unavailable carbohydrates in mixed diets. Also, five DF supplements from different sources representing NSP of differing *D* were included rather than a mixture from different foods in which the *D* of unavailable carbohydrate tends towards an average value of about 0.7, at all intakes of unavailable carbohydrate (Livesey, 1990). Further, the rat was used in place of humans since *D* of NSP in the two species appear to be similar (Nyman *et al.* 1986; Livesey, 1992; Bach-Knudsen *et al.* 1994) and an investigation of methods performance in humans of the present complexity would have been impractically expensive.

The DEV_s of the Solka-floc cellulose and sugar-beet-DF supplements have been determined previously at 0 and 9 kJ/g respectively (Harley *et al.* 1989; Johnson *et al.* 1990); the 0.3 and 9.5 kJ/g respectively obtained at present agree with those values (Table 3).

It is very clear that for the five DF supplements investigated neither the DEV_s nor the *D* of the NSP depend on the amount in the diet between 50 and 100 g DF supplement/kg basal diet. Indeed, the similarities in DEV_s at each dose are within 3% of the heat of combustion (i.e. within 0.5 kJ/g) and differences in *D* between doses also did not exceed 0.03 (3%; Tables 3 and 6). The rat, therefore, was well within its capacity to accommodate the higher intake of NSP and did so within the 2 weeks adaptation period before the energy balance measurements. Moreover, within experimental error, the DEV_s of the five NSP-rich products could be anticipated from their *D* and the theoretical relationship between the two values suggested previously (Livesey 1990, 1992). We conclude, therefore, that doses in the range 50–100 g DF supplement/kg basal diet are suitable for the determination of energy value and that *D* is a reasonable index of the energy value.

The procedure for the determination of energy value performed reasonably well, but that for the determination of *D* was not good. The less than 2% among-laboratory variation in heat of combustion compares with expectations of approximately 2% for the analysis of a 100% 'pure' material by any satisfactory analytical method (Horwitz *et al.* 1980). The among-laboratory variation in the *DE* evaluation procedure, usually about 1 kJ/g (Table 3) or 6% of the heat of combustion, is twice that expected of an analytical procedure with a material of 0.05–0.1 'purity', probably due to biological variation. It should be noted that any differences in animals from the different suppliers used by the participating laboratories, although all having the same specification, will contribute to the among-laboratory variance.

The ratio, among-replicate variation within laboratories: among-laboratory variation is an indicator of the quality of the data in analytical methods performance, typically values in the range 0.5–0.7 are expected (Horwitz *et al.* 1980). Variation within laboratories,

among animals (SD_A) is higher than variation among laboratories (SD_L), by about $\times 1.6$ (Table 3); this value is above the 0.5–0.7 expected of an analytical method. Again biological variation is probably the limiting factor in the precision of the method and the explanation of the high $SD_A:SD_L$ value. However, biological variation has less influence on the among-occasion:among-laboratory variance ratio and over all the DF supplements and doses the ratio was 0.5 (2.3/4.6; Table 4). This ratio compares with mean values of 0.5–0.7 expected of analytical methodology that is applied equally well in the participating laboratories (Horwitz *et al.* 1980).

$SD_A:SD_L$ for the present *D* determinations is about 1.6 (Table 6), and similar to that for the *DE* determinations (Table 3). Again the value is high probably because of biological variation, and possibly also because of difficulties in the execution of the analytical procedure for NSP. $SD_O:SD_L$ tended to be below 1 (except for pectin) indicating tighter control over the determinations within laboratories than among them.

While the analysis of NSP is not the focus of the present work certain pertinent points are made using apple pectin and Solka-floc cellulose as examples that help interpretation of the *D* data. With apple pectin the among-laboratory variation in NSP determination was unacceptably high at 0.3 (30%) (Table 5) reflecting difficulties among the participating laboratories when analysing uronic acids in this particular source by the method of Scott (1979). However, variation in the determined *D* for pectin among laboratories, among occasions and among animals (Table 6) was small for two reasons. First, *D* is a ratio of NSP determinations so any analytical bias within laboratories is largely cancelled, and second, the amount of NSP in the faeces is small giving rise to little analytical variation. The situation with Solka-floc cellulose is different. The among-laboratory variation in NSP determination by the method of Faulks & Timms (1985) in the cellulose supplement was small, 0.03 (3%) and very close to the 0.02 expected for the analysis of a 'pure' substance (Horwitz *et al.* 1980). However, the among-laboratory and, to a lesser extent, the among-animal variations in the determined *D* for Solka-floc cellulose were high (0.1–0.2; Table 6) indicating either a high biological variation (including variation among animals from the different suppliers) or a high analytical variation when measuring large amounts of substrate in faeces. These different circumstances probably explain the low variation about the *D* of the more-easily-fermentable NSP and the high variation about the *D* of poorly-fermentable NSP. A different result might have been reached had either a more sophisticated procedure been employed for the NSP determinations, such as in some previous studies on fermentation measured with GLC analysis (Nyman & Asp, 1985, 1988), or had the researchers learned more about the colorimetric methods before use here. Nevertheless, the finding of a large among-laboratory variation in *D* of the difficult-to-ferment NSP indicates that the present procedure is not yet sufficiently well developed for predicting energy value, which is better determined directly.

We have determined DEV_s , but food labelling practice with other fermentable carbohydrates and sugar alcohols uses NE_m values (European Council Directive, 1990). Conversion of *D* and DEV_s to NE_m values (Table 7) is simpler than direct determination (Livesey, 1992). This is because with several test products it is not practical to determine the sum of the energy expended on the heat of fermentation, the additional heat generated when short-chain fatty acids are oxidized in place of glucose and the transfer of energy to combustible gases. The two conversions to NE_m give similar values, within 2 kJ/g of each other (Table 7), so that in principle either approach would seem satisfactory. The small discrepancies in the values obtained by the two approaches could well arise because of combustible impurities in the dietary fibre preparations; for example, with maize bran the higher NE_m calculated from DEV_s than from *D* may be due to small amounts of starch being present. Such may also explain the departure of observations from the expected

relationship between DEV_s and D in Fig. 2. Nevertheless, with the present methods the conversion of DEV_s to NE_m would be more reliable than prediction from D because of the large among-laboratory variations in the D of difficult-to-ferment NSP.

In conclusion, the method described for determining the energy values of different non-starch polysaccharide-rich dietary fibre supplements is both reproducible and repeatable. The result is independent of dietary dosage with supplements over the range used so that it is feasible to apply the method using a single dose within this range. Provided that the user of the method is reasonably proficient in animal husbandary and general analytical procedures there is no evidence that the result is dependent on the experience gained by the user of the method. We suggest the method is acceptable as an official method for obtaining energy values of dietary fibre supplements for the purpose of food labelling, tables and databases.

This study was coordinated and supported by the Dietary Fibre Task Force of the International Life Sciences Institute – ILSI Europe, Avenue E. Mounier, 83 – Box 6, B-1200 Brussels. Coordinators were Drs J.-L. Barry (INRA, Nantes, France), T. F. S. (present author), G. L. (present author), and U. Priebe (ILSI Europe). Thanks are due to R. M. Faulks for guidance on the use of the Faulks & Timms (1985) colorimetric method for NSP. The authors are grateful for technical help from Mr F. Montigon, Mrs D. Cavin and Mrs S. Reimann of Nestlé, Mr K. Olsson and Mrs M. Stenberg of Lund University.

REFERENCES

- Bach-Knudsen, K. E., Wisker E., Daniel, F., Feldheim, W. & Eggum, B. O. (1994). Digestibility of energy, protein, fat and non-starch polysaccharides in mixed diets: comparative studies between man and rat. *British Journal of Nutrition* **71**, 471–487.
- Barnett, V. & Lewis, T. (1978). *Outliers in Statistical Data*, p. 99. Chichester: John Wiley & Sons.
- Barry, J.-L., Hoebler, C., MacFarlane, G. T., MacFarlane, S., Mathers, J. C., Reed, K. A., Mortensen, P. B., Norgaard, I., Rowland, I. R. & Rumney, C. J. (1995). Estimation of the fermentability of dietary fibre *in vitro*: a European interlaboratory study. *British Journal of Nutrition* **74**, 303–322.
- Boyer, K. W., Horwitz, W. & Albert, R. (1985). Interlaboratory variance in trace element analysis. *Analytical Chemistry* **57**, 454–459.
- British Nutrition Foundation (1990). *Complex Carbohydrates in Foods: The Report of the British Nutrition Foundation's Task Force*. London: Chapman & Hall.
- Brown, J. C. (1993). Energy: measurement of food energy. In *Encyclopaedia of Food Science, Food Technology and Nutrition*, vol. 13, pp. 1590–1595; [R. Macrae, R. K. Robinson and M. J. Sadler, editors]. London: Academic Press Ltd.
- Brown, J. C. & Livesey, G. (1994). Urinary energy excretion in rats fed guar gum: implications for putative thermogenesis. *Food Hydrocolloids* **8**, 69–72.
- European Council Directive (1990). Nutrition labelling for foodstuffs. *Official Journal of the European Community* **33**, L276.
- Faulks, R. M. & Timms, S. B. (1985). A rapid method for determining the carbohydrate component of dietary fibre. *Food Chemistry* **17**, 273–287.
- Göranzon, H. & Forsum, E. (1987). Metabolisable energy in humans in two diets. Calculation and analysis. *Journal of Nutrition* **117**, 267–273.
- Göranzon, H., Forsum, E. & Thilén, M. (1983). Calculation and determination of the metabolisable energy in mixed diets to humans. *American Journal of Clinical Nutrition* **38**, 954–963.
- Harley, L. J., Davies, I. R. & Livesey, G. (1989). Digestible energy value of gums in the rat – data on gum arabic. *Food Additives and Contaminants* **6**, 13–70.
- Horwitz, W., Kamps, L. R. & Boyer, K. W. (1980). Quality assurance for the analysis of foods for trace constituents. *Journal of the Association of Official Analytical Chemists* **63**, 1344–1354.
- Johnson, I. T., Livesey, G., Gee, J. M. G., Brown, J. C. & Wortley, G. M. (1990). The biological effects and digestible energy value of a sugar-beet fibre preparation in the rat. *British Journal of Nutrition* **64**, 187–199.
- Livesey, G. (1989). Procedure for calculating the digestible and metabolizable energy values of food components making a small contribution to the diet. *Journal of the Science of Food and Agriculture* **48**, 475–481.
- Livesey, G. (1990). Energy values of unavailable carbohydrates and diets: an inquiry and analysis. *American Journal of Clinical Nutrition* **51**, 617–636.

- Livesey, G. (1991). Calculating the energy values of foods: towards empirical formulae based on diets with varied intakes of unavailable complex carbohydrates. *European Journal of Clinical Nutrition* **45**, 1–22.
- Livesey, G. (1992). The energy values of diets and sugar alcohols for man. *Nutrition Research Reviews* **5**, 61–84.
- Miles, C. (1990). The metabolisable energy of diets differing in dietary fat and fibre in humans. *Journal of Nutrition* **122**, 306–311.
- Nyman, M. & Asp, N.-G. (1985). Dietary fibre fermentation in the rat intestinal tract: effect of adaptation period, protein and fibre level, and particle size. *British Journal of Nutrition* **54**, 635–643.
- Nyman, M. & Asp, N.-G. (1988). Fermentation of oat fibre in the rat intestinal tract: A study of different cellular areas. *American Journal of Clinical Nutrition* **48**, 274–279.
- Nyman, M., Asp, N.-G., Cummings, J. H. & Wiggins, H. (1986). Fermentability of dietary fibre in the intestinal tract: comparison between man and rat. *British Journal of Nutrition* **55**, 487–496.
- Prosky, L. (1985). Total dietary fibre in foods. Enzymatic-gravimetric method. *Journal of the Association of Official Analytical Chemists* **68**, 399.
- Prosky, L., Asp, N.-G., Schweizer, T. M., DeVries, J. W. & Furda, I. (1988). Determination of insoluble, soluble, and total dietary fibre in foods and products: interlaboratory study. *Journal of the Association of Official Analytical Chemists* **71**, 1017–1023.
- Roberfroid M., Gibson, G. R. & Delzenne, N. (1993). The biochemistry of oligofructose, a non-digestive fiber: an approach to calculate its caloric value. *Nutrition Research* **51**, 137–148.
- Scott, R. W. (1979). Colorimetric determination of hexuronic acids in plant materials. *Analytical Chemistry* **51**, 936–941.
- Southgate, D. A. T. & Durnin, J. V. G. A. (1970). Calorie conversion factors: an experimental reassessment of the factors used in the calculation of the energy values of human diets. *British Journal of Nutrition* **24**, 517–535.
- Wisker, E. & Feldheim, W. (1992). Faecal bulking and energy value of dietary fibre. In *Dietary Fibre – A Component of Food*, pp. 233–246 [T. F. Schweizer and C. A. Edwards, editors]. London: Springer-Verlag.
- Wisker, E., Maltz, A. & Feldheim, W. (1988). Metabolisable energy of diets low or high in dietary fibre from cereals eaten by humans. *Journal of Nutrition* **118**, 945–952.