



N-acetylcysteine manipulation fails to elicit an increase in glutathione in a teleost model

Birnie-Gauvin, Kim; Larsen, Martin Hage; Aarestrup, Kim; Willmore, William G; Cooke, S.J.

Published in:
Fish Physiology & Biochemistry

Link to article, DOI:
[10.1007/s10695-017-0419-3](https://doi.org/10.1007/s10695-017-0419-3)

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Birnie-Gauvin, K., Larsen, M. H., Aarestrup, K., Willmore, W. G., & Cooke, S. J. (2018). N-acetylcysteine manipulation fails to elicit an increase in glutathione in a teleost model. *Fish Physiology & Biochemistry*, 44(1), 137-142. DOI: 10.1007/s10695-017-0419-3

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3
4
5 **N-acetylcysteine manipulation fails to elicit an increase in glutathione in a teleost model**
6
7

8
9 *In press in Fish Physiology and Biochemistry*
10
11
12
13
14
15
16

17
18 Kim Birnie-Gauvin^{1,2,3†}, Martin H. Larsen^{3,4}, Kim Aarestrup³, William G. Willmore², Steven J.
19 Cooke¹
20
21
22
23
24
25

26 ¹ Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of
27 Environmental Science, Carleton University, 1125 Colonel By Drive, Ottawa, ON, Canada K1S
28 5B6
29

30 ² Department of Biology and Institute of Biochemistry, Carleton University, 1125 Colonel By
31 Drive, Ottawa, ON, Canada K1S 5B6
32

33 ³ DTU AQUA, National Institute of Aquatic Resources, Section for Freshwater Fisheries
34 Ecology, Technical University of Denmark, Vejlsovej 39, 8600 Silkeborg, Denmark
35

36 ⁴ Danish Centre for Wild Salmon, Brusgårdsvej 15, 8960 Randers, Denmark.
37
38
39
40

41 † Author for Correspondence: kbir@aqu.dtu.dk
42
43

44 **Running title:** N-acetylcysteine in brown trout
45
46

47 **Abstract**

48 Levels of oxidative stress can be affected by a range of compounds including toxins and
49 pharmaceuticals. Antioxidants are important protective compounds which counteract the
50 damaging effects of oxidative stress. Glutathione (GSH) is one of the main antioxidants for many
51 organisms, and can be synthesized from administered N-acetylcysteine (NAC). NAC has
52 therefore often been used in a wide range of taxa to manipulate levels of GSH. Our objective was
53 to validate this approach in a wild temperate teleost fish model, the brown trout (*Salmo trutta*).
54 We used intracoelomic injections of NAC in saline and vegetable shortening, at two different
55 concentrations (100 and 400mg/kg), with the appropriate controls and shams, under controlled
56 laboratory settings. We found that NAC failed to elicit an increase in GSH over three time
57 periods and concluded that NAC is not an effective method to enhance GSH levels in teleost fish
58 using the concentrations and vehicles tested here. We emphasize the importance of validation
59 studies across all new species/taxa when possible and suggest that more investigation is required
60 with regards to NAC manipulation in fish if this approach is to be used.

61

62 **Keywords:** glutathione, N-acetylcysteine, teleost fish, saline, validation studies, vegetable
63 shortening

64

65 **Introduction**

66 Antioxidants, and more generally oxidative stress, have received much attention in recent years.
67 The field of oxidative ecology has emerged with the growing body of evidence that oxidative
68 stress processes are linked with life history strategies, coping mechanisms associated with
69 environmental alterations, and pathogenesis (Kehrer 1993; Beaulieu et al. 2013; Speakman et al.

70 2015). Reactive oxygen species (ROS) are continuously generated through mitochondrial
71 respiration (Halliwell and Gutteridge 2015) as well as by the actions of various hormones and
72 neurotransmitters (Finkel 1998). The presence of ROS, if unquenched, can be highly detrimental
73 to cellular macromolecules, and cause oxidative stress (Halliwell and Gutteridge 2015). Of all
74 the forms of ROS, hydrogen peroxide (H₂O₂) is of particular interest as it is one of the most
75 stable and long-lived ROS (Kress et al. 1995).

76 N-acetylcysteine (NAC) is a known thiolic antioxidant, and is a precursor to glutathione
77 synthesis as it provides cysteine groups for γ -glutamylcysteine synthetase, an essential enzyme
78 required for the generation of glutathione (Pena-Llopis et al. 2003; Gutierrez-Praena et al. 2012).
79 NAC also protects against cellular damage through its direct reaction with ROS and cannot be
80 obtained from diet (Aruoma et al. 1989). NAC manipulation has been used in a wide range of
81 taxa (mammals (Reid et al. 1994; Tomkiewicz et al. 1994); amphibians (Giniatullin and
82 Giniatullin 2003); birds (Valdivia et al. 2001)) in an attempt to manipulate glutathione levels. All
83 studies dedicated to NAC manipulation in fish thus far have focused on the investigation of its
84 protective effects against pesticides and pathogens under controlled laboratory conditions (e.g.,
85 Pena-Llopis et al. 2003; Sevgiler et al. 2006; Puerto et al. 2009; Üner et al. 2009; Gutierrez-
86 Praena et al. 2012), but none have documented its potential effects on wild populations.
87 Moreover, to our knowledge, no studies have tested vegetable shortening as a carrier. Vegetable
88 shortening is commonly used when manipulating wild fish from temperate regions (such as the
89 brown trout) because the vehicle will solidify and hence prolong the effects of the injected
90 substance.

91 Given that studies in the wild involving NAC manipulation have yet to be performed, our
92 aim was to determine the best method to administer NAC in a wild population of brown trout

93 under a highly controlled laboratory setting in an attempt to induce an increase in glutathione.
94 Our goal was to bring a more experimental approach to a body of literature that is dominated by
95 correlations (i.e., this approach would provide us with a way of manipulating antioxidants in
96 wild fish, which may help us understand the role of oxidative stress processes in an ecological
97 context). We tested saline and vegetable shortening as vehicles for intracoelomic NAC
98 injections. We hypothesized that saline injections containing NAC would be more readily
99 absorbed given that NAC is highly soluble in water, and therefore predicted that glutathione
100 (GSH) would become elevated more quickly than in the vegetable shortening-treated fish.
101 Furthermore, we predicted that the increase in GSH using saline would be short-lived, given that
102 NAC will be absorbed more rapidly than with the vegetable shortening. We also hypothesized
103 that vegetable shortening injections containing NAC would take longer to be absorbed given that
104 vegetable shortening solidifies after the injection, and thus predicted that elevated GSH may take
105 longer to appear, but that its presence will be long-lasting in comparison to saline-injected fish.

106

107 **Material and Methods**

108 On July 1st, 2016, wild juvenile brown trout (n = 240) were captured from the Kastbjerg stream,
109 Jutland, Denmark, using backpack electrofishing (Scubla ELT 60 II GI; 300 volts). Fish were
110 transported to the laboratory facilities in a 100L tank of fresh oxygenated stream water, and were
111 randomly attributed to one of three identical 4000L tanks (n = 80 per tank). The tanks had a
112 constant circulating flow of fresh oxygenated water, held at a constant temperature of
113 $13.5 \pm 0.4^\circ\text{C}$ (average temperature in the wild typically fluctuates between 10 and 15°C during the
114 summer). All fish were kept at a 17:7 light:dark photoperiod (representative of daylight in

115 Denmark during the summer months), and fed daily with mosquito larvae, starting one day after
116 the manipulation.

117 Fish were left to acclimate for 24 hours, prior to manipulation. Fish were anesthetized
118 using a solution of benzocaine (0.03g l^{-1} ethyl-*p*-aminobenzoate; Sigma) in water, then weighed
119 ($\pm 0.01\text{g}$), measured for total length ($\pm 0.1\text{cm}$), and tagged using a 12mm PIT tag (Texas
120 Instruments, RI-TRP-RRHP, 134Hz, 0.01g mass in air, Plano, Texas, USA). Fish were randomly
121 assigned to one of seven treatment groups: (1) control, (2) sham-saline, (3) sham-shortening, (4)
122 100mg/kg NAC in saline (sal-low), (5) 100mg/kg NAC in vegetable shortening (veg-low), (6)
123 400mg/kg NAC in saline (sal-high), and (7) 400mg/kg NAC in vegetable shortening (veg-high),
124 each group containing 30 fish (10 fish from each tank). In addition, some fish were simply left in
125 the tank (i.e., totally undisturbed), and remained untouched until sampling (i.e., not tagged,
126 weighed or measured) so as to detect tagging effects if necessary, despite evidence that tagging
127 has minimal impacts on salmonids (Larsen et al. 2013). Control fish were recovered in a 60L
128 tank of fresh water following tagging. NAC-treated fish received an intracoelomic injection of a
129 suspension of physiological saline (0.59% NaCl in pure water) or vegetable shortening (100%
130 vegetable shortening, Crisco, OH, USA) mixed with N-acetylcysteine (NAC; Sigma-Aldrich, St.
131 Louis, MO, USA, Product A7250) using a dosage of 0.01 mL vehicle (concentration of 0.01g or
132 0.04 NAC per mL) per 1 g of fish (equivalent to 100 or 400 mg kg⁻¹, respectively). Sham fish
133 were injected with only 0.01mL g⁻¹ saline or vegetable shortening. NAC-treated fish were
134 recovered separately from control and sham fish to prevent cross-treatment contamination of
135 NAC. Once recovered, all fish were returned to the tank.

136 After 3 days, all fish from tank 1 (10 fish from each treatment group) were anesthetized
137 and weighed as per the above description. Fish were sampled for blood (0.1ml) from the caudal

138 vasculature using a 25-gauge heparinized needle. Fish were then immediately euthanized using a
139 lethal percussion. All samples were immediately flash-frozen with liquid nitrogen, and then
140 stored at -80°C until analyzed. The same sampling technique was used at 6 and 9 days post-
141 treatment using fish from tank 2 and 3, respectively. This method was used to avoid disturbing
142 fish until sampling. These standardized techniques were approved by the Danish Animal
143 Experiments Inspectorate (License Number: 2013-15-2934-00808).

144 Glutathione (GSH) was measured in red blood cells (RBCs) samples using a glutathione
145 assay as described in Birnie-Gauvin et al. (2017). This assay measures total glutathione (TGS_H)
146 and oxidized glutathione (GSSG). The concentration of reduced glutathione (GSH), the
147 antioxidant, can then be derived from these values. Final values of GSH were reported in μM.

148 Statistical analyses were conducted using JMP v12.0.1 (SAS Institute Inc.,
149 Buckinghamshire, UK). A two-way ANOVA followed by a Tukey *post hoc* was used to evaluate
150 differences in mass changes among the seven treatment groups, as well as differences in
151 glutathione concentration.

152

153 **Results**

154 Fish initially weighed between 22.8 and 28.0g. Fish in each treatment and day did not
155 differ in condition initially ($F_{6,179} = 0.94$, $p = 0.47$). An interaction between treatment and day
156 was detected for change in mass ($F_{12,179} = 2.12$, $p = 0.0178$, Table 1) such that sal-high fish
157 gained the most mass on day 3. Generally, fish progressively decreased in mass over the course
158 of the study. Time had a significant effect on glutathione concentration, such that day 6 had
159 significantly elevated glutathione in comparison to days 3 and 9 ($F_{2,206} = 20.01$, $p < 0.0001$).
160 This elevation was observed in all groups including the control fish and the shams. On day 5 of

161 the study, 6 and 3 fish, from tanks 2 and 3 respectively, were found dead. These fish all belonged
162 to the sal-high treatment group. Additionally, one fish from the veg-low group was found dead in
163 tank 3 as a result of jumping out of the tank. No other mortality occurred and at sampling all fish
164 were vigorous.

165

166 **Discussion**

167 The objective of our study was to validate the use of N-acetylcysteine (NAC) as a method
168 to increase glutathione (GSH) in a teleost model; the brown trout. Based on the literature we
169 anticipated that we would see an increase in GSH with NAC injections. However, our results fail
170 to demonstrate that NAC elicited such a response, given that no differences were observed
171 among treatments. These findings pose a set of important concerns with regards to the use of
172 NAC in teleost fish.

173 While a number of studies have claimed the protective effects of NAC against oxidative
174 stress processes via an increase in glutathione synthesis (Peña-Llopis et al. 2003; Gutiérrez-
175 Praena et al. 2012), none that we know of have properly validated this in fish. A common caveat
176 to these types of experimental studies is in fact the lack of appropriate controls, shams, or
177 validations (Cooke et al. in press). In the majority of studies currently in the literature, either
178 shams or controls are missing, making it rather difficult to interpret results and draw conclusions.
179 Many of the studies investigating the protective effects of NAC have been performed on human
180 patients with various illnesses (e.g., Horowitz et al. 1988; Prescott et al. 1989) or other
181 mammalian models such as the rat (e.g., Moussawi et al. 2009). It is therefore possible that the
182 physiological mechanisms by which NAC acts in mammals differ from those in fish.
183 Alternatively, it is possible that the effects of NAC take longer to appear in tissues in wild brown

184 trout given that antioxidant capacity and glutathione concentrations are already high (Birnie-
185 Gauvin et al. 2017). No differences were observed in the control, shams and untagged fish, when
186 compared to treated fish, suggesting that NAC injections had no effect on GSH at all.
187 Additionally, the increase in GSH observed in all groups at day 6 is likely not the result of the
188 NAC injections themselves, given that control, shams and untagged fish showed the same
189 increase. It is also highly unlikely that laboratory conditions caused the observed day 6 increase,
190 given that temperature, lighting, flow and overall fish conditions were monitored at least 4 times
191 a day.

192 Though NAC injections failed to increase GSH, a number of fish from the sal-high group
193 were found dead on day 6 (from both the day 6 and day 9 tanks), suggesting that high
194 concentrations of NAC absorbed at a rapid rate may have lethal impacts in fish. Similar results
195 were found in a study on rats where low doses of NAC had protective effects against
196 lipopolysaccharide toxicity, but high doses had the opposite effect and even increased mortality
197 (Sprong et al. 1998). While it is possible that keeping wild fish in captivity has caused unknown
198 physiological alterations where glutathione synthesis could be affected, previous studies have
199 demonstrated that after 24h in captivity, wild salmonids are typically calm with normal baseline
200 levels of cortisol in comparison to captive-bred counterparts (e.g., Lepage et al. 2000; Patterson
201 et al. 2004; Portz et al. 2006). Alternatively, GSH may have increased prior to the first sampling
202 period at 3 days. Peña-Llopis et al. (2003) detected an increase in GSH as early 12 hours post-
203 injection with saline, though only sham fish were used in this study (no controls), making it
204 difficult to properly interpret the results. Another alternative hypothesis as to why GSH increased
205 in all treatments on day 6 is coincidental fluctuations in normal GSH levels. Further investigation

206 is required to better understand natural patterns of GSH as well as the mechanistic basis for NAC
207 in fish.

208 Validation studies, such as the present one, are crucial components of proper
209 experimental science. We therefore urge other groups to take a similar approach to test the
210 fundamental concepts applied to their study, and for each new species, when possible. We
211 conclude that further studies are required to investigate whether NAC injection is an adequate
212 method to manipulate glutathione levels in teleost fish. We acknowledge that other vehicles or
213 concentrations could have yielded different findings, but the ones used here are common carriers
214 for other taxa. It may be worthwhile to explore other manipulation methods such of those that
215 involve dietary manipulation (e.g., NAC infused in food items) or use of mini-osmotic pumps.
216 Clearly, additional detailed validation work is needed before NAC is used to manipulate
217 oxidative status in wild fish. Given the interest in bringing a more experimental approach to
218 oxidative ecology, such validations are pressing. Until then, we caution against using NAC to
219 manipulate oxidative status in teleosts.

220

221 **Acknowledgments**

222 W.G.W. and S.J.C. are supported by NSERC Discovery Grants. S.J.C. is supported by the
223 Canada Research Chairs Program. We would like to thank Andreas Svarer for his help in the
224 field, as well as Anders Koed and Henrik Baktoft for their help in setting up the laboratory
225 experiments.

226

227

228

229

230

231 **References**

- 232
- 233 Aruoma, O.I., Halliwell, B., Hoey, B.M., Butler, J., 1989. The antioxidant action of N-
- 234 acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide, and
- 235 hypochlorous acid. *Free. Rad. Biol. Med.* 6, 593-597.
- 236
- 237 Beaulieu, M., Thierry, A.M., González-Acuña, D., Polito, M.J., 2013. Integrating oxidative
- 238 ecology into conservation ecology. *Cons. Physiol.* 1, cot004.
- 239
- 240 Birnie-Gauvin, K., Peiman, K.S., Larsen, M.H., Baktoft, H., Aarestrup, K., Willmore, W.G.,
- 241 Cooke, S.J., 2017. Short-term and long-term effects of exogenous cortisol manipulation on
- 242 oxidative stress in juvenile brown trout. *Journal of Experimental Biology*, 220, 1693-1700.
- 243
- 244 Cooke, S.J., Birnie-Gauvin, K., Lennox, R.J., Taylor, J.J., Rytwinski, T., Rummer, J.L., Franklin,
- 245 C.E., Bennett, J.R., Haddaway, N.R., in press. How experimental biology and ecology can
- 246 support evidence-based decision-making in conservation: avoiding pitfalls and enabling
- 247 application. *Cons. Physiol.*
- 248
- 249 Finkel, T., 1998. Oxygen radicals and signaling. *Curr. Opinion Cell. Biol.* 10, 248-253.
- 250
- 251 Giniatullin, A.R., Giniatullin, R.A., 2003. Dual action of hydrogen peroxide on synaptic
- 252 transmission at the frog neuromuscular junction. *J. Physiol.* 552, 283-293.
- 253
- 254 Gutiérrez-Praena, D., Puerto, M., Prieto, A.I., Jos, Á., Pichardo, S., Vasconcelos, V., Cameán,
- 255 A.M., 2012. Protective role of dietary N-acetylcysteine on the oxidative stress induced by
- 256 cylindrospermopsin in tilapia (*Oreochromis niloticus*). *Environ. Toxicol. Chem.* 31, 1548-1555.
- 257
- 258 Halliwell, B., Gutteridge, J.M.C., 2015. Free radicals in biology and medicine. Oxford
- 259 University Press, USA.
- 260
- 261 Horowitz, J.D., Henry, C.A., Syrjanen, M.L., Louis, W.J., Fish, R.D., Antman, E.M., Smith,
- 262 T.W., 1988. Combined use of nitroglycerine and N-acetylcysteine in the management of unstable
- 263 angina pectoris. *European Heart J.* 9, 95-100.
- 264
- 265 Kehrer, J., 1993. Free radicals as mediators of tissue injury and disease. *Critical Rev. Toxicol.*
- 266 23, 21-48.
- 267
- 268 Kress, M., Riedl, B., Reeh, P.W., 1995. Effect of oxygen radicals on nociceptive afferents in the
- 269 rat skin in vitro. *Pain* 62, 87-94.
- 270
- 271 Larsen, M.H., Thorn, A.N., Skov, C., Aarestrup, K., 2013. Effects of passive integrated
- 272 transponder tags on survival and growth of juvenile Atlantic salmon *Salmo salar*. *Anim.*
- 273 *Biotelem.* 1, 19-26.
- 274
- 275 Lepage, O., Øverli, Ø., Petersson, E., Järvi, T., Winberg, S., 2000. Differential stress coping in
- 276 wild and domesticated sea trout. *Brain Behav. Evol.* 56, 259-268.

277
278 Moussawi, K., Pacchioni, A., Moran, M., Foster Olive, M., Gass, J.T., Lavin, A., Kalivas, P.W.,
279 2009. N-acetylcysteine reverses cocaine-induced mataplasticity. *Nature Neurosci.* 12, 182-189.
280
281 Patterson, A.D., Macdonald, J.S., Hinch, S.G., Healey, M.C., Farrell, A.P., 2004. The effect of
282 exercise and captivity on energy partitioning, reproductive maturation and fertilization success in
283 adult sockeye salmon. *J. Fish Biol.* 64, 1039-1059.
284
285 Pena-Llopis, S., Ferrando, M.D., Peña, J.B., 2003. Fish tolerance to organophosphate-induced
286 oxidative stress is dependent on the glutathione metabolism and enhanced N-acetylcysteine.
287 *Aquatic Toxicol.* 65, 337-360.
288
289 Portz, D.E., Woodley, C.M., Cech, J.J., 2006. Stress-associated impacts of short-term holding on
290 fishes. *Rev. Fish Biol. Fish.* 16, 125-170.
291
292 Prescott, L.F., Donovan, J.W., Jarvie, D.R., Proudfoot, A.T., 1989. The disposition and kinetics
293 of intravenous N-acetylcysteine in patients with paracetamol overdose. *European J. Clin.*
294 *Pharmacol.* 37, 501-506.
295
296 Puerto, M., Prieto, A.I., Pichardo, S., Moreno, I., Jos, Á., Moyano, R., Cameán, A.M., 2009.
297 Effects of dietary N-acetylcysteine on the oxidative stress induced in tilapia (*Oreochromis*
298 *niloticus*) exposed to a microcystin-producing cyanobacterial water bloom. *Environ. Toxicol.*
299 *Chem.* 28, 1679-1686.
300
301 Reid, M.B., Stokić, D.S., Koch, S.M., Khawli, F.A., Leis, A.A., 1994. N-acetylcysteine inhibits
302 muscle fatigue in humans. *J Clin. Investigat.* 94, 2468-2474.
303
304 Sevgiler, Y., Piner, P., Durman, H., Üner, N., 2007. Effects of acetylcysteine on oxidative
305 responses in the liver of fenthion exposed *Cyprinus carpio*. *Pesticide Biochem. Physiol.* 87, 248-
306 254.
307
308 Speakman, J.R., Blount, J.D., Bronikowski, A.M., Biffenstein, R., Isaksson, C., Kirkwood,
309 T.B.L., Monaghan, P., Ozanne, S.E., Beaulieu, M., Briga, M., Carr, S.K., Christensen, L.L.,
310 Cocheme, H.M., Cram, D.L., Dantzer, B., Harper, J.M., Jurk, D., King, A., Noguera, J.C., Salin,
311 K., Sild, E., Simons, M.J.P., Smith, S., Stier, A., Tobler, M., Vitikainen, E., Peaker, M., Selman,
312 C., 2015. Oxidative stress and life histories: unresolved issues and current needs. *Ecol. Evol.* 5,
313 5745-5757.
314
315 Sprong, R.C., Winkelhuyzen-Janssen, A.M.L., Aarsman, C.J.M., van Oirschot, F.M.L.M., van
316 der Bruggen, T., van Asbeck, B.S., 1998. Low-dose N-acetylcysteine protects rats against
317 endotoxin-mediated oxidative stress, but high-dose increases mortality. *Am. J. Resp. Critical*
318 *Care Med.* 157,1283-1293.
319
320 Tomkiewicz, R.P., App, E.M., Coffiner, M., Fossion, J., Maes, P., King, M., 1994. Mucolytic
321 treatment with N-acetylcysteine L-lysinate metered dose inhaler in dogs: airway epithelial
322 function changes. *European Resp. J.* 7, 81-87.

- 323
324 Üner, N., Sevgiler, Y., Durmaz, H., Piner, P., Çinkiloglu, E., 2009. N-acetylcysteine provides
325 dose-dependent protection against fenthion toxicity in the brain of *Cyprinus carpio* L. *Comp.*
326 *Biochem. Physiol. C* 150, 33-38.
327
328 Valvidia, A.G., Martinez, A., Damian, F.J., Quezada, T., Ortiz, R., Martinez, C., Llamas, J.,
329 Rodriguez, M.L., Yamamoto, L., Jaramillo, F., Loarca-Piña, M.J., Reyes, J.L., 2001. Efficacy of
330 N-acetylcysteine to reduce the effects of aflatoxin B₁ intoxication in broiler chickens. *Poultry*
331 *Sci.* 80, 727-734.


332 **Table 1. Change in mass of treated brown trout.** Average change in mass (g) for each
 333 treatment groups, across sampling days (\pm SEM). Sample sizes are shown in parentheses.
 334 Treatment (and day had significant effects on changes in mass (Tukey *post-hoc*, $p < 0.001$).
 335 Asterisk represents significant difference from control of the same day.

336
 337
 338

Treatment	Sampling day		
	3	6	9
Control	-0.38 \pm 0.18 (10)	-0.92 \pm 0.20 (10)	-0.83 \pm 0.21 (10)
Veg	0.16 \pm 0.14 (10)	-0.25 \pm 0.22 (10)	-0.89 \pm 0.19 (10)
Veg-low	0.22 \pm 0.29 (10)	-0.36 \pm 0.17 (10)	-1.13 \pm 0.26 (9)
Veg-high	0.18 \pm 0.20 (10)	-0.55 \pm 0.27 (10)	-1.02 \pm 0.30 (10)
Sal	-0.28 \pm 0.21 (10)	-0.97 \pm 0.23 (10)	-1.56 \pm 0.27 (10)
Sal-low	-0.33 \pm 0.17 (10)	-1.04 \pm 0.25 (10)	-1.16 \pm 0.17 (10)
Sal-high	1.54 \pm 0.23 (10)*	0.10 \pm 0.21 (4)	-1.01 \pm 0.28 (7)

339

340 **Figure 1. Levels of glutathione in treated brown trout.** Glutathione concentration (μM) \pm
341 SEM, across treatments and days. Time had a significant effect at day 6 on all treatments (Tukey
342 *post-hoc*, $p < 0.0001$).
343
344



345