

Technical University of Denmark



## The impact of a diet with fructan-rich chicory roots on Oesophagostomum dentatum worm population dynamics and host immune responses in pigs

Thamsborg, Stig Millan; Mejer, Helena; Skovgaard, Kerstin; Sengupta, Mita E.; Kringel, Helene; Petersen, Heidi Huus; Jensen, Bent Borg; Andreasen, Annette

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Thamsborg, S. M., Mejer, H., Skovgaard, K., Sengupta, M. E., Kringel, H., Petersen, H. H., ... Andreasen, A. (2017). The impact of a diet with fructan-rich chicory roots on Oesophagostomum dentatum worm population dynamics and host immune responses in pigs. Abstract from 26th International Conference of World Association for the Advancement of Veterinary Parasitology (WAAVP), Kuala Lumpur, Malaysia.

**DTU Library**  
Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## 26<sup>th</sup> International Conference of the World Association for Advancement of Veterinary Parasitology

Theme: **“Combating Zoonoses: Strength in East-West Partnerships”**

Time: Monday 4 September – Friday 8 September

Venue: Kuala Lumpur Convention Centre, Malaysia

---

### **The impact of a diet with fructan-rich chicory roots on *Oesophagostomum dentatum* worm population dynamics and host immune responses in pigs**

Stig Milan Thamsborg<sup>\*1</sup> ; Helena Mejer<sup>1</sup> ; Kerstin Skovgaard<sup>1</sup> ; Mita E. Sengupta<sup>1</sup> ; Helene Kringel<sup>1</sup> ; Heidi H. Petersen<sup>1</sup> ; Bent Borg Jensen<sup>2</sup> ; Annette Andreasen<sup>1</sup>

<sup>1</sup> *Veterinary and Animal Sciences / University of Copenhagen / Denmark*, <sup>1</sup> *Veterinary Institute / Technical University of Denmark / Denmark*, <sup>2</sup> *Animal Science/ Aarhus University / Denmark*

*Oesophagostomum* infections in pigs persist for months. We hypothesized that feeding fructans (dried chicory roots) may improve immunity and facilitate worm expulsion. We therefore examined the effects of long-term chicory on *O. dentatum* population dynamics and host immune responses. Methods: Seventy-two pigs were allocated to four groups in a 2-factorial design. Group O was fed regular feed and trickle inoculated with 15 *O. dentatum* L3/kg/day 0-12 weeks post-infection (pi.) start. Group OC was also trickle inoculated but switched to a chicory-rich diet (12% inulin in DM) weeks 3-12 pi. Group C was uninfected but switched to chicory diet while Group Ctr remained uninfected on regular feed. Six pigs per group were necropsied 5, 9 and 12 weeks pi. for worm counts and qRT-PCR for gene expression in the gut. Faecal egg counts (FEC) and specific antibody levels were assessed regularly. Results: When group OC switched to chicory diet, FECs dropped within 3-4 days and remained very low. Worm counts were reduced 50-65% by chicory feeding (Group OC versus O;  $p < 0.001$ ) and was accompanied by a 2-fold higher *O. dentatum*-specific IgG1 response. In group O, a build-up of a typical Th2-type immune response was seen but leveled out later and worm counts remained stable. Group C had a down-regulated Th1-type response and thus an anti-inflammatory effect in colon. Conclusions: We found little evidence that chicory feeding improved host protective immunity against *Oesophagostomum*. It seems more likely, as previously suggested, that physico-chemical changes in caeco-colon are responsible for the observed anthelmintic effects.

*Keywords: swine; oesophagostomum; chicory; prebiotics; immune response*