Technical University of Denmark



### Survival and infectivity of chicken ascarid eggs in soil after exposure to an eggdegrading microfungus

Mejer, Helena; Thamsborg, Stig M.; Meyling, Nicolai V.; Rui, Wang; Dalgaard, Tina S.; Petersen, Heidi Huus; Thapa, Sundar

Publication date: 2017

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Mejer, H., Thámsborg, S. M., Meyling, N. V., Rui, W., Dalgaard, T. S., Petersen, H. H., & Thapa, S. (2017). Survival and infectivity of chicken ascarid eggs in soil after exposure to an egg-degrading microfungus. Abstract from 26th International Conference of World Association for the Advancement of Veterinary Parasitology (WAAVP), Kuala Lumpur, Malaysia.

## DTU Library Technical Information Center of Denmark

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# 26<sup>th</sup> International Conference of the World Association for Advancement of Veterinary Parasitology

Theme: "Combating Zoonoses: Strength in East-West Partnerships"

Time: Monday 4 September – Friday 8 September

Venue: Kuala Lumpur Convention Centre, Malaysia

# Survival and infectivity of chicken ascarid eggs in soil after exposure to an egg-degrading microfungus

Helena Mejer<sup>\*1</sup>; Stig M. Thamsborg<sup>1</sup>; Nicolai V. Meyling<sup>1</sup>; Wang Rui<sup>2</sup>; Tina S. Dalgaard<sup>3</sup>; Heidi H. Petersen<sup>4</sup>; Sundar Thapa<sup>1</sup>

<sup>1</sup>Department of Veterinary and Animal Sciences/ University of Copenhagen/ Denmark, 1Department of Plant and Environmental Sciences/ University of Copenhagen/ Denmark, <sup>2</sup>College of Veterinary Medicine/ Inner Mongolia Agricultural University/ China, <sup>3</sup>Department of Animal Science/ Aarhus University/ Denmark, <sup>4</sup>National Veterinary Institute/ Technical University of Denmark/ Denmark.

The microfungus *Pochonia chlamydosporia* has been shown to kill high numbers of chicken ascarid (*Ascaridia galli* and *Heterakis* spp.) eggs *in vitro* but it is not known if surviving eggs may be infective. Unembryonated ascarid eggs (predominantly *A. galli*) were therefore isolated from faeces and added to sterilised (S) or non-sterilised (N) soil in Petri dishes that were either treated with *P. chlamydosporia* (F) or left untreated (C) during incubation at 22°C for 35 days. Egg recovery was estimated before (day 0) and after (day 35) treatment. Thereafter, each of four groups of parasite-free egg-laying hens was exposed to the soil from one of the four treatments in the feed over 12 days. The hens were necropsied day 42 post first exposure. The number of surviving eggs was most substantially reduced in SF soil and SF hens had statistically lower worm burdens (both parasites) compared to SC, NC and NF hens. However, adult *A. galli* were primarily found in SF hens while the other groups mainly harboured immature *A. galli*. Accordingly, SF hens also had the highest ascarid faecal egg counts and lowest serum *A. galli* IgY titre. Overall, *A. galli* recovery increased with increasing exposure, but contrastingly resulted in a reduced risk of egg-producing adult worms, at least short-term. Eggs not destroyed by *P. chlamydosporia* were clearly infective to the hens. The fungus did not appear to be sufficiently effective as a biocontrol agent of ascarid eggs in natural (i.e. non-sterilised) soil unless its effect can be further optimised.

Keywords: Ascaridia galli; Pochonia chlamydosporia; hens; soil