



## Short-term effect of oral amoxicillin treatment on the gut microbial community composition in farm mink (Neovison vison)

**Marker, Laura Meier; Hammer , Anne Sofie ; Andresen, Lars; Isaack, Pernille; Clausen, Tove; Byskov, Kevin; Honoré, Oliver Lykke; Jensen, Søren Krogh; Bahl, Martin Iain**

*Published in:*  
Fems Microbiology Ecology

*Link to article, DOI:*  
[10.1093/femsec/fix092](https://doi.org/10.1093/femsec/fix092)

*Publication date:*  
2017

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Marker, L. M., Hammer , A. S., Andresen, L., Isaack, P., Clausen, T., Byskov, K., ... Bahl, M. I. (2017). Short-term effect of oral amoxicillin treatment on the gut microbial community composition in farm mink (Neovison vison). *Fems Microbiology Ecology*, 93(7), [fix092]. DOI: 10.1093/femsec/fix092

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Short-term effect of oral Amoxicillin treatment on the Gut Microbial Community**  
2 **Composition in Farm Mink (*Neovison vison*)**

3

4 Laura Meier Marker<sup>1</sup>, Anne Sofie Hammer<sup>1</sup>, Lars Andresen<sup>1</sup>, Pernille Isaack<sup>1</sup>, Tove Clausen<sup>2</sup>,  
5 Kevin Byskov<sup>2</sup>, Oliver Lykke Honoré<sup>1</sup>, Søren Krogh Jensen<sup>3</sup>, Martin Iain Bahl<sup>4\*</sup>

6

7 <sup>1</sup>Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences,  
8 University of Copenhagen, Frederiksberg C, Denmark

9

10 <sup>2</sup>Danish Fur Breeders Research Centre, Holstebro, Denmark

11

12 <sup>3</sup>Department of Animal Science, Aarhus University, Foulum, Denmark

13

14 <sup>4</sup>National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark

15

16

17 **\*Corresponding author:** National Food Institute, Technical University of Denmark, Kemitorvet  
18 Building 202, DK-2800 Kgs. Lyngby, Denmark. Tel: +45 35887036; E-mail: mbah@food.dtu.dk

19

20 **One Sentence abstract:** Amoxicillin affects the intestinal mucosa-associated bacterial community  
21 in mink despite the very fast gastro-intestinal transit time of these carnivorous animals.

22

23 **Keywords:** mink, *Neovison vison*, carnivore, gut microbiota, amoxicillin, 16S rRNA gene  
24 sequencing

## 25 ABSTRACT

26 It is well documented that antibiotics have pronounced modulatory effects on the intestinal bacterial  
27 community of both humans and animals, with potential health consequences. The gut microbiota of  
28 mink has however attracted little attention due to low bacterial load and fast gastrointestinal transit  
29 time, questioning its relevance. In the present study we hypothesise that oral amoxicillin treatment  
30 affects the gut microbiota in mink. This was investigated in a controlled trial including 24 animals  
31 of which 12 were treated with amoxicillin for seven days. By applying 16S rRNA gene sequencing  
32 we found that the faecal microbiota was markedly altered already after two days of treatment, with  
33 a surprising increase in diversity to resemble the feed. The diversity within the mucosa at  
34 termination was however reduced, which indicates this compartment as an important colonization  
35 site in mink. No impact on blood biochemistry, lipid metabolism, serum amyloid A, vitamins A and  
36 E and histomorphology of the gut and liver was found, however, a slight decrease in fat digestibility  
37 was observed. We suggest that early life use of amoxicillin in mink production may be  
38 counteractive as dysbiosis of the microbiota during infancy is increasingly being recognized as a  
39 risk factor for future health.

40

## 41 INTRODUCTION

42 The rich microbial community residing in the intestinal tract of humans and animals is collectively  
43 termed the gut microbiota. There is a growing body of evidence manifesting that the function of this  
44 gut microbiota is not restricted to general digestion of nutrients in the intestine but also plays a  
45 tremendously important role in relation to general health including programming of the immune  
46 system especially in early life (Matamoros *et al.* 2013; Albenberg and Wu 2014; Round and  
47 Mazmanian 2014). In addition, the gut microbiota has the capacity to synthesize vitamins B and K  
48 (LeBlanc *et al.* 2013), deconjugate bile acids (Jones *et al.* 2008) and produce short chain fatty acids

49 and antimicrobial compounds, which overall may limit pathogen colonization and proliferation  
50 (Kamada *et al.* 2013). Perturbation of the intrinsic balance between the hundreds of different  
51 species within this ecosystem is especially profound following exposure to antimicrobial  
52 compounds and it is well established that oral antibiotic treatment may cause large shifts in  
53 intestinal bacterial community composition in both humans (Panda *et al.* 2014), rodents (Tulstrup *et*  
54 *al.* 2015) and numerous other animals representing both vertebrate- (Grønvold *et al.* 2010) and  
55 invertebrate species (Raymann, Shaffer and Moran 2016). The effect of antibiotic treatment on the  
56 gut microbiota is highly dependent on the class and spectrum of the drug in question (Tulstrup *et al.*  
57 2015) and different studies have demonstrated both short- and long-lasting effects of antibiotics not  
58 only on pathogens but also on the resident gut microbiota (Cochetière *et al.* 2005; Jernberg *et al.*  
59 2007; Dethlefsen *et al.* 2008). The extent of the disturbances additionally depends on the dose and  
60 the route of administration as well as the general pharmacokinetic and pharmacodynamic properties  
61 of the drug (Edlund and Nord 2000; Sullivan, Edlund and Nord 2001).

62 To date, very few studies have investigated the effects of antibiotics on the gut microbiota in  
63 carnivorous animals (Suchodolski *et al.* 2009; Grønvold *et al.* 2010) and to our knowledge, no  
64 studies have specifically investigated the impact of antibiotics on farm mink (*Neovison vison*).  
65 Recently, high throughput techniques using 16S rRNA gene sequencing of mucosa associated  
66 intestinal samples obtained from mink has showed that the gut microbiota in adult mink is  
67 commonly dominated by bacteria belonging to the phylum *Firmicutes* (especially the class  
68 *Clostridia*) with few *Bacteroidetes*, but may also contain high numbers of both *Proteobacteria* and  
69 *Fusobacteria* (Bahl *et al.* 2017). The dominance of *Firmicutes* is in accordance with similar studies  
70 sequencing faecal samples from cats and dogs (Handl *et al.* 2011; Garcia-Mazcorro *et al.* 2012),  
71 indicating that mink may be a useful animal model for studying other carnivorous animals as well.  
72 Mink have a relatively short intestinal tract and a fast gastro-intestinal transit time of 3-5 hours

73 (Bleavins and Aulerich 1981; Szymeczko and Skrede 1990). A bacterial density up to  $10^8$  CFU g<sup>-1</sup>  
74 has been detected in the colon of mink, which is much lower than numbers of bacteria found in  
75 many other mammals including dogs (Davis *et al.* 1977; Williams, Buddington and Elnif 1998).  
76 Collectively these characteristics could suggest a relatively restricted effect of antibiotics  
77 administered orally on the gut microbiota.

78 In Danish mink production, the largest amount of antibiotics is used during the lactation period  
79 in May until August (Chriél *et al.* 2012; Jensen *et al.* 2016). At this stage mink kits have a thinner  
80 mucus layer lining the gut wall, different enzyme-related properties in the gastrointestinal tract  
81 (Hedemann, Clausen and Jensen 2011), and a different microbial community composition  
82 compared to adult mink (Williams, Buddington and Elnif 1998) potentially making them more  
83 sensitive than adult animals. The broad-spectrum antibiotic, amoxicillin, is the most frequently used  
84 antibiotic in mink and is utilized for treatment of various infectious diseases including diarrhoea,  
85 urinary tract infections, pneumonia, pleuritis, and abscesses (Pedersen *et al.* 2009; Chriél *et al.*  
86 2012). Reports of antimicrobial use in mink in recent years reveal that the oral administration route  
87 constitutes 98% of the antimicrobial use measured in Defined Animal Daily Dose (Jensen *et al.*  
88 2016). Since oral administration of antimicrobials for mink may involve treatment of either part of  
89 or the entire farm, including both sick and asymptomatic animals, there is a need to understand the  
90 potential impact of the oral antimicrobial administration on the microbiota of the healthy mink gut.

91 The present study was designed to investigate the induced changes in the bacterial community  
92 composition of the gut microbiota as well as changes related to biochemical profile, vitamin A and  
93 E levels, lipid metabolism, levels of acute phase reactant serum amyloid A, and histomorphological  
94 parameters in the gut and liver following a standard treatment period of 7 days with daily oral  
95 amoxicillin administered to clinically healthy growing male mink.

96

## 97 MATERIALS AND METHODS

### 98 **Ethical statement**

99 All institutional and national guidelines for the care and use of laboratory animals were followed.

100 The handling of the animals and the experimental procedures were approved by the Danish Animal  
101 Experiments Inspectorate (licence no. 2016-15-0201-00965) and all personnel involved in handling  
102 and care of the animals were trained for carrying out animal experiments (FELASA certification).

103

### 104 **Animals and housing**

105 The study was carried out at the Danish Fur Breeders Association research farm, Kopenhagen Fur,  
106 Holstebro, Denmark from the 12<sup>th</sup> to the 26<sup>th</sup> of September 2016. Twenty-four male mink (*Neovison*  
107 *vison*) at 4.5 months of age of the colour type Brown (also referred to as colour type Wild) were  
108 selected for the study. The mink were all littermates in pairs from 12 different litters with a  
109 minimum of 5 kits. The average body mass was  $2,809 \pm 300$  g (mean  $\pm$  SD) with 584 g as the  
110 maximum body mass variance between siblings. All animals had received standard vaccinations  
111 (Distemink<sup>®</sup> Vet., Biovet ApS and Biocom-P Vet., Biovet ApS) at the age of 10 weeks and the farm  
112 was tested negative for Aleutian Disease Virus. The mink were housed indoors individually in  
113 metabolic cages modified from Jørgensen, G. & Glem Hansen (1973) with the possibility for  
114 manual ventilation. The average ambient temperature and humidity was  $21 \pm 1.8^{\circ}\text{C}$  and  $68.3 \pm$   
115  $4.2\%$ , respectively. All mink were fed 180 g of a diet produced at the local central of feedstuff once  
116 a day in the morning and had free access to tap water. The same batch of feed was used throughout  
117 the entire study.

118

119 **Experimental design**

120 The study was designed as a parallel group experiment. Mink were allocated in two groups by  
121 randomized stratification with one littermate in each group (Fig. 1). Amoxicillin (Octacillin Vet.  
122 800 mg g<sup>-1</sup>, amoxicillin trihydrate corresponding to 697 mg amoxicillin), 20 mg kg<sup>-1</sup> was  
123 administered orally to the treatment group (AMX) from Day 0 and for 7 consecutive days while no  
124 treatment was given to the control group (CON). The antimicrobial compound was mixed in the  
125 feed ration. General health checks were made in the morning and afternoon throughout the study.

126

127 **Gastrointestinal transit time**

128 On Day -4 (relative to the antibiotic intervention) a study of the feed transit time was conducted to  
129 verify that the transit time in the gastrointestinal tract (GI-transit time) did not differ between the  
130 two groups before the intervention (Fig. 1). All mink received 100 plastic Hama beads (2.5x2.5  
131 mm, assorted colours, Malte Haaning Plastic A/S), which have been used as a marker of GI-transit  
132 time elsewhere (Hernot *et al.* 2005). The beads were mixed in 50 g of feed in the morning. When  
133 the 50 g ration was ingested or after one hour the rest of the daily ration without beads was given.  
134 Two hours after beads were ingested the mink started defecating and faeces were collected every 15  
135 minutes. Beads were counted after washing the faeces in a sieve under running water. The time  
136 lapse until the first bead appeared in the faeces was considered the feed transit time for the  
137 individual animal.

138

139 **Collection of blood and faeces and biochemical analyses**

140 On Days -5, 2 and 7 blood was obtained by puncture of vena cephalica antebrachii and directly  
141 transferred to EDTA-coated (plasma) and non-coated (serum) tubes (BD Vacutainer Systems,  
142 Preanalytical Solutions, Belliver Industrial Estate, UK). Tubes were centrifuged at 2000 G for 10

143 minutes and isolated plasma and serum was transferred to 0.5 ml eppendorf tubes, respectively.  
144 Samples for analysing biochemical parameters in serum as well as parameters related to lipid  
145 metabolism in plasma were kept at 5°C and analysed by Advia® 1800 Clinical Chemistry System,  
146 Siemens, at Centrallaboratoriet, Department of Veterinary and Animal Sciences, University of  
147 Copenhagen, Denmark. Serum samples for analysing acute phase reactant serum amyloid A were  
148 stored at -18°C and analysed by a standard ELISA-kit (Phase SAA assay, Tridelta Development  
149 Ltd., Kildare, Ireland) at the National Veterinary Institute at the Technical University of Denmark.  
150 Plasma samples for analysing vitamin A and E were stored at -18°C and subsequently analysed by  
151 PerkinElmer Series 200 HPLC System at Foulum, University of Aarhus, Denmark (Jensen, Engberg  
152 and Hedemann 1999).

153 On Days -5 and 2 faeces was collected from a sterile plastic drape placed under the cages using  
154 disposable forceps and directly transferred to cryovials after voiding (CryoPure Tube 1.8 ml,  
155 SARSTEDT, Germany). Samples were kept on dry ice until they were transferred to storage at -  
156 80°C. On Day 7 all mink were euthanized by initial sedation with a mixture of 0.6 ml ketamine  
157 (Ketaminol® Vet. 50 mg ml<sup>-1</sup>, MSD Animal Health) and medetomidin (Domitor® Vet. 1.0 mg ml<sup>-1</sup>,  
158 Orion Pharma Animal Health) administered intramuscularly followed by an intracardiac injection  
159 with 1 ml pentobarbital (Euthanimal 200 mg ml<sup>-1</sup>, corresponding to 182 mg ml<sup>-1</sup> pentobarbital,  
160 ScanVet Animal Health A/S).

161

## 162 **Digestibility trial**

163 A digestibility trial was performed on all animals during the intervention period Day 0 – Day 4  
164 (Fig. 1). Total faeces from each animal was collected separately once a day in the morning from  
165 a clean chute placed under the wire cage. The faeces was transferred to an individual foil tray  
166 and stored at -18°C. In addition, feed not consumed was also collected for each animal



167 transferred to a foil tray and stored at -18°C. Accumulated feed weighing between 20 and 100 g  
168 at the end of the trial, was analysed for content of dry matter. Faeces and samples of the feed  
169 were analysed for dry matter, protein, fat and ash. Chemical analysis was made in accordance  
170 with EU regulations (Commission Regulation 152/09/EC, 2009) at the analysis laboratory at  
171 Dansk Pelsdyr Foder A/S. Calculations of the apparent digestibility was made in accordance  
172 with existing literature (Ahlstrom & Skrede 1998).

173

#### 174 **Dissection of animals**

175 Dissection was performed *lege artis* immediately after euthanasia. Tissue samples from the liver,  
176 the duodenum as well as the colon and the mesenteric lymph node were removed and transferred to  
177 10% neutral buffered formalin prior to processing for histology. Intestinal content obtained from the  
178 distal colon was collected and stored as described for faecal samples by the use of a disposable  
179 forceps. The collection and storage of mucus (and associated bacteria) was done as previously  
180 described (Bahl *et al.* 2017). After processing and haematoxylin and eosin staining, liver sections  
181 were graded for hepatic steatosis based on a scoring system applied for classification of non-  
182 alcoholic fatty liver disease (Brunt 2007), while the intestine and the mesenteric lymph node were  
183 evaluated in the software programme ZEN 2.3 blue edition (Carl Zeiss microscopy GmbH, 2011,  
184 Göttingen, Germany) after scanning by Zeiss type Axiostar Plus (Carl Zeiss; Göttingen, Germany).  
185 For the section of intestine to be evaluated a minimum of five representative areas on each slide  
186 were to be present.

187

#### 188 **Extraction of bacterial community DNA**

189 Total community DNA from mucus, faeces, and feed samples was extracted using the MoBio  
190 Power Soil®-htp 96 Well Soil DNA Isolation Kit (MoBio Laboratories, Carlsbad, CA) according to

191 the manufacturer's recommendations, with minor modifications as previously reported (Tulstrup *et*  
192 *al.* 2015). DNA concentrations were measured with the Quant-iT dsDNA HS kit (Life  
193 Technologies).

194

### 195 **Bacterial community composition and diversity**

196 The bacterial community composition was determined by partial 16S rRNA gene sequencing of the  
197 extracted community. DNA amplification of the V3-region was performed following a dual PCR  
198 strategy with universal bacterial primers; PBU (5'-CCTACGGGAGGCAGCAG-3') and PBR (5'-  
199 ATTACCGCGGCTGCTGG-3') as described previously (Bahl *et al.* 2017). Sequencing of the 16S  
200 rRNA gene libraries was performed on the Ion OneTouch<sup>TM</sup> and Ion PGM platform with a 318-  
201 Chip v2. Sequencing was performed by the DTU in-house facility (DTU Multi-Assay Core  
202 (DMAC), Technical University of Denmark. Sequencing data were imported into CLC Genomic  
203 Workbench (version 8.5. CLC bio, Qiagen, Aarhus, Denmark) and were de-multiplexed trimmed to  
204 remove barcodes and PCR primers. Quality filtering (maxee 1.5), dereplication, OTU clustering  
205 (minsize 2), chimera filtering (RDP\_gold database), mapping of reads to OTUs (97% similarity)  
206 and generation of an OTU table was done according to the UPARSE pipeline (Edgar 2013),  
207 generating a total of 2254 non-chimeric OTUs. Taxonomy was assigned using the RDP  
208 multiclassifier ver. 2.10.1 with a confidence threshold set to 0.5 recommended for sequences shorter  
209 than 250bp (Wang *et al.* 2007). A phylogenetic tree was created with FastTree based on PyNAST  
210 alignment of representative OTU sequences with an archaea (*Methanosarcina*) sequence added as  
211 outgroup for rooting. In QIIME (Caporaso *et al.* 2010), the OTU table was filtered to exclude OTUs  
212 classified as Cyanobacteria/Chloroplast and to exclude OTUs with average relative abundance  
213 below 0.005% of the total community (Bokulich *et al.* 2013), resulting in 377 OTUs. Relative taxon  
214 abundances and alpha diversity (Observed OTUs and Shannon diversity index) and beta diversity

215 based on unweighted UniFrac distances were calculated in QIIME (core\_diversity\_analyses.py),  
216 with the sequencing depth rarefied to 4,700 sequences per sample. Sequencing data are deposited in  
217 the NCBI Sequence Read Archive with the accession number SRP110733.

218

## 219 **Data handling and statistics**

220 Statistical analysis was performed in GraphPad Prism (version 7.0b; GraphPad Software Inc., La  
221 Jolla, CA). Differences between groups were assessed by paired t-test, unpaired Student's t-test or t-  
222 test with Welch correction if data had unequal variances. For non-parametric data, Mann-Whitney,  
223 Wilcoxon or Fischer's exact test were used as appropriate. Levels of biochemical blood parameters  
224 and vitamins were compared between groups on separate days (Day -5, Day 2, and Day 7) as well  
225 as through a trend analysis comparing differences between the sampling days. Levels of acute phase  
226 reactant serum amyloid A, and parameters related to lipid metabolism were compared between the  
227 two groups at the end of the study (Day 7). Statistical significance threshold was set at  $P < 0.05$ .  
228 Identification of differentially abundant bacterial taxa based on 16S rRNA gene sequencing was  
229 performed by permutation based tests adjusting for multiple comparisons, with a false discovery  
230 rate threshold  $q = 0.05$  (Pike 2011).

231

## 232 **Results**

### 233 **Effects on the host animals**

234 No morbidity or mortality was recorded during the animal experiment and no gross pathological  
235 findings were recorded in necropsy procedures.

236 At Day -4 prior to the intervention, no differences were found in gastro-intestinal transit time  
237 between the CON group and the AMX group ( $2.9 \pm 0.65$  and  $3.1 \pm 0.98$  h, respectively) (Fig. 2A).

238 Generally, mink in both groups lost body mass during the study (Day -5 to Day 7), but no

239 difference was seen between groups (Table 1) and no significant difference in feed intake was  
240 observed during the intervention (Fig. 2B) or before treatment (data not shown).

241 In the present study, no apparent changes in histomorphological parameters related to neither the  
242 duodenum nor colon were seen between the two groups after the treatment (Table 1). Furthermore,  
243 no reaction in the mesenteric lymph nodes was recorded, and no difference in grading of hepatic  
244 steatosis was present between groups (data not shown).

245 Treatment with amoxicillin for 7 days had no significant influence on uptake of fat-soluble  
246 vitamins A and E, the level of acute phase reactant serum amyloid A, nor parameters related to  
247 lipolysis e.g. triglycerides and non-esterified fatty acids (data not shown), however, significantly  
248 higher levels of  $\beta$ -hydroxybutyrate was found in the AMX group on Day 7 (Fig. 2C). There were no  
249 differences between groups in the parameters included in the biochemical profile (data not shown).

250 A slight increase in digestibility of fat was seen in the group of mink receiving amoxicillin when  
251 compared to the control group during the treatment period (Fig. 2D-F). Data from one control mink  
252 in the digestibility trial was excluded from the study because of extremely abnormal values.

253

#### 254 **Effects on the microbiota**

255 On Day -5 before the treatment period, no differences in the community composition of the faecal  
256 microbiota between the CON and AMX groups were observed (Fig. 3 and 4). Furthermore no  
257 differences in alpha diversity indices were found between groups before the intervention (Fig. 5)  
258 and also no separation of groups according to beta diversity was seen on Day -5 as assessed by  
259 PCoA analysis of unweighted UniFrac distances (Fig. 6). After two days of amoxicillin treatment  
260 (Day 2) the number of observed species in the AMX group was significantly higher than in the  
261 CON group ( $P < 0.05$ ), and the Shannon index was also higher although not significantly (Fig. 5A-  
262 B). Principle coordinate analysis of beta diversity showed that faecal samples obtained from

263 amoxicillin treated animals on Day 2 appeared to cluster together with feed samples and separate  
264 from the control animals (Fig. 6B), which is further supported by community composition  
265 histograms, where samples from the AMX group Day 2 and Feed samples appear very similar (Fig.  
266 3). On Day 2 the relative abundance of numerous bacterial genera were different between the CON  
267 and AMX groups (Fig. 4; Table S1, Supporting Information).

268 Analysis of mucus associated bacterial community obtained at the end of the treatment period  
269 (Day 7) revealed significantly lower alpha diversity in the AMX group compared to the CON group  
270 in terms of both number of observed species and Shannon diversity index (Fig. 5A-B) which  
271 appeared consistent with the community composition analysis based on relative abundance of  
272 bacterial classes (Fig. 3) as well as the separation of the two groups by PCoA (Fig. 6D). Three  
273 bacterial genera belonging to the Clostridia class, namely *Clostridium sensu stricto*, *Anaerococcus*  
274 and *Clostridium XI*, were found to have a lower relative abundance in the mucus layer in the AMX  
275 group compared to the CON group.

276 No faecal samples were obtained on Day 7; however, intestinal luminal samples were taken with  
277 disposable forceps directly from the distal part of the colon during the dissection. These samples  
278 appeared similar to mucus samples obtained on the same day in terms of bacterial composition (Fig.  
279 3). The same tendency is also seen in beta-diversity (principal coordinate analysis) where intestinal  
280 samples from Day 7 and mucosa samples from Day 7 appear to be similar (Fig. 6C-D), which may  
281 be due to the fact that the colon contained a sparse amount of luminal content making it difficult to  
282 avoid collecting mucus in those samples. For this reason, luminal samples obtained from Day 7 are  
283 not included in further analysis, as they are not deemed comparable to faecal sample taken on Day -  
284 5 and Day 2.

285

286 **DISCUSSION**

287 To our knowledge this is the first study to demonstrate that the intestinal microbial community  
288 composition of mink, changes considerably during treatment with amoxicillin. However, during the  
289 short-term intervention, no changes in parameters included in the biochemical profile, vitamin A  
290 and E levels, level of acute phase reactant serum amyloid A, parameters related to the general lipid  
291 metabolism nor histomorphological parameters related to the gut nor the liver were observed. The  
292 general loss in body mass observed in both groups may be due to a less intensive feeding strategy as  
293 well as stress factors during the study period.

294 No difference in GI-transit time was observed prior to the intervention and the community  
295 composition in the faecal microbiota between the two groups before the intervention did also not  
296 differ (Fig. 2, 3 and 4). Thus, the observed difference in microbiota composition during the  
297 administration of amoxicillin is a direct consequence of treatment. The early onset of changes in the  
298 gut microbiota after two days of treatment with amoxicillin is consistent with other studies  
299 (Cochetière *et al.* 2005). The observed increase in alpha diversity in the faecal microbiota after two  
300 days of treatment (Fig. 5) was surprising and not consistent with previous studies in dogs, rats and  
301 humans, which all show a significantly decreased microbial alpha diversity after treatment with  $\beta$ -  
302 lactams and vancomycin (Grønvold *et al.* 2010; Panda *et al.* 2014; Vrieze *et al.* 2014; Tulstrup *et al.*  
303 2015). In the present study we find it very likely that the inhibitory effect of amoxicillin combined  
304 with the very fast intestinal transit time in mink and the high bacterial load associated with mink  
305 feed (Bahl *et al.* 2017) result in a population structure in faecal samples from amoxicillin treated  
306 animals comparable to that found in the feed. As mink feed has a relatively high microbial alpha  
307 diversity in terms of both number of observed species and Shannon diversity this results in an  
308 apparent increased count of observed species in these animals, which is thus a reflection of  
309 allochthonous microbes from the feed ingredients (Savage 1977). Interestingly, within the mucosa  
310 the amoxicillin treatment has the opposite effect of reducing the alpha diversity significantly (Fig.

311 5) which is consistent with other studies (Grønvold *et al.* 2010; Panda *et al.* 2014; Vrieze *et al.*  
312 2014; Tulstrup *et al.* 2015). Within the mucosa (Day 7) a couple of specific genera within the  
313 Clostridia were found to be affected by the amoxicillin treatment after correcting for multiple  
314 testing, which is supported by the global shift in community composition (Fig. 6D); however, a  
315 much more pronounced effect was found at the genera level in the faecal samples obtained on Day  
316 2 (Fig. 4). This supports the notion that genetic material detected in the faecal samples on Day 2  
317 (AMX group) is predominantly from feed-associated bacteria and thus not colonizing microbes. To  
318 investigate whether these bacteria are living or dead, a quantification of genetic material could have  
319 been applied as performed in other studies (Pérez-Cobas *et al.* 2012). The observed effects within  
320 the mucosa-associated bacteria (Fig. 3, 4 and 6) occur despite the very fast transit time in the  
321 gastrointestinal tract in mink (Bleavins and Aulerich 1981; Szymeczko and Skrede 1990) allowing  
322 limited time for uptake or direct effect of antimicrobials administered in the feed. The observed  
323 faecal microbiota in the mink may be partly driven by the resident bacterial community in the  
324 mucosa and partly by feed associated bacteria. We therefore speculate that the colonizing  
325 microbiota associated with the mucus layer is substantially inhibited by amoxicillin on Day 2 since  
326 the composition ingested appears similar to that detected in the faeces. An increased renewal of the  
327 colonic mucosa is essential given the fast transit time in the gastrointestinal tract. This is consistent  
328 with a recent study showing that a fast transit time is associated with an increased mucosal turnover  
329 in the gut and generally related to the bacterial composition and diversity (Roager *et al.* 2016). As  
330 there is a link between gastrointestinal-transit time and bacterial composition and diversity, it would  
331 have been interesting to have investigated transit time during the treatment period as well.

332 The elevated level of  $\beta$ -hydroxybutyrate in the AMX group could be caused by reduced appetite  
333 in this group even though no significant difference in feed intake was observed (Fig. 2). Fasting is  
334 known to induce lipolysis from adipose tissue and the formation of ketone bodies (VanItallie and

335 Nufert 2003); however, no alterations in triglycerides or non-esterified fatty acids were observed as  
336 would have been expected in the case of a physiological reaction to fasting (Mustonen *et al.* 2005;  
337 Rouvinen-Watt *et al.* 2010).

338 In the present study, no changes were seen in histomorphological parameters related the gut and  
339 the liver regarding characteristics of villus-crypt complex, goblet cells, crypt depth, thickness of  
340 tunica muscularis or grading of hepatic steatosis, respectively. A previous study with broilers,  
341 which are also characterised by having a fast GI-transit time (Ferrando *et al.* 1987), reported  
342 increased villus height and improved performance, in terms of significant weight gain after  
343 antibiotic growth promoter administration (Sayrafi *et al.* 2011).

344 Maintenance of the host barrier function in the gut is enhanced by the commensal bacterial  
345 community, which forms a colonization barrier by e.g. competitive metabolic interactions,  
346 occupying intestinal niches and induction of the host immune system (Kamada *et al.* 2013).  
347 Furthermore some metabolites, namely butyric acid, from the digestion of oligosaccharides serve as  
348 an important energy source for colonocytes (Wang *et al.* 2012) and modulate tight-junctions near  
349 the apical surface of epithelial cells (Ohata, Usami and Miyoshi 2005). A compromised barrier  
350 function may lead to the invasion of bacteria or bacterial by-product initiating an inflammatory  
351 response in the host. Results from the present study, however, did not show any increased levels of  
352 acute phase reactant serum amyloid A in the amoxicillin treated animals, which is a marker of  
353 inflammation (Jain, Gautam and Naseem 2011).

354 In broilers, antimicrobials have been shown to enhance apparent absorption of alpha-tocopheryl  
355 acetate (vitamin E) and fatty acids due to decreased deconjugation of bile acids by *Clostridium*  
356 *perfringens* (Knarreborg *et al.* 2004). In the current study, the amoxicillin treated animals showed a  
357 lower proportion of *Clostridium* sensu stricto (group containing *C. perfringens*) and a slightly  
358 higher digestibility with no effect on protein or carbohydrate digestibility nor the levels of vitamins



359 A and E. According to Hedemann et al. (Hedemann, Clausen and Jensen 2011) the biliary bile acids  
360 are exclusively taurine conjugated, but it is not known to which extent they are deconjugated by the  
361 gut microbiota.

362 It has been reported for laboratory rodents that antibiotic-induced perturbations of the intestinal  
363 microbiota may alter the host susceptibility to enteric infection (Stecher *et al.* 2007; Sekirov *et al.*  
364 2008; Endt *et al.* 2010). It would be of great practical relevance for the mink production to  
365 investigate whether the observed alterations in the gut microbiota following antimicrobial treatment  
366 in mink are associated with increased susceptibility to pathogens. This may especially be relevant in  
367 young animals. The current study contributes with useful information regarding experimental  
368 design for future studies within this field. Additionally, our findings may contribute to the  
369 understanding of gut health in other carnivorous animals subjected to antibiotic treatment.

370

## 371 FUNDING

372 This work was supported by the Danish Fur Breeders Research Foundation and the University of  
373 Copenhagen.

374

## 375 ACKNOWLEDGEMENTS

376 We thank Susanne Kronborg, Bettina Gjedsted Andersen, Elisabeth Wairimu Petersen, Peter  
377 Heegaard and Ronja Mathiesen for technical assistance and Marlene Danner Dalgaard at the DTU  
378 Multi-Assay Core facility for performing the 16S rRNA sequencing. The authors declare no conflict  
379 of interest.

380

## 381 REFERENCES

382 Albenberg LG, Wu GD. Diet and the intestinal microbiome: Associations, functions, and

383 implications for health and disease. *Gastroenterology* 2014;**146**:1564–72.

384 Bahl MI, Hammer AS, Clausen T *et al.* The gastrointestinal tract of farmed mink (*Neovison vison*)  
385 maintains a diverse mucosa-associated microbiota following a 3-day fasting period. *Microbiol*  
386 *Open* 2017, DOI: 10.1002/mbo3.434.

387 Bleavins MR, Aulerich RJ. Feed consumption and food passage time in mink (*Mustela vison*) and  
388 European ferrets (*Mustela putorius furo*). *Lab Anim Sci* 1981;**31**:268–9.

389 Bokulich NA, Subramanian S, Faith JJ *et al.* Quality-filtering vastly improves diversity estimates  
390 from Illumina amplicon sequencing. *Nat Methods* 2013;**10**:57–9.

391 Brunt EM. Pathology of fatty liver disease. *Mod Pathol* 2007;**20 Suppl 1**:S40-8.

392 Caporaso JG, Kuczynski J, Stombaugh J *et al.* QIIME allows analysis of high-throughput  
393 community sequencing data. *Nat Methods* 2010;**7**:335–6.

394 Chriél M, Harslund JE, Jensen TH *et al.* *Antimicrobials in Mink—consumption and Resistance*  
395 *Patterns: 2012. Proceedings of the Xth International Scientific Congress in Fur Animal*  
396 *Production*. Larsen, P.F.; Møller, S.H.; Clausen, T.; Hammer, A.S.; Lásen, T.M.; Nielsen,  
397 V.H.; Tauson, A.H.; Jeppesen, L.L.; Hansen, S.W.; Elnif, J.; Malmkvist J (ed.). Wageningen  
398 Academic Publishers, 2012.

399 Cochetière MFD La, Durand T, Lepage P *et al.* Resilience of the Dominant Human Fecal  
400 Microbiota upon Short-Course Antibiotic Challenge. *J Clin Microbiol* 2005;**43**:5588.

401 Commission Regulation 152/09/EC on laying down the methods of sampling and analysis for the  
402 official control of feed O.J. L 54. *Off J Eur Union* 2009.

403 Davis CP, Cleven D, Balish E *et al.* Bacterial association in the gastrointestinal tract of beagle dogs.  
404 *Appl Environ Microbiol* 1977;**34**:194–206.

405 Dethlefsen L, Huse S, Sogin ML *et al.* The pervasive effects of an antibiotic on the human gut  
406 microbiota, as revealed by deep 16s rRNA sequencing. *PLoS Biol* 2008;**6**:2383–400.

407 Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods*  
408 2013;**10**:996–8.

409 Edlund C, Nord CE. Effect on the human normal microflora of oral antibiotics for treatment of  
410 urinary tract infections. *J Antimicrob Chemother* 2000;**46 Suppl 1**:41-48-65.

411 Endt K, Stecher B, Chaffron S *et al.* The Microbiota Mediates Pathogen Clearance from the Gut  
412 Lumen after Non-Typhoidal Salmonella Diarrhea. *PLoS Pathog* 2010;**6**:e1001097.

413 Ferrando C, Vergara P, Jiménez M *et al.* Study of the rate of passage of food with chromium-  
414 mordanted plant cells in chickens (*Gallus gallus*). *Q J Exp Physiol* 1987;**72**:251–9.

415 Garcia-Mazcorro JF, Dowd SE, Poulsen J *et al.* Abundance and short-term temporal variability of  
416 fecal microbiota in healthy dogs. *Microbiologyopen* 2012;**1**:340–7.

417 Grønvold AMR, L’Abée-Lund TM, Sørum H *et al.* Changes in fecal microbiota of healthy dogs  
418 administered amoxicillin. *FEMS Microbiol Ecol* 2010;**71**:313–26.

419 Handl S, Dowd SE, Garcia-Mazcorro JF *et al.* Massive parallel 16S rRNA gene pyrosequencing  
420 reveals highly diverse fecal bacterial and fungal communities in healthy dogs and cats. *FEMS*  
421 *Microbiol Ecol* 2011;**76**:301–10.

422 Hedemann MS, Clausen TN, Jensen SK. Changes in digestive enzyme activity, intestine  
423 morphology, mucin characteristics and tocopherol status in mink kits (*Mustela neovision*)  
424 during the weaning period. *Animal* 2011;**5**:394–402.

425 Hernot DC, Biourge VC, Martin LJ *et al.* Relationship between total transit time and faecal quality  
426 in adult dogs differing in body size. *J Anim Physiol Anim Nutr (Berl)* 2005;**89**:189–93.

427 Jain S, Gautam V, Naseem S. Acute-phase proteins: As diagnostic tool. *J Pharm Bioallied Sci*  
428 2011;**3**:118–27.

429 Jensen SK, Engberg RM, Hedemann MS. All-rac-alpha-tocopherol acetate is a better vitamin E  
430 source than all-rac-alpha-tocopherol succinate for broilers. *J Nutr* 1999;**129**:1355–60.

431 Jensen VF, Sommer HM, Struve T *et al.* Factors associated with usage of antimicrobials in  
432 commercial mink (*Neovison vison*) production in Denmark. *Prev Vet Med* 2016;**126**:170–82.

433 Jernberg C, Lofmark S, Edlund C *et al.* Long-term ecological impacts of antibiotic administration  
434 on the human intestinal microbiota. *ISMEJ* 2007;**1**:56–66.

435 Jones B V., Begley M, Hill C *et al.* Functional and comparative metagenomic analysis of bile salt  
436 hydrolase activity in the human gut microbiome. *Proc Natl Acad Sci U S A* 2008;**105**:13580–5.

437 Jørgensen, G.; Glem Hansen N. A Cage designed for Metabolism- and Nitrogen Balance Trials with  
438 Mink. *Acta Agric Scand* 1973;**23**:3–4.

439 Kamada N, Chen GY, Inohara N *et al.* Control of pathogens and pathobionts by the gut microbiota.  
440 *Nat Immunol* 2013;**14**:685–90.

441 Knarreborg A, Lauridsen C, Engberg RM *et al.* Dietary antibiotic growth promoters enhance the  
442 bioavailability of alpha-tocopheryl acetate in broilers by altering lipid absorption. *J Nutr*  
443 2004;**134**:1487–92.

444 LeBlanc JG, Milani C, de Giori GS *et al.* Bacteria as vitamin suppliers to their host: A gut  
445 microbiota perspective. *Curr Opin Biotechnol* 2013;**24**:160–8.

446 Matamoros S, Gras-Leguen C, Le Vacon F *et al.* Development of intestinal microbiota in infants  
447 and its impact on health. *Trends Microbiol* 2013;**21**:167–73.

448 Mustonen AM, Paakkonen T, Ryökkynen A *et al.* Adaptations to fasting in the American mink  
449 (*Mustela vison*): Carbohydrate and lipid metabolism. *Comp Biochem Physiol - A Mol Integr*  
450 *Physiol* 2005;**140**:195–202.

451 Ohata A, Usami M, Miyoshi M. Short-chain fatty acids alter tight junction permeability in intestinal  
452 monolayer cells via lipoxygenase activation. *Nutrition* 2005;**21**:838–47.

453 Panda S, El khader I, Casellas F *et al.* Short-term effect of antibiotics on human gut microbiota.  
454 *PLoS One* 2014;**9**:e95476.

455 Pedersen K, Hammer AS, Sørensen CM *et al.* Usage of antimicrobials and occurrence of  
456 antimicrobial resistance among bacteria from mink. *Vet Microbiol* 2009;**133**:115–22.

457 Pérez-Cobas AE, Gosalbes MJ, Friedrichs A *et al.* Gut microbiota disturbance during antibiotic  
458 therapy: a multi-omic approach. *Gut* 2012;**62**:1–11.

459 Pike N. Using false discovery rates for multiple comparisons in ecology and evolution. *Methods*  
460 *Ecol Evol* 2011;**2**:278–82.

461 Raymann K, Shaffer Z, Moran NA. Antibiotic exposure perturbs the gut microbiota and elevates  
462 mortality in honeybees. *PLoS Biol* 2016:1–22.

463 Roager HM, Hansen LBS, Bahl MI *et al.* Colonic transit time is related to bacterial metabolism and  
464 mucosal turnover in the gut. *Nat Microbiol* 2016;**1**:16093.

465 Round J, Mazmanian S. The gut microbiota shapes intestinal immune responses during health and  
466 disease. *Nat Rev Immunol* 2014;**9**:313–23.

467 Rouvinen-Watt K, Mustonen A-M, Conway R *et al.* Rapid development of fasting-induced hepatic  
468 lipidoses in the American mink (*Neovison vison*): effects of food deprivation and re-  
469 alimentation on body fat depots, tissue fatty acid profiles, hematology and endocrinology.  
470 *Lipids* 2010;**45**:111–28.

471 Savage DC. Microbial Ecology Of The Gastrointestinal Tract. *Annu Rev Microbiol* 1977;**31**:107–33.

472 Sayrafi R, Soltanlinejad F, Shahrooz R *et al.* Effect of butyric acid glycerides and antibiotic growth  
473 promoter on the performance and intestinal histomorphometry of broiler chickens. *Journal*  
474 *Food, Agric Environ* 2011;**9**:285–8.

475 Sekirov I, Tam NM, Jogova M *et al.* Antibiotic-induced perturbations of the intestinal microbiota  
476 alter host susceptibility to enteric infection. *Infect Immun* 2008;**76**:4726–36.

477 Stecher B, Robbiani R, Walker AW *et al.* *Salmonella enterica* serovar typhimurium exploits  
478 inflammation to compete with the intestinal microbiota. *PLoS Biol* 2007;**5**:2177–89.

- 479 Suchodolski JS, Dowd SE, Westermarck E *et al.* The effect of the macrolide antibiotic tylosin on  
480 microbial diversity in the canine small intestine as demonstrated by massive parallel 16S  
481 rRNA gene sequencing. *BMC Microbiol* 2009;**9**:210.
- 482 Sullivan Å, Edlund C, Nord CE. Effect of new antimicrobial agents on the ecological balance of  
483 human microflora. *Lancet* 2001;**18**:249–53.
- 484 Szymeczko R, Skrede A. Protein digestion in mink. *Acta Agric Scand* 1990;**40**:189–200.
- 485 Tulstrup MVL, Christensen EG, Carvalho V *et al.* Antibiotic Treatment Affects Intestinal  
486 Permeability and Gut Microbial Composition in Wistar Rats Dependent on Antibiotic Class.  
487 *PLoS One* 2015;**10**:1–17.
- 488 VanItallie TB, Nufert TH. Ketones: metabolism’s ugly duckling. *Nutr Rev* 2003;**61**:327–41.
- 489 Vrieze A, Out C, Fuentes S *et al.* Impact of oral vancomycin on gut microbiota, bile acid  
490 metabolism, and insulin sensitivity. *J Hepatol* 2014;**60**:824–31.
- 491 Wang HB, Wang PY, Wang X *et al.* Butyrate enhances intestinal epithelial barrier function via up-  
492 regulation of tight junction protein claudin-1 transcription. *Dig Dis Sci* 2012;**57**:3126–35.
- 493 Wang Q, Garrity GM, Tiedje JM *et al.* Naive Bayesian classifier for rapid assignment of rRNA  
494 sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 2007;**73**:5261–7.
- 495 Williams C, Buddington RK, Elnif J. The Gastrointestinal Bacteria of Mink (*Mustela vison* L):  
496 Influence of Age and Diet. *Acta Vet Scand* 1998;**39**:473–82.

497

## 498 SUPPLEMENTARY DATA

499 Additional Supporting Information may be found in the online version of this article:

500

501 **Table S1.** Relative abundance of bacterial groups at different taxonomical levels for individual  
502 animals at timepoints Day -5, Day 2, Day 7 as well as Mucus- and Feed samples.

503

504 **Figure legends:**

505

506 **Fig. 1.** Timeline of experimental period. The acclimatization period was from Day -7 to Day 0 and  
507 treatment period from Day 0 to Day 6. All animals were euthanized on Day 7. Blood samples (**B**) as  
508 well as faecal samples (**F**) and mucosal samples (**M**) were collected as indicated. Gastrointestinal  
509 transit time analysis (**T**) was performed on Day -4 and pathological examination (**P**) was performed  
510 during dissection. A digestibility trial (box) was performed from Day 0 to Day 4 during the  
511 treatment period.

512

513 **Fig. 2.** Gastrointestinal transit time calculated at Day -4 (A), Percentage feed intake during the  
514 intervention calculated from Day 0-7 (B), serum  $\beta$ -hydroxybutyrate concentration determined on  
515 Day 7 (C) and digestibility of Fat (D), Protein (E) and Carbohydrate (F) during the intervention  
516 (Day 0-4). Boxplots show median and first and third quartile for the control group (CON, n=11-12),  
517 and the amoxicillin group (AMX, n=12) with whiskers indicating range. \*  $P < 0.05$ .

518

519 **Fig. 3.** Average bacterial community composition at the class level in faecal samples collected on  
520 Day -5, Day 2, and Day 7, as well as in mucus samples obtained on Day 7 for the control group  
521 (CON, n=12) and the amoxicillin group (AMX, n=11-12). The bacterial community composition in  
522 feed samples collected at the end of the study (Feed, n=5) is also shown.

523

524 **Fig. 4.** Heatmap of z-scores calculated based on average relative abundance within each row for the  
525 control group (CON, n=12) and the group receiving amoxicillin (AMX, n=11-12) on Day -5, Day 2  
526 and mucus sample (Day 7). The bacterial genera depicted represent above 25% prevalence  
527 collectively in all samples. Asterisks indicate significant difference between the control group and  
528 the antibiotic treatment group compared at the same treatment time (FDR corrected permutation

529 based t-test). \* P < 0.05, \*\* P < 0.01. The left-hand colour bar shows the taxonomic classification  
530 of bacterial genera at the class level (Orange: *Actinobacteria*, dark-green: *Bacteroidia*, light-gray:  
531 *Flavobacteria*, light-blue: *Bacilli*, blue: *Clostridia*, grey: *Erysipelotrichia*, dark-grey: *Negativicutes*,  
532 yellow: *Fusobacteria*, faint-pink:  $\alpha$ -*Proteobacteria*, salmon:  $\beta$ -*Proteobacteria*, dark-red:  $\chi$ -  
533 *Proteobacteria*, red:  $\gamma$ -*Proteobacteria* and violet: *Mollicutes*.

534

535 **Fig. 5.** Alpha-diversity of the bacterial communities in the faeces samples, mucosal scraps as well  
536 as feed samples. Bacterial richness (number of observed OTU's) (A) and Shannon diversity index  
537 (B) are shown for faecal samples obtained on Day -5 and Day 2, mucus samples obtained post  
538 mortem on Day 7 as well as feed samples obtained at the end of study. Columns show mean values  
539 of the control group (CON, n=12), the amoxicillin group (AMX, n=12) as well as feed samples  
540 (Feed, n=5) with standard error of the means indicated by error bars. Significant differences  
541 between groups are indicated by asterisks \* P < 0.05 and \*\* P < 0.01.

542

543 **Fig. 6.** Beta-diversity represented as principle coordinate analysis based on un-weighted UniFrac  
544 distance matrix at Day -5 (A), during the treatment period at Day 2 (B), at the end of treatment at  
545 Day 7 (C) as well as mucosa associated microbiota samples collected post-mortem at Day 7 (D).  
546 Each dot represents a faecal- or mucus sample from individual animals in the control group (blue,  
547 n=12) and amoxicillin group (red, n=12). Feed samples obtained at the end of the study are also  
548 included (grey, n=5).

549

550



551 **Tables**

552

553 Table 1. Animal body mass and histomorphological parameters of the intestine.

554


	<b>Control group</b>	<b>Amoxicillin group</b>	<b><i>p</i>-value</b>
Initial body mass (g) <sup>a</sup>	2789 ± 93	2829 ± 83	0.75
Final body mass (g) <sup>a</sup>	2661 ± 86	2652 ± 73	0.94
<i>p</i> -value	<0.001	<0.001	
Duodenum			
Villus-crypt complex (μm) <sup>b</sup>	1053 ± 85.6 (n=12)	1096 ± 76.2 (n=11)	0.78
Goblet cells <sup>c</sup>	27.1 ± 12.0 (n=12)	28.7 ± 12.3 (n=11)	0.93
Tunica muscularis (μm)	413.3 ± 15.0 (n=10)	424 ± 30.8 (n=11)	0.75
Colon			
Crypt depth (μm)	459.5 ± 25.5 (n=12)	409.9 ± 20.2 (n=11)	0.15
Tunica muscularis (μm)	932.7 ± 96.8 (n=10)	1117 ± 71.4 (n=12)	0.13

567 <sup>a</sup>Body mass measured at day -5 and 7 respectively. <sup>b</sup>Villus-crypt complex measured from villus568 apex to the base of the crypt. <sup>c</sup>Goblet cells counted on a 400 μm piece of villus apex. Means ±

569 SEM are shown.


570

Figure 1




# Figure 2


**A**




**B**




**C**




**D**




**E**




**F**



# Figure 3




# Figure 4




# Figure 5

## A



## B



# Figure 6

