

Technical University of Denmark



The *Campylobacter jejuni* contamination level in houseflies after exposure to materials contaminated with *Campylobacter*

Jensen, Annette Nygaard

Published in:

Proceeding of EAAP - 68th annual meeting

Publication date:

2017

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Jensen, A. N. (2017). The *Campylobacter jejuni* contamination level in houseflies after exposure to materials contaminated with *Campylobacter*. In Proceeding of EAAP - 68th annual meeting (pp. 666-666). Tallinn, Estonia.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Campylobacter contamination level in houseflies after exposure to materials containing *Campylobacter*

A.N. Jensen

Technical University of Denmark, National Food Institute, Division of Food Microbiology and Production, Anker Engélunds Vej 1, 2800 Kgs. Lyngby, Denmark; anyj@food.dtu.dk

Houseflies have been found to carry *Campylobacter jejuni* and play a role in infection of poultry flocks. This study aimed to elucidate the quantitative campylobacter level in naturally infected houseflies depending on their previous exposure to campylobacter-contaminated material in terms of campylobacter level, duration and type of material. Houseflies (*Musca domestica*) reared under laboratory conditions were placed in 250-ml cups containing 5 g of chicken faeces or 1 ml liquid and spiked with 3, 4, 5 or 7 Log cfu *C. jejuni*. Sixteen houseflies were added to each cup at a time. After approx. 1 h of exposure, 4 flies were removed from the cup for direct enumeration of campylobacter by plate spreading. Another 4 flies were transferred onto an Abeyta-Hunt-Bark (AHB) agar plate (9 cm) to assess if flies will contaminate surfaces. After 1 h on the plate, each fly was tested for campylobacter. This procedure was repeated after approx. 4 h of exposure for the remaining eight flies. The percentage of houseflies positive for campylobacter as well as the Log₁₀ cfu recovered per fly depended on the contamination level and the material exposed to. For faeces, 90.0% (n=80), 48.4% (n=64), 6.3% (n=48) and 0% (n=16) of flies were campylobacter positive when exposed to 7, 5, 4, and 3 Log₁₀ cfu with a mean (±SE) of 2.0±0.1, 0.8±0.1, 0.3±0.0 and 0 Log₁₀ cfu recovered per campylobacter-positive fly, respectively. For liquid, 95.7% (n=47), 91.4% (n=47), 20.8% (n=48) and 6.3% (n=16) of flies were campylobacter positive when exposed to 7, 5, 4, and 3 Log₁₀ cfu with a mean of 3.3±0.2, 2.0±0.1, 0.8±0.2 and 0.3±0.0 Log₁₀ cfu recovered per campylobacter-positive fly, respectively. *Campylobacter* seemed to be taken up readily as there was no significant effect of exposure time (1 vs 4 h) and the uptake was higher from contaminated liquid than faeces. The surface of the AHB plates was only contaminated by houseflies previously exposed to either 5 or 7 Log₁₀ cfu and most by flies exposed to liquid (45% n=48) opposed to faeces (31% n=72). The results support that houseflies are likely to become contaminated with campylobacter if exposed to material containing >4 Log₁₀ cfu.