

# AMCoR

Asahikawa Medical University Repository <http://amcor.asahikawa-med.ac.jp/>

Environmental Health and Preventive Medicine (2010) 15(5):276–284.

Dampness, food habits, and sick building syndrome symptoms in elementary school pupils

Saijo, Yasuaki ; Nakagi, Yoshihiko ; Ito, Toshihiro ;  
Sugioka, Yoshihiko ; Endo, Hitoshi ; Yoshida, Takahiko

1 **Dampness, Food Habits, and Sick Building Syndrome Symptoms among Elementary School**  
2 **Pupils**

3 \_\_\_\_\_

4 Yasuaki SAIJO, Yoshihiko NAKAGI, Toshihiro ITO, Yoshihiko SUGIOKA, Hitoshi ENDO,  
5 and Takahiko YOSHIDA

6

7 <sup>1</sup> *Department of Health Science, Asahikawa Medical College*

8

9

10 Reprint requests: Yasuaki SAIJO

11 Department of Health Science, Asahikawa Medical College, Midorigaoka, E2-1-1-1

12 Asahikawa, Hokkaido 078-8510, Japan

13 TEL: +81 166 68 2402, FAX: +81 166 68 2409

14 E-mail: y-saijo@asahikawa-med.ac.jp

1 **Abstract**

2           *Objective*     We investigated the dampness/mold in schools and dwellings, food habits,  
3 and subjective symptoms among elementary school pupils in order to clarify the relationship  
4 between dampness and food habits on subjective symptoms in elementary school pupils.

5           *Methods*     Questionnaires were used to investigate the dampness in classrooms and  
6 dwellings in Hokkaido, Japan, and its effect on subjective symptoms involving 1,077 pupils in 8  
7 elementary schools. A dampness index for both the home and classroom was the sum of the  
8 presence of four dampness indicators: (1) visible mold, (2) moldy odor, (3) water leakage, and (4)  
9 condensation on windowpanes. The questionnaire contained queries about food habits as follows:  
10 the frequency of eating breakfast, whether the energy provided by school lunch was sufficient, and  
11 whether too many snacks and/or sweets were consumed. Adjusted logistic regression was used to  
12 determine whether dampness and food habits were related to symptoms.

13           *Results*     The home dampness index was significantly related to cough, general symptoms,  
14 and having at least one of symptoms; the classroom dampness index was significantly related to  
15 nose symptoms in fully adjusted models. In addition, usually not eating breakfast was significantly  
16 related to eye symptoms, and too many snacks and/or sweets was significantly related to eye, nose,  
17 and general symptoms.

18           *Conclusions*     Both home and classroom dampness can affect pupils' health. Home

1 dampness, in particular, was significantly related to cough and general symptoms, and classroom  
2 dampness was significantly related to nose symptoms. Furthermore, favorable food habits have a  
3 positive effect on pupils' subjective symptoms.

4

5 **Keywords** Dampness, Mold, Sick Building Syndrome, School Pupils, Food Habits

1 **Introduction**

2           There has been much concern about the effect of indoor environment on residents' health.  
3 Many studies have demonstrated that building dampness and mold have adverse health effects on  
4 residents. [1-3]

5           The school environment affects children's health. Adverse health effects of dampness and  
6 mold in school buildings on school children have been reported. [4-7] Moreover, an  
7 interventional study, in which a moisture-damaged school was repaired, showed a positive effect  
8 on school children's health. [8] Meanwhile, home dampness and mold have adverse health effects  
9 on children, [9-11] and home remediation against moisture sources has a protective effect in  
10 asthmatic children. [12]

11           The dampness index is the sum of the presence of several dampness indicators. [13] Its  
12 relation to sick building syndrome (SBS) symptoms and asthmatic symptoms was initially  
13 reported in Swedish multifamily building studies. [13,14] We also have reported its significant  
14 relation to SBS symptoms in newly build dwellings [15-17] and public apartment houses. [18]  
15 However, to our knowledge, there has been no report on simultaneous evaluation of home and  
16 school dampness and their association with children's health.

17           It has been reported that psychosocial factors are associated with SBS. [19] Lifestyle  
18 factors—such as work time, sleep time, and alcohol consumption—are also related to SBS. [20]

1 Among children, the relationship of unhealthy eating patterns to unfavorable overall school  
2 performance has been reported, [21] though its relation to SBS has not been reported. Thus,  
3 lifestyle factors including food habits may affect children's subjective symptoms.

4 From the point of view of improving children's health and their overall school performance,  
5 it is important to clarify improvable environmental and lifestyle risk factors for children's subjective  
6 symptoms including SBS. In this study, we explored dampness in schools and dwellings, food  
7 habits and subjective symptoms among elementary school pupils in Hokkaido Prefecture (north  
8 island of Japan) to clarify the relationship between home dampness, classroom dampness or food  
9 habits and subjective symptoms.

1 **Methods**

2 Study Population

3           Nayoro city and Asahikawa city are located in Hokkaido, the north island of Japan. The  
4 eight elementary schools under study included all 7 municipal elementary schools in the Nayoro  
5 district of Nayoro city (the other district, Furen, had five) and one elementary school affiliated with  
6 the university in Asahikawa city. Nayoro is located near Asahikawa (60 kilometer apart), and both  
7 cities have a similar inland climate.

8           We distributed anonymous parent-administered questionnaires to 1,753 pupils through  
9 classroom teachers between November and December 2007. The questionnaires included questions  
10 about subjective symptoms, food habits, and dwelling characteristics. Each classroom teacher  
11 retrieved the questionnaires of his or her pupils. The classroom teachers were also asked to answer a  
12 questionnaire about classroom dampness. School characteristics were gathered from the vice  
13 principal of each school. Of the 1,753 pupils, 1,141 (65.1%) answered the questionnaires. We  
14 finally analyzed 1,077 (61.4%) questionnaires, after excluding questionnaires that had any missing  
15 information. All classroom characteristics questionnaires were answered by the classroom teachers  
16 (n = 61). This study was conducted after obtaining informed consent from all subjects and approval  
17 from the institutional ethical board for epidemiological studies at Asahikawa Medical College.

18

## 1 Subjective Symptoms, Food Habits, and Dwelling Characteristics

2 We used a modified parent-administered questionnaire, MM080 for schools, designed for  
3 epidemiological assessment of SBS involving elementary school pupils. [22] We translated the  
4 English version of MM080 into Japanese for schools, based on the Japanese version of MM040EA,  
5 which is a validated self-administered questionnaire and also designed for epidemiological  
6 assessment of adults SBS. [23] We translated the English version of MM080 into Japanese for  
7 schools, based on the Japanese version of MM040EA. The questionnaire contained information on  
8 grade, sex, food habits, dwelling characteristics, and subjective symptoms.

9 Symptoms surveyed during the preceding three months were as follows: fatigue; headache;  
10 sleep problems; itching, burning, or irritation of the eyes; irritated, stuffy, or runny nose; cough; dry  
11 or flushed facial skin; scaling/itching of the scalp or ears; and dry, itching, or red-skinned hands.  
12 Each question had three alternative answers: *Yes, often* (every week); *Yes, sometimes*; and *No, never*.  
13 An additional query concerning the attribution of a symptom to the child's school environment,  
14 present in the original questionnaire, [22] was not used in this study.

15 Symptoms that occurred often (weekly) were defined as positive. For the analysis,  
16 symptoms were categorized as follow: general symptoms (fatigue, headache, and sleep problems)  
17 and symptoms involving the eyes, nose, cough, and skin. The questionnaires also contained queries  
18 about the history of asthma, hay fever, eczema, and food allergies in the previous year.



1           Queries about food habits concerned with the frequency of eating breakfast (always, almost  
2 always, 1–3 times per week, never : later two categories were defined as “usually not eating  
3 breakfast”), whether the energy provided by school lunch was sufficient or not (yes/no), and  
4 whether too many snacks and/or sweets were consumed (yes/no).

5           The self-administered questionnaire contained queries about the housing type (solitary or  
6 other), year of construction (before 1980, later, or unknown), heating (electric stove or other; gas,  
7 kerosene, or wood stove), ventilation (natural, mechanical, or unknown), wall-to-wall carpeting in a  
8 child’s room (yes/no), furry animals or birds in the home (yes/no), a smoker in the home (yes/no),  
9 anyone smoking indoors at home (yes/no), signs of moisture/mold damage (yes/no), and  
10 condensation on windowpanes (yes/no). In the original MM080 for schools, the year of construction  
11 of dwelling was categorized as *before 1960 or later*, but we revised it to *before 1980 or later*  
12 because the life span of Japanese buildings is relatively short. We considered that one of the original  
13 query, *Are there signs of moisture/mold damage*, was difficult for Japanese participants to  
14 understand. Therefore, we used queries about visible mold (yes/no), perception of moldy odor  
15 (yes/no), and episodes of water leakage during past 5 years (yes/no).

16           We constructed a home dampness index. The index considered four dampness indicators: (1)  
17 visible mold, (2) perception of moldy odor, (3) episodes of water leakage during past 5 years, and  
18 (4) condensation on windowpanes. The home dampness index was estimated by the sum of the

1 presence of these indicators.

2

### 3 Classroom Characteristics

4 We distributed the questionnaire about classroom characteristics to classroom teachers. It  
5 contained queries about the number of pupils, perception of visible mold (yes/no), moldy odor  
6 (yes/no), episodes of water leakage during past 5 years (yes/no), and condensation on windowpanes  
7 (yes/no). By using these results, we calculated a class dampness index. It was estimated by the  
8 sum of the presence of four dampness indicators in the classroom, the indicators are: (1) visible  
9 mold, (2) perception of moldy odor, (3) episodes of water leakage during past 5 years, and (4)  
10 condensation on windowpanes.

11

### 12 Statistical Analysis

13 The prevalence of subjective symptoms in boys and girls were compared by the  $\chi^2$  test. Odds  
14 ratios (ORs) for each subjective symptom were analyzed using logistic regression. To obtain  
15 multivariate adjusted ORs for each subjective symptoms of home and classroom dampness  
16 indicators, their dampness indexes, and food habits, we controlled the possible confounders,  
17 including age, sex, 1-year history of allergic diseases, housing type (solitary or other), construction  
18 year of dwelling ( $\leq 1980$ ,  $>1980$ , or unknown), type of heating of dwelling (electric or other),

1 ventilation of dwelling (mechanical, natural, or unknown), construction year of school, ventilation  
2 of school, and number of pupils in class (<20, 21–30, or >31), and each dampness indicator and  
3 food habit was introduced separately in the model. Dampness indicators, dampness indexes, and  
4 food habits were introduced separately in the model. Moreover, to estimate statistical dose-response  
5 relationships, *P* values for trends of home and classroom dampness indexes were analyzed.

6 Finally, fully adjusted model analyses were performed to obtain the ORs of home and  
7 classroom dampness indexes and food habits after adjustment of age, sex, 1-year history of allergic  
8 diseases, housing type (solitary or other), construction year of dwelling ( $\leq 1980$ ,  $>1980$ , or  
9 unknown), type of heating of dwelling (electric or other), ventilation of dwelling (mechanical,  
10 natural, or unknown), construction year of school, ventilation of school, and number of pupils in  
11 class (category).

12 For all statistical analyses, a 5% level of significance was applied. All statistical analyses  
13 were conducted using SPSS for Windows version 17.0 (SPSS, Inc., Chicago, U.S.A.).

## 1 **Results**

2           Table 1 shows the characteristics of pupils and their homes. The status of home dampness  
3 was as follow: 29.0% had visible mold, 8.3% had moldy odor, 16.6% had water leakage, and 55.2%  
4 had condensation on windowpanes. Food habits were as follows: 2.6% usually did not eat breakfast  
5 (1–3 times per week or never), 9.0% had no sufficient energy following a school lunch, and 22.7%  
6 had too many snacks and/or sweets.

7           Table 2 shows the characteristics of schools and classrooms. The status of classroom  
8 dampness was as follow: 37.5% had visible mold, 25.0% had moldy odor, 37.5% had water leakage,  
9 and 37.5% had condensation on windowpanes.

10           Table 3 shows subjective symptoms observed in this survey. Eye symptoms, nose symptoms,  
11 cough, skin symptoms, and general symptoms were found in 4.4%, 14.0%, 4.8%, 11.3%, and 4.8%  
12 of children, respectively. Any symptoms (at least one of skin, eye, nose, throat, or general  
13 symptoms) were found in 25.2% of children. Boys had a significantly higher prevalence of nose  
14 symptoms and any symptoms compared to girls.

15           Table 4 shows the ORs of the dampness indicator and food habits for subjective symptoms  
16 adjusted by age, sex, allergy, construction year of dwelling, housing type (solitary or other),  
17 ventilation of dwelling, type of heating of dwelling, construction year of school, ventilation of  
18 school, and number of pupils per class. Usually not eating breakfast had significantly higher ORs

1 for eye symptoms. Too many snacks and/or sweets had significantly higher ORs for eye, nose,  
2 general symptoms, and any symptoms. Water leakage of dwelling had significantly higher ORs for  
3 cough, general symptoms, and any symptoms. The home dampness index had significantly higher  
4 ORs for cough, general symptoms, and any symptoms. The classroom dampness index had  
5 significantly lower ORs for general symptoms and marginally significantly higher ORs for cough.

6 Table 5 shows fully adjusted ORs of home and classroom dampness indexes and food habits  
7 for subjective symptoms. Usually not eating breakfast had a significantly higher OR for eye  
8 symptoms. Too many snacks and sweets had significantly higher ORs for eye, nose, and general  
9 symptoms. The home dampness index had significantly higher ORs for cough ( $P$  for trend = 0.007;  
10 dampness index 4 vs. 0: OR 5.43, 95% CI: 1.41–20.9); general symptoms (dampness index 4 vs. 0:  
11 OR 3.57, 95% CI: 1.51–12.5); and any symptoms ( $P$  for trend = 0.016, dampness index 4 vs. 0: OR  
12 2.92, 95% CI: 1.22–6.99). Classroom dampness index had a significantly higher OR for nose  
13 symptoms ( $P$  for trend = 0.040).

## 1 **Discussion**

2           In this study, we found significant relationships between both home and classroom  
3 dampness and subjective symptoms. In fully adjusted models, home dampness index was  
4 significantly related to cough, general symptoms, and any symptoms, and classroom dampness  
5 index was significantly related to nasal symptoms. In addition, usually not eating breakfast was  
6 significantly related to eye symptoms, and too many snacks and/or sweets was significantly related  
7 to eye, nose, and general symptoms. To our knowledge, this is the first report on the relationship  
8 between both home and classroom dampness and subjective symptoms of pupils with  
9 simultaneous evaluation of home and classroom dampness.

10           For the difference between sexes, boys had more symptoms than girls in this study. In a  
11 previous study, no significant difference in the prevalence of atopic dermatitis between boys and  
12 girls in Japan has been reported. [24] Meanwhile, the prevalence of asthma and rhinitis in boys is  
13 greater than that in girls. [25-27] Similar to previous studies, our study showed a higher symptom  
14 prevalence among boys than girls.

15           We used the MM080 questionnaire. [22] There has been no report on the prevalence of  
16 symptoms in pupils using the MM080 for schools in Japan. The MM080 developers reported that  
17 the prevalence rate of each symptom was 2–7% ([http://www.orebroll.se/uso/page\\_\\_\\_17918.aspx](http://www.orebroll.se/uso/page___17918.aspx)).  
18 The prevalence of nasal symptoms found in this study was rather high. The reason for this was

1 probably because allergic rhinitis has been the most prevalent allergic disease and its prevalence has  
2 been increasing in Japan. [25]

3 In a previous study of newly built dwellings in Hokkaido, Japan, visible mold and  
4 condensation on windowpanes and/or walls was 15.6% and 41.7%, respectively. [16] Another study  
5 of newly built dwellings in Japan (including six prefectures), found visible mold, moldy odor, water  
6 leakage, and condensation on windowpanes and/or walls in 39.8%, 7.8%, 4.7%, and 51.8% of  
7 dwellings, respectively. [15] A study of old apartment houses in Hokkaido, Japan, found visible  
8 mold, moldy odor, water leakage, and condensation on windowpanes in 59.5%, 62.1%, 20.8%, and  
9 81.8% of dwellings, respectively. [18] Meanwhile, in a Swedish multifamily dwellings study, moldy  
10 odor, water leakage, and condensation on windowpanes, was 12.4%, 12.7%, and 9.0%, respectively.  
11 [13] Moreover, in the Swedish dwellings study, water leakage (during a previous year), visible  
12 dampness (mold or damp spot), and condensation on windowpanes was seen in 17.8%, 1.5%, and  
13 14.3% [9] of dwellings, respectively. Thus, Japanese dwellings have higher dampness rates.  
14 Meanwhile, the annual average of outdoor temperature and relative humidity in Asahikawa and  
15 Stockholm were 6.7°C and 76% and 6.7°C and 76%, respectively (World Meteorological  
16 Organization; Japan Meteorological Agency). However, inadequate waterproofing of the  
17 constructions, water leakage, and moisture from humans and indoor activities are important  
18 contributors to building dampness, [2] and those differences between Japan and Nordic countries

1 are little investigated. One possible explanation of these differences is that many Japanese people  
2 regard low humidity as a risk factor for respiratory infections, such as influenza virus infection in  
3 winter [28] and adenovirus infection in summer, [29] and may prefer high humidity, especially in  
4 winter. However, further studies will be needed to clarify the cause of these differences in dampness  
5 and preferable constructions and indoor activities.

6         The home dampness index was significantly associated with cough (3 and 4 vs. 0; *P* for  
7 trend), general symptoms (4 vs. 0), and any symptoms (4 vs. 0; *P* for trend), and the classroom  
8 dampness index was significantly associated with increased nasal symptoms (*P* for trend) in fully  
9 adjusted models. Because the significances of *P* for trend possibly indicated the dose-response of  
10 dampness exposure, those significances may reflect the effect of dampness on the symptoms.

11         A meta-analysis showed that home dampness was significantly associated with children's  
12 asthma and cough. [3] Home dampness reflects on evening and night exposure, and nocturnal cough  
13 is common in asthmatic patients. [30] The adverse effect of school dampness on respiratory  
14 symptoms has also been reported. [31] However, we saw no significant relationships between  
15 classroom dampness and cough. This may be attributed to shorter time in the school and/or  
16 inadequate statistical power.

17         Dampness relates to general symptoms in adults [1,13] and children. [31,32] In this study,  
18 only the home dampness index was significantly related to general symptoms. This also may be due



1 to the longer time in home and/or inadequate statistical power. Classroom dampness, not home  
2 dampness, was significantly related to nasal symptoms. Since nasal symptoms are the strongest just  
3 after waking up in the morning, [33] exposure to home dampness seems to be important. These  
4 nasal symptoms remain strong until noon; [33] therefore, the classroom dampness can possibly  
5 affect the late circadian peak of nasal symptoms.

6 As for the dampness index, the index consisting of condensation on windows, high air  
7 humidity in the bathroom, moldy odor, and water leakage had significantly increased ORs for SBS  
8 symptoms in Swedish multiresident houses. [13] In Taipei office buildings, the dampness index had  
9 significantly increased ORs for eye irritation, cough, and lethargy/fatigue. [34] Moreover, several  
10 studies of Japanese dwellings reported that the dampness index had significantly increased ORs for  
11 SBS symptoms. [15-18]

12 As previously mentioned, it has been reported that lifestyle can affect SBS symptoms. [20]  
13 In this study, usually not eating breakfast had significantly higher ORs for eye symptoms, and too  
14 many snacks and/or sweets had higher ORs for eye, nose, general symptoms, and any symptoms.  
15 Thus, favorable food habits probably have a positive effect on subjective symptoms. However, there  
16 is a possibility that favorable food habits are related to lower ORs for subjective symptoms via other  
17 favorable lifestyle factors.

18 This study had several limitations. First, if participants knew that dampness was a risk factor

1 for the subjective symptoms, people whose houses had dampness problems might tend to report  
2 subjective symptoms, which might could have caused a bias. However, exposure to chemicals has  
3 been spotlighted in Japan, and many Japanese may prefer higher humidity in winter. Besides,  
4 classroom dampness was estimated by classroom teachers who did not know pupils' symptoms.  
5 Therefore, such bias seems to have occurred rarely in this study. If nonrespondents were prone to  
6 have no subjective symptoms, the symptom prevalence rate may have been overestimated. Secondly,  
7 a lower response rate may also affect the significant relationship between dampness and subjective  
8 symptoms because the statistical power was reduced. Thus, in each dampness indicator, only water  
9 leakage of dwellings had significantly higher ORs for cough and general symptoms in multivariate  
10 adjustment analyses. Secondly, in fully adjusted analyses, *P* for trends of dampness index was not  
11 necessarily consistent with *P* of each dampness index number (OR of home dampness index for  
12 general symptoms; OR of class dampness index for nasal symptoms). This may also have been  
13 caused due to reduced statistical power. A seasonal variation of SBS symptoms—an increase in  
14 mucosal and skin symptoms during winter and spring—has been reported. [35] Fourthly, the season  
15 of this study may have affected the prevalence of subjective symptoms. Regarding the symptom  
16 part of the MM040 questionnaire, several studies have defined positive symptoms as those  
17 occurring often, but “whether the symptoms were attributed to the work environment or not” was  
18 not included in the definition of SBS symptoms. [13,23] Fifthly, the MM080 includes queries about

1 “whether the symptoms were attributed to the school environment or not.” Because our study  
2 focused on both the home and school environment, “whether the symptoms were attributed to the  
3 school environment or not” was not used in our subjective symptom definition. The Japanese  
4 version of MM080 was not validated by another Japanese study. However, the queries of subjective  
5 symptoms contain common SBS symptoms that are included in the MM040. Therefore, we believe  
6 that we were able to estimate the SBS symptoms properly. Finally, This study was conducted in two  
7 cities of Hokkaido, Japan. Therefore, generalization of this study to other districts in Japan and in  
8 other countries is difficult. But many studies have reported the effect of dampness on pupils’ health.

9 In conclusion, both home and classroom dampness can affect pupils’ health. Home dampness,  
10 in particular, was significantly related to cough and general symptoms, and classroom dampness  
11 was significantly related to nose symptoms. Furthermore, favorable food habits have a positive  
12 effect on pupils’ subjective symptoms. Thus, to improve both home and school dampness, pupils’  
13 food habits is one of the measures that can be used to protect pupils’ health, especially in the case of  
14 SBS. However, further studies are required to investigate the dampness environment and lifestyle  
15 factors for improving pupils’ health.

16

1 **Acknowledgment**

2 This work was supported by a grant from Foundation for Total Health Promotion and a Health  
3 Science Research Grant from the Japan Ministry of Health, Labor and Welfare.

4

1 **References**

2

3 1. Bornehag CG, Blomquist G, Gyntelberg F, Jarvholm B, Malmberg P, Nordvall L, et al. Dampness  
4 in buildings and health. Nordic interdisciplinary review of the scientific evidence on  
5 associations between exposure to "dampness" in buildings and health effects  
6 (NORDDAMP). *Indoor Air*. 2001;11:72-86.

7 2. Bornehag CG, Sundell J, Bonini S, Custovic A, Malmberg P, Skerfving S, et al. Dampness in  
8 buildings as a risk factor for health effects, EUROEXPO: a multidisciplinary review of the  
9 literature (1998-2000) on dampness and mite exposure in buildings and health effects.  
10 *Indoor Air*. 2004;14:243-257.

11 3. Fisk WJ, Lei-Gomez Q, Mendell MJ. Meta-analyses of the associations of respiratory health  
12 effects with dampness and mold in homes. *Indoor Air*. 2007;17:284-296.

13 4. Rylander R, Norrhall M, Engdahl U, Tunsater A, Holt PG. Airways inflammation, atopy, and  
14 (1--> 3)-beta-D-glucan exposures in two schools. *Am J Respir Crit Care Med*.  
15 1998;158:1685-1687.

16 5. Savilahti R, Uitti J, Roto P, Laippala P, Husman T. Increased prevalence of atopy among children  
17 exposed to mold in a school building. *Allergy*. 2001;56:175-179.

18 6. Smedje G, Norback D, Edling C. Asthma among secondary schoolchildren in relation to the

- 1 school environment. *Clin Exp Allergy*. 1997;27:1270-1278.
- 2 7. Taskinen T, Hyvarinen A, Meklin T, Husman T, Nevalainen A, Korppi M. Asthma and respiratory  
3 infections in school children with special reference to moisture and mold problems in the  
4 school. *Acta Paediatr*. 1999;88:1373-1379.
- 5 8. Meklin T, Potus T, Pekkanen J, Hyvarinen A, Hirvonen MR, Nevalainen A. Effects of  
6 moisture-damage repairs on microbial exposure and symptoms in schoolchildren. *Indoor Air*.  
7 2005;15 Suppl 10:40-47.
- 8 9. Bornehag CG, Sundell J, Hagerhed-Engman L, Sigsggard T, Janson S, Aberg N. 'Dampness' at  
9 home and its association with airway, nose, and skin symptoms among 10,851 preschool  
10 children in Sweden: a cross-sectional study. *Indoor Air*. 2005;15 Suppl 10:48-55.
- 11 10. Haverinen-Shaughnessy U, Pekkanen J, Hyvarinen A, Nevalainen A, Putus T, Korppi M, et al.  
12 Children's homes--determinants of moisture damage and asthma in Finnish residences.  
13 *Indoor Air*. 2006;16:248-255.
- 14 11. Simoni M, Lombardi E, Berti G, Rusconi F, La Grutta S, Piffer S, et al. Mould/dampness  
15 exposure at home is associated with respiratory disorders in Italian children and adolescents:  
16 the SIDRIA-2 Study. *Occup Environ Med*. 2005;62:616-622.
- 17 12. Kerckmar CM, Dearborn DG, Schluchter M, Xue L, Kirchner HL, Sobolewski J, et al.  
18 Reduction in asthma morbidity in children as a result of home remediation aimed at

- 1 moisture sources. *Environ Health Perspect.* 2006;114:1574-1580.
- 2 13. Engvall K, Norrby C, Norback D. Sick building syndrome in relation to building dampness in  
3 multi-family residential buildings in Stockholm. *Int Arch Occup Environ Health.*  
4 2001;74:270-278.
- 5 14. Engvall K, Norrby C, Norback D. Asthma symptoms in relation to building dampness and odour  
6 in older multifamily houses in Stockholm. *Int J Tuberc Lung Dis.* 2001;5:468-477.
- 7 15. Kishi R, Saijo Y, Kanazawa A, Tanaka M, Yoshimura T, Chikara H, et al. Regional differences  
8 in residential environments and the association of dwellings and residential factors with the  
9 sick house syndrome: a nationwide cross-sectional questionnaire study in Japan. *Indoor Air.*  
10 2009;243-254.
- 11 16. Saijo Y, Kishi R, Sata F, Katakura Y, Urashima Y, Hatakeyama A, et al. Symptoms in relation to  
12 chemicals and dampness in newly built dwellings. *Int Arch Occup Environ Health.*  
13 2004;77:461-470.
- 14 17. Takeda M, Saijo Y, Yuasa M, Kanazawa A, Araki A, Kishi R. Relationship between sick  
15 building syndrome and indoor environmental factors in newly built Japanese dwellings. *Int*  
16 *Arch Occup Environ Health.* 2009;82:583-593.
- 17 18. Saijo Y, Nakagi Y, Ito T, Sugioka Y, Endo H, Yoshida T. Relation of dampness to sick building  
18 syndrome in Japanese public apartment houses. *Environ Health Prev Med.* 2009;14:26-35.

- 1 19. Runeson R, Wahlstedt K, Wieslander G, Norback D. Personal and psychosocial factors and  
2 symptoms compatible with sick building syndrome in the Swedish workforce. *Indoor Air*.  
3 2006;16:445-453.
- 4 20. Nakayama K, Morimoto K. Relationship between, lifestyle, mold and sick building syndromes  
5 in newly built dwellings in Japan. *Int J Immunopathol Pharmacol*. 2007;20:35-43.
- 6 21. Fu ML, Cheng L, Tu SH, Pan WH. Association between unhealthy eating patterns and  
7 unfavorable overall school performance in children. *J Am Diet Assoc*. 2007;107:1935-1943.
- 8 22. Andersson K, Aslaksen W. Can the parents give a reliable assessment of  
9 the children's school environment? *Indoor Air*. 2005;15 issue s11:214.
- 10 23. Mizoue T, Reijula K, Andersson K. Environmental tobacco smoke exposure and overtime work  
11 as risk factors for sick building syndrome in Japan. *Am J Epidemiol*. 2001;154:803-808.
- 12 24. Saeki H, Iizuka H, Mori Y, Akasaka T, Takagi H, Kitajima Y, et al. Prevalence of atopic  
13 dermatitis in Japanese elementary schoolchildren. *Br J Dermatol*. 2005;152:110-114.
- 14 25. Nishima S, Chisaka H, Fujiwara T, Furusho K, Hayashi S, Hiraba K, et al. Surveys on the  
15 Prevalence of Pediatric Bronchial Asthma in Japan: A Comparison between the 1982, 1992,  
16 and 2002 Surveys Conducted in the Same Region Using the Same Methodology. *Allergol Int*.  
17 2009;58:37-53.
- 18 26. Ozasa K, Hama T, Dejima K, Watanabe Y, Hyo S, Terada T, et al. A 13-year study of Japanese



- 1 cedar pollinosis in Japanese schoolchildren. *Allergol Int.* 2008;57:175-180.
- 2 27. Postma DS. Gender differences in asthma development and progression. *Gend Med.* 2007;4  
3 Suppl B:S133-146.
- 4 28. Hayashi T. A study on the causes of an epidemic of influenza, especially an analysis of relative  
5 humidity as a main cause. *Nippon Ika Daigaku Zasshi.* 1985;52:272-280 (in Japanese).
- 6 29. Matsumoto I, Yoshida S, Kawana R. Virological surveillance of acute respiratory tract illnesses  
7 of children in Morioka, Japan. II. Rhinovirus infection. *Kansenshogaku Zasshi.*  
8 1991;65:1286-1296 (in Japanese).
- 9 30. Brooke AM, Lambert PC, Burton PR, Clarke C, Luyt DK, Simpson H. Night cough in a  
10 population-based sample of children: characteristics, relation to symptoms and associations  
11 with measures of asthma severity. *Eur Respir J.* 1996;9:65-71.
- 12 31. Mudarri D, Fisk WJ. Public health and economic impact of dampness and mold. *Indoor Air.*  
13 2007;17:226-235.
- 14 32. Meyer HW, Wurtz H, Suadicani P, Valbjorn O, Sigsgaard T, Gyntelberg F, et al. Molds in floor  
15 dust and building-related symptoms in adolescent school children. *Indoor Air.*  
16 2004;14:65-72.
- 17 33. Smolensky MH, Reinberg A, Labrecque G. Twenty-four hour pattern in symptom intensity of  
18 viral and allergic rhinitis: treatment implications. *J Allergy Clin Immunol.*

1           1995;95:1084-1096.

2   34. Wan GH, Li CS. Dampness and airway inflammation and systemic symptoms in office building  
3           workers. Arch Environ Health. 1999;54:58-63.

4   35. Mizoue T, Andersson K, Reijula K, Fedeli C. Seasonal variation in perceived indoor  
5           environment and nonspecific symptoms in a temperate climate. J Occup Health.  
6           2004;46:303-309.

7

8

Table 1 Characteristics of Pupils and their Homes (n = 1,077)

	n	%
Grade		
1 <sup>st</sup>	190	17.6
2 <sup>nd</sup>	165	15.3
3 <sup>rd</sup>	198	18.4
4 <sup>th</sup>	156	14.5
5 <sup>th</sup>	179	16.6
6 <sup>th</sup>	189	17.5
Sex		
Boys	516	47.9
Girls	561	52.1
Allergy within 1 yr	379	35.2
Type of housing		
Solitary house	791	73.4
Year of construction		
≤1980	250	23.2
>1980	730	67.8
Unknown	97	9.0
Heating		
Electric	129	12.0
Ventilation		
Natural	553	51.3
Mechanical	456	42.3
Unknown	68	6.3
Wall-to-wall carpeting in a child's room	523	48.6
Furry animals or birds in the home	263	24.4
Smoke indoors at home	557	51.7
Visible mold	312	29.0
Moldy odor	89	8.3
Water leakage	179	16.6
Condensation on windowpanes	595	55.2
Usually not eating breakfast (1–3 times per week, never)	28	2.6
No sufficient energy at school lunch	97	9.0
Too many snacks and/or sweets	245	22.7

Table 2 Characteristics of Schools (n = 8) and Classrooms (n = 61)

	School		Classroom	
	n	%	n	%
Year of construction				
>1995	3	37.5	24	39.3
<1987	5	62.5	37	60.7
Ventilation system				
Natural	4	50	29	47.5
Mechanical	4	50	32	52.5
Moldy odor	3	37.5	9	14.8
Visible mold	2	25	9	14.8
Water leakage	3	37.5	7	11.5
Condensation on windowpanes	3	37.5	22	36.1
Number of pupils per class				
≤20			8	14.8
21–30			23	37.7
≥31			29	47.5

Table 3 Prevalence of Subjective Symptoms\* in Boys and Girls

	Boys (n = 516)		Girls (n = 561)		All pupils (n = 1077)		<i>P</i> (Boys vs. girls)
	n	%	n	%	n	%	
Eye	22	4.3	25	4.5	47	4.4	0.877
Nose	87	16.9	64	11.4	151	14.0	0.010
Cough	29	5.6	23	4.1	52	4.8	0.245
Skin	57	11.0	65	11.6	122	11.3	0.780
General	22	4.3	28	5.0	52	4.8	0.795
Any**	144	27.9	127	22.6	271	25.2	0.047

\* occurred often (weekly)

\*\* having at least one of above symptoms

Table 4 Multivariate Adjusted Odds Ratios of Dampness and Food Habits for Subjective Symptoms

	Eye		Nose		Cough		Skin		General		Any	
	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>
Usually not eating breakfast	3.84 (1.22–12.5)	0.022	1.72 (0.68–1.34)	0.246	0.66 (0.08–5.00)	0.687	0.75 (0.21–2.70)	0.653	2.70 (0.85–8.33)	0.093	1.69 (0.75–3.85)	0.208
No sufficient energy following school lunch	1.17 (0.44–3.10)	0.755	1.69 (0.97–2.95)	0.065	1.86 (0.83–4.17)	0.134	1.59 (0.86–2.91)	0.137	0.75 (0.26–2.14)	0.586	1.28 (0.79–2.07)	0.310
Too many snacks and/or sweets	2.56 (1.38–4.75)	0.003	1.64 (1.09–2.45)	0.017	1.31 (0.69–2.48)	0.405	0.99 (0.62–1.58)	0.973	2.63 (1.46–4.76)	0.001	1.49 (1.07–2.08)	0.018
Home												
Visible mold	1.28 (0.66–2.48)	0.465	1.18 (0.79–1.77)	0.427	1.60 (0.87–2.95)	0.129	0.94 (0.60–1.47)	0.771	1.65 (0.90–3.03)	0.106	1.20 (0.87–1.66)	0.270
Moldy odor	1.06 (0.35–3.28)	0.915	1.00 (0.52–1.94)	0.998	1.78 (0.75–4.22)	0.192	0.97 (0.47–2.04)	0.943	2.05 (0.90–4.64)	0.086	1.37 (0.82–2.30)	0.229
Water leakage	0.79 (0.32–1.94)	0.604	1.28 (0.79–2.06)	0.315	2.54 (1.34–4.83)	0.004	1.19 (0.71–1.99)	0.506	2.42 (1.27–4.64)	0.008	1.42 (0.97–2.07)	0.073
Condensation on windowpanes	1.02 (0.52–2.00)	0.967	1.18 (0.78–1.78)	0.428	1.53 (0.77–3.03)	0.226	1.16 (0.74–1.81)	0.526	1.15 (0.59–2.23)	0.675	1.30 (0.94–1.81)	0.117
Dampness Index												
0	Reference		Reference		Reference		Reference		Reference		Reference	
1	0.63 (0.27–1.47)	0.283	0.96 (0.59–1.58)	0.881	0.92 (0.39–2.15)	0.847	1.13 (0.67–1.90)	0.653	0.63 (0.26–1.53)	0.309	1.12 (0.76–1.65)	0.577
2	1.11 (0.49–2.50)	0.807	1.33 (0.80–2.20)	0.279	1.08 (0.44–2.64)	0.871	1.08 (0.61–1.93)	0.787	1.18 (0.51–2.73)	0.697	1.32 (0.87–2.01)	0.189
3	0.92 (0.30–2.81)	0.807	1.22 (0.62–2.39)	0.567	2.74 (1.09–6.86)	0.032	0.90 (0.42–1.93)	0.782	1.73 (0.70–4.48)	0.258	1.27 (0.74–2.19)	0.390
4	1.03 (0.12–8.70)	0.982	1.35 (0.42–4.42)	0.615	4.94 (1.32–18.5)	0.018	1.98 (0.64–6.14)	0.238	4.82 (1.40–16.6)	0.012	3.20 (1.36–7.54)	0.008
<i>P</i> for trend	0.853		0.281		0.008		0.702		0.017		0.030	

Classroom												
Visible mold	0.82 (0.30–2.26)	0.705	1.53 (0.87–2.68)	0.140	1.20 (0.48–2.98)	0.698	1.17 (0.62–2.21)	0.628	0.63 (0.25–1.64)	0.357	1.39 (0.88–2.20)	0.155
Moldy odor	1.35 (0.55–3.32)	0.520	1.46 (0.87–2.45)	0.148	1.63 (0.68–3.91)	0.275	1.82 (1.00–3.32)	0.052	0.58 (0.22–1.53)	0.272	1.32 (0.86–2.02)	0.211
Water leakage	0.55 (0.18–1.73)	0.309	1.24 (0.71–2.16)	0.442	1.42 (0.61–3.28)	0.414	1.02 (0.55–1.90)	0.940	0.24 (0.06–1.03)	0.055	0.94 (0.60–1.48)	0.790
Condensation on windowpanes	0.77 (0.28–2.13)	0.618	1.72 (0.90–3.29)	0.102	1.79 (0.61–5.30)	0.292	0.93 (0.48–1.82)	0.839	0.47 (0.19–1.17)	0.104	0.92 (0.57–1.48)	0.727
Dampness index												
0	Reference		Reference		Reference		Reference		Reference		Reference	
1	0.58 (0.17–1.98)	0.381	1.71 (0.90–3.25)	0.101	1.39 (0.48–4.05)	0.544	0.72 (0.34–1.51)	0.379	0.42 (0.13–1.33)	0.139	0.74 (0.44–1.26)	0.272
2	0.85 (0.25–2.89)	0.799	1.34 (0.65–2.77)	0.424	1.40 (0.44–4.46)	0.573	0.83 (0.37–1.86)	0.641	0.64 (0.23–1.74)	0.378	0.92 (0.53–1.61)	0.768
3	1.31 (0.32–5.30)	0.704	2.03 (0.86–4.80)	0.107	1.15 (0.25–5.28)	0.859	1.42 (0.56–3.61)	0.463	0.27 (0.05–1.49)	0.133	0.99 (0.49–2.00)	0.969
4	0.51 (0.09–2.81)	0.437	2.30 (0.99–5.38)	0.054	2.33 (0.67–8.14)	0.184	1.51 (0.61–3.74)	0.376	0.16 (0.02–1.41)	0.099	1.42 (0.72–2.77)	0.312
<i>P</i> for trend	0.675		0.050		0.244		0.401		0.042		0.443	

<sup>a</sup>Adjusted for age, sex, allergy, construction year of dwelling, housing type (solitary or other), ventilation of dwelling, type of heating of dwelling, construction year of school, ventilation of school, and number of pupils per class. Each dampness indicator and food habit was introduced separately in the model.

Table 5 Fully Adjusted Odds Ratios of Dampness and Food Habits for Subjective Symptoms

	Eye		Nose		Cough		Skin		General		Any	
	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>	OR <sup>a</sup> 95% CI	<i>P</i>
Usually not eating breakfast	3.7 (1.11–12.5)	0.033	1.89 (0.72–4.76)	0.200	0.61 (0.07–5.00)	0.642	0.71 (0.19–2.63)	0.593	2.12 (0.63–7.14)	0.223	1.59 (0.68–3.70)	0.286
No sufficient energy following school lunch	1.25 (0.47–3.38)	0.656	1.6 (0.91–2.81)	0.102	1.78 (0.78–4.06)	0.171	1.59 (0.86–2.94)	0.141	0.71 (0.24–2.11)	0.534	1.26 (0.77–2.05)	0.357
Too many snacks and/or sweets	2.43 (1.29–4.57)	0.006	1.57 (1.04–2.37)	0.034	1.28 (0.67–2.47)	0.454	0.96 (0.60–1.54)	0.865	2.45 (1.33–4.53)	0.004	1.4 (1.00–1.97)	0.051
Home												
Dampness index												
0	Reference		Reference		Reference		Reference		Reference		Reference	
1	0.6 (0.25–1.43)	0.249	0.96 (0.59–1.59)	0.874	0.93 (0.40–2.19)	0.867	1.15 (0.68–1.95)	0.593	0.62 (0.26–1.52)	0.298	1.12 (0.76–1.66)	0.578
2	1.01 (0.44–2.33)	0.988	1.31 (0.78–2.20)	0.316	1.04 (0.42–2.55)	0.934	1.09 (0.61–1.96)	0.778	0.93 (0.40–2.19)	0.871	1.27 (0.83–1.94)	0.278
3	0.95 (0.31–2.94)	0.927	1.27 (0.64–2.51)	0.498	2.8 (1.11–7.07)	0.029	0.92 (0.43–1.99)	0.833	1.65 (0.63–4.30)	0.309	1.29 (0.75–2.23)	0.365
4	0.67 (0.07–3.02)	0.72	1.3 (0.39–4.31)	0.67	5.43 (1.41–20.9)	0.014	2.08 (0.65–6.60)	0.215	3.57 (1.01–12.5)	0.047	2.92 (1.22–6.99)	0.016
<i>P</i> for trend	0.645		0.184		0.007		0.566		0.056		0.047	
Classroom												
Dampness Index												
0	Reference		Reference		Reference		Reference		Reference		Reference	
1	0.6 (0.17–2.11)	0.419	1.82 (0.95–3.49)	0.074	1.5 (0.50–4.48)	0.472	0.71 (0.34–1.50)	0.368	0.49 (0.15–1.62)	0.243	0.79 (0.46–1.34)	0.379
2	0.93 (0.27–3.21)	0.903	1.41 (0.68–2.92)	0.354	1.46 (0.45–4.77)	0.53	0.81 (0.36–1.83)	0.61	0.74 (0.26–2.08)	0.565	0.96 (0.55–1.70)	0.898



3	1.65 (0.39–6.92)	0.496	2.28 (0.95–5.44)	0.063	1.26 (0.27–5.94)	0.771	1.39 (0.54–3.55)	0.497	0.33 (0.06–1.85)	0.207	1.07 (0.53–2.19)	0.848
4	0.51 (0.09–2.98)	0.451	2.38 (1.00–5.66)	0.051	2.43 (0.67–8.85)	0.179	1.41 (0.56–3.54)	0.463	0.19 (0.02–1.70)	0.139	1.48 (0.75–2.94)	0.258
<i>P</i> for trend	0.763		0.04		0.232		0.469		0.09		0.348	

<sup>a</sup>Adjusted for age, sex, allergy, construction year of dwelling, housing type (solitary or other), ventilation of dwelling, type of heating of dwelling, construction year of school, ventilation of school, and number of pupils per class.

All above variables included in the model.