


FERTILIZER RECOMMENDATIONS

for the High Plains



Adapted from Texas Agricultural Experiment Station Bulletin 431, by W. T. Carter.

TEXAS AGRICULTURAL EXTENSION SERVICE
 J. E. Hutchison, Director, College Station, Texas

FERTILIZER RECOMMENDATIONS

for the High Plains

William F. Bennett, Extension Soil Chemist
Chas. E. Fisher, Superintendent, Substation No. 8
Texas A. & M. College System

The fertilizer recommendations contained herein are general in scope. Soils vary in fields and area so much that soil tests should be made in order to recommend more definite and economical fertilizer applications.

For best results with fertilizers, other factors should be favorable, such as a well-prepared seed bed, good stand, absence of disease, adequate moisture, aeration and good cultural practices. Good cropping systems with legumes in rotation aid in a favorable response of crops to fertilizers. Where soil conditions are very favorable even higher rates of fertilization than those shown may be economically advantageous.

High-analysis fertilizers usually are cheaper. Low-analysis fertilizers cost less per bag, but the cost per acre is greater for the same amount of nutrients. The grades, 5-10-5 and 10-20-10, have the same ratio (1-2-1) of nutrients, but 10-20-10 has twice as much fertilizing value as 5-10-5. It requires only one-half as many pounds of 10-20-10 per acre to supply the same amount of plant nutrients.

Fertilizers containing phosphorus should be drilled or plowed into the land. Phosphorus does not move freely into the soil. Liquid fertilizers may be used instead of solid fertilizers at the same rate per acre. Liquid fertilizers usually are much more expensive per unit of nutrients.

Anhydrous ammonia is an excellent source of nitrogen when properly applied. It is the cheapest nitrogen fertilizer when used at moderate to high rates on large acreages. When anhydrous ammonia is used as a fertilizer, the opening made by the applicator should be covered immediately to prevent loss of ammonia. Likewise, the soil should be in good tilth.

FERTILIZER RECOMMENDATIONS:

The following recommendations are expressed in pounds of nutrients per acre and do not represent fertilizer grades. The nutrients must be obtained from materials or fertilizer mixtures sold on the market.

For example, a recommendation calling for 30-60-30, which is a 1-2-1 ratio, can be obtained by applying 600 pounds of 5-10-5 or 250 pounds of 12-24-12 or 300 pounds of 10-20-10. Again, if a recommendation calls for 15-60-0, this may be obtained by applying about 400 pounds of a 4-16-0 or 125 pounds of 11-48-0.

Row Crops: Fertilizer usually is applied at the time of planting or just before. Fertilizers are more efficiently used by most crops when applied in a band 10 inches to the side and 5 inches below the seed.

If equipment for applying fertilizers in bands while planting or cultivating is not available, apply the fertilizer in the water furrow and bed on it when the land is prepared for planting. Avoid putting the seed too close to the fertilizer as germination may be impaired.

Small Grains: Fertilizers for small grains may be broadcast, drilled in or plowed in. Fertilizers containing nitrogen and potash should not be allowed to touch the seed.

Phosphorus, potash and part of the nitrogen should be applied at or before seeding. The rest of the nitrogen should be applied in the spring before plants begin to joint.

Pastures: For establishing improved pastures, fertilizer should be applied in bands when possible. Otherwise, it should be broadcast, drilled or plowed in. For maintenance, topdress with 30-0-0 as needed. Repeat basic fertilizer treatment annually as suggested or according to a soil test.

Recommendations for fertilizers in this circular are those found best by experiments, tests and practical experience in the field. They range from the calcareous (limy) river valley clays to the sands of the uplands. If your farm contains both clays and loams, there will be two recommendations for your land.

When crops follow legumes turned under, the amount of fertilizer to be applied at planting may be reduced. Side or topdress with the amount of fertilizer suggested.

The letters NR mean that the crop is not recommended for this class of soils.

High Plains

(Irrigated Land)

POUNDS OF NUTRIENTS TO BE APPLIED PER ACRE AT OR BEFORE PLANTING

	Clays and clay loams	Loams and sandy loams	Sands and loamy sands	Additional Treatment
FIELD CROPS				
Alfalfa & biennial sweetclovers	20-60-0	20-60-0	30-60-30	For maintenance 0-60-0 annually
Corn Grain sorghum	90-0-0	90-30-0	30-60-30	Sidedress sands and loamy sands with 30-0-0 in 35 days after planting if sufficient water is applied
Sorghum for hay Sudan	30-0-0	30-30-0	30-60-30	Sidedress with 30-0-0 after each cut- ting
Cotton, sesame	30-30-0	60-60-0	60-60-30	
Annual legumes	0-30-0	0-40-0	15-60-30	
Oats and wheat	60-0-0	60-30-0	30-60-30	Topdress sands and loamy sands with 40-0-0 in January or early Feb- ruary if irrigation is adequate
Oats and wheat to be grazed	60-0-0	60-30-30	60-60-30	Topdress with 40-0-0 in January or early February
Pastures (cultivated) Grasses only	20-0-0	20-60-0	30-60-30	Topdress with 30-0-0 each time grazed down or cut
Sugar beets, stock beets	30-0-0	30-60-0	30-60-30	Sidedress with 60-0-0
TRUCK CROPS				
Peas, black-eye, etc.	0-60-0	20-60-0	20-60-30	
Beans	20-40-0	40-80-0	40-80-40	
Cabbage, lettuce, mustard, etc.	30-30-0	40-80-0	40-80-40	Sidedress with 60-0-0 when plants begin to bud
Cucumbers, squash	30-30-0	40-80-0	40-80-40	Sidedress with 30-0-0 when vines begin to bloom
Cantaloupes	30-30-0	40-80-0	40-80-40	
Carrots, beets, turnips	0-30-0	30-60-0	30-60-30	
Irish potatoes	60-0-0	60-60-30	80-80-30	
Onions	40-40-0	40-80-0	40-80-40	
Sweet potatoes	NR	30-60-30	30-60-60	
Tomatoes, peppers	40-40-0	50-100-0	50-100-50	
Watermelons	20-40-0	30-60-0	30-60-30	

Fertilizers not recommended on dryland. Irrigated land probably will not respond to fertilizer for the first 2 years it is irrigated.