

The bimodal regulation of vascular function by superoxide anion: role of endothelium

Buket Demirci, MD¹; Pascal P. McKeown, MD¹; Ulvi Bayraktutan DVM, PhD²

¹Department of Medicine and Therapeutics, Institute of Clinical Science, Queen's University Belfast, Belfast and ²Division of Stroke Medicine, Clinical Sciences Building, University of Nottingham, Nottingham, UK

Short Title: superoxide anion regulates vasorelaxation

Address for correspondence

Ulvi Bayraktutan DVM, PhD

Associate Professor

Division of Stroke Medicine,

Clinical Sciences Building,

The University of Nottingham,

Hucknall Road,

Nottingham NG5 1PB

United Kingdom

Tel: +44 115 8231764

Fax: +44 115 8231767

E-mail: ulvi.bayraktutan@nottingham.ac.uk

Word count of the abstract: 200

Word count of the body: 4548 including title page, abstract and references.

Abstract

Reactive oxygen species (ROS) are implicated in vascular homeostasis. This study investigated whether $O_2^{\cdot-}$, the foundation molecule of all ROS, regulates vasomotor function. Vascular reactivity was measured by isometric tension studies using thoracic aortas from 12-14 week-old Sprague Dawley rats in the absence or presence of pyrogallol, an $O_2^{\cdot-}$ generator. Pyrogallol significantly enhanced intracellular $O_2^{\cdot-}$ level in a dose-dependent fashion without affecting the overall tissue viability. It also dose-dependently regulated both α -adrenergic agonist-mediated contractility to phenylephrine and endothelium-dependent relaxations to acetylcholine. Pyrogallol improved and attenuated responses to acetylcholine at its lower (10 nM - 1 μ M) and higher (10 - 100 μ M) concentrations, respectively while producing the inverse effects in response to phenylephrine at the same concentrations. The inactivation of nitric oxide synthase by L-NAME (50 μ M) abolished acetylcholine-induced vasodilatations but increased phenylephrine and KCl-induced vasoconstrictions irrespective of the concentration of the pyrogallol used. Relaxant responses to sodium nitroprusside, a nitric oxide donor, were not affected by pyrogallol. Other ROS i.e. peroxynitrite, H_2O_2 and hydroxyl radical that may be produced during experiments did not alter vascular functions. These findings suggest that the nature of $O_2^{\cdot-}$ -evoked vascular function is determined by its local concentration and the presence of a functional endothelium.

Keywords: Superoxide anion; Endothelium; Acetylcholine; Relaxation; Contractility

Introduction

The endothelium plays a pivotal role in the maintenance of normal vascular tone. However, its properties in pathological conditions change leading to a phenomenon termed “endothelial dysfunction” which is characterized by impaired endothelium-dependent relaxation (Bayraktutan, 2002). Endothelial dysfunction has been reported in conduit and resistance arteries of patients with vascular diseases including hypertension and in various animal models of human diseases (Panza et al., 1993; Taddei et al., 1993; Tesmaferiam et al., 1988; Luscher et al., 1986). Despite current data on the etiology of endothelial dysfunction being rather diverse, much of the recent data have implicated oxidative stress in its pathogenesis (Ohara et al., 1993; Kerr et al., 1999). Oxidative stress is associated with enhanced production of reactive oxygen species (ROS) in particular superoxide anion ($O_2^{\cdot-}$) that is normally converted to hydrogen peroxide (H_2O_2) by superoxide dismutases (SODs). H_2O_2 is in turn metabolized to H_2O by antioxidant enzymes; catalase and glutathione peroxidase (Yu, 1994). However, in pathological states the balance between the synthesis and metabolism of ROS may be impaired thus leading to generation of oxidative stress. $O_2^{\cdot-}$ can elicit direct vasoconstriction and scavenge nitric oxide (NO), the most potent endogenous vasodilator, to produce peroxynitrite ($OONO^{\cdot-}$) which oxidizes proteins, breaks DNA strands and reduces intracellular antioxidants such as glutathione and cysteine (Bayraktutan, 2002; Gryglewski et al., 1986). Contrary to these findings, other studies have revealed that low levels of $O_2^{\cdot-}$ are required for normal vascular cell proliferation and migration and contribute to endothelium-dependent vasodilatation in some vascular beds particularly after dismutation to H_2O_2 (Bayraktutan, 2004; Bayraktutan, 2005; Wei et al., 1996).

In the light of these contradictory findings, the aim of the current study was to investigate whether local concentrations of $O_2^{\cdot-}$ determine the nature of vascular functions using aortic rings from Sprague Dawley (SD) rats. To this end, relaxant responses to endothelium-

(in)dependent agents and contractile responses to an α -adrenergic agonist phenylephrine (PE) and receptor-independent constrictor KCl were investigated in vessels exposed to incremental concentrations of an $O_2^{\cdot-}$ -generating agent, pyrogallol. The requirement for an intact endothelium and the involvement of other ROS including H_2O_2 and $OONO^{\cdot-}$ in putative functional changes were also examined in the study.

Materials and methods

All the experiments described in this article were performed in accordance with the UK “Home Office Guidance on the Operation of the Animals (Scientific Procedures) Act 1986”.

Drugs and Chemicals

All drugs and chemicals, other than Mn(III)TMPyP (IDS Ltd., UK), used in this study were from Sigma Chemical Company (UK).

Animals

The studies were conducted using the thoracic aortas obtained from 12-14 week old male SD rats. The aortas were removed from anticoagulated rats (100 U heparin, IV) under deep pentobarbitone sodium anesthesia (100 mg/kg body, IP) and carefully cleaned of adhering tissue.

Vascular reactivity studies

Thoracic aortas were cut into six 2-3 mm rings and mounted horizontally on stainless hooks in 25 ml organ baths filled with carbonated (95% O_2 /5% CO_2) Krebs buffer [in (mM): NaCl 118.3, KCl 4.7, $MgSO_4$ 1.2, KH_2PO_4 1.22, $CaCl_2$ 2.5, $NaHCO_3$ 25, glucose 11.1, pH 7.4]. Rings were equilibrated for 90 min under 2 g resting tension and only one vasodilator was used for each ring experiment. Changes in isometric tension were detected and recorded by a force transducer and an 8 channel transducer data acquisition system (LE-TR201 and PowerLab/8S, ADInstruments), respectively. After initial equilibration period, ordinary Krebs

solution was replaced with fresh solution containing 1 μM indomethacin to dismiss the relaxant effects of prostanoids. All experiments were performed in the latter solution.

At the beginning of each experiment dose-response curves to PE (0.003-10 μM) were obtained. Rings were then washed and equilibrated prior to contraction with submaximal concentration of PE before detecting endothelium-dependent or endothelium-independent relaxant responses to acetylcholine (ACh, 0.03-1 μM) and sodium nitroprusside (SNP, 0.01-10 μM), respectively. For contractility studies, the rings were incubated with cumulative concentrations of PE (0.003-10 μM) or a single dose of KCl (120 mM) until the respective contractions reached a plateau. For pyrogallol experiments, rings that were used in the abovementioned experiments were washed and equilibrated prior to 20 min incubation with the required concentration of pyrogallol [$\text{C}_6\text{H}_6\text{O}_3$; 1,2,3-trihydroxybenzene] and repetition of the same experiments. This experimental pattern allowed each vascular ring to serve as its own control for a given chemical.

At the end of each experiment, the rings were blotted dry, weighed and the lengths were measured to calculate tension as normalized for cross-sectional area as previously reported (Abebe et al., 1990).

Detection of O_2^-

O_2^- generation in aortic rings was measured using three different approaches. Firstly, thoracic aortic rings (~10 mm) were placed in Krebs solution for 30 min at 37 °C prior to transfer into Krebs solution containing 5 μM of lucigenin. The chemiluminescence was recorded every 60 seconds for 10 min using a luminometer and the levels of O_2^- were calculated by taking into account the differences in the readings before and after addition of the aortic segments to the buffer (Bayraktutan et al; 1998).

Secondly, level of O_2^- was measured by SOD-inhibitable cytochrome C reduction assay. Briefly, aortic homogenates were incubated with 50 μM cytochrome C for 60 min at 37 °C in

Krebs buffer with or without SOD (125 U/ml). Absorbances were recorded using a GENios plate reader (TECAN) at 550 nm with 60 seconds intervals for 8 min. Production of $O_2^{\cdot-}$ was calculated as previously described and results were expressed as pmole $O_2^{\cdot-}$ per mg tissue (Horie and Kita; 1994).

Finally, the level of $O_2^{\cdot-}$ in aortic homogenates was measured using dihydroethidium (DHE) assay. DHE is oxidized by $O_2^{\cdot-}$ thereby yielding a fluorescent compound whose emission relates to the amount of $O_2^{\cdot-}$ present in the system. This assay was performed in 96-well microplates. Each well contained 100 μ l of homogenate obtained from aortas (~10 mm) exposed to different concentrations of pyrogallol (10 nM -100 μ M) in phosphate buffer (pH 7.8) plus 10 μ M of DHE in a volume of 100 μ l of phosphate buffer. The reaction was performed at 25 °C and the extent of DHE oxidation was followed by measuring the increase of the fluorescence using a microplate reader for 60 min (excitation= 485 nm and emission= 595 nm). Three independent experiments were performed for each concentration of pyrogallol. The percentage generation of fluorescence with aortic homogenates was calculated using the pyrogallol-untreated aortic homogenates as positive controls.

Detection of H_2O_2

H_2O_2 synthesis was measured using 2',7'-dichlorodihydrofluorescein (DCHF) diacetate (DA) as a fluorescent probe for intracellular H_2O_2 measurement (López-Ongil et al; 1998). DCHF-DA diffuses readily to the intracellular compartment, where it is desacetylated to the non-membrane-permeable DCHF. Then, during the cellular production of H_2O_2 , DCHF is oxidized and emits a fluorescent signal. To this end, the normal and pyrogallol-incubated aortic segments were homogenised and loaded with 20 μ M DCHF-DA for 30 min at 37 °C and fluorescence generation at 488 nm (excitation) and 525 nm (emission) was detected. The fluorescent signal was registered as a function of the time.

Detection of OONO⁻

Thoracic aortas were incubated with different concentrations of pyrogallol (10 nM- 100 μ M) in the presence of 20 μ M 123-dihydrorhodamine (123-DHR) for 2 h at 37 °C in Krebs solution. Following incubation, 123-DHR conversion to 123-rhodamine was measured by fluorometric analysis at excitation/emission wavelengths of 485 and 530 nm, respectively. Fluorescence due to autooxidation of 123-DHR was deducted from the original measurements (Muijsers et al; 2000).

LDH measurement

The vascular rings were exposed to different concentrations of pyrogallol in a 3 mL of phenol-free cell culture medium (DMEM) for 3 h at 37 °C in a humidified tissue culture incubator (5% CO₂ / 95% air) prior to removal of an 100 μ l of aliquot. The LDH release was measured spectrophotometrically from the changes in absorbance at 560 nm using 0.25 mM NADH and 0.75 mM pyruvate as substrates following subtraction of the background values obtained from the tubes without rings (Bergmeyer and Bernt; 1974).

Statistical Analysis

Data are expressed as mean \pm SEM. Each of the vascular reactivity was performed using 8 rats. Dose-response curves were fitted by non-linear regression with simplex algorithm and E_{max} values, the maximum contractile response of the tissue that indicates the potency, were calculated. Relaxant responses were given as the percentages of PE precontraction. Comparisons of dose-response curves were evaluated by two-way analysis of variance (ANOVA) for repeated measures. Molecular biological experiments were performed using aortic samples from 3 different rats. Statistical significances for LDH release and ROS production were assessed by two-way ANOVA followed by Bonferroni-Dunn's *post hoc* analysis. *P* values less than 0.05 were considered to be statistically significant.

Results

Intracellular levels of $O_2^{\cdot-}$ in aortic rings subjected to different levels of pyrogallol

In the current study, pyrogallol was employed as a generator of $O_2^{\cdot-}$ as previously described. Pyrogallol produced significant dose-dependent increases in the levels of aortic $O_2^{\cdot-}$ as detected by lucigenin enhanced-chemiluminescence, cytochrome C reduction and DHE oxidation assays without altering the level of LDH release, an indicator of cellular damage (Table 1).

Vasodilator responses in thoracic aortic rings and the effects of $O_2^{\cdot-}$ on vascular relaxation

In PE-precontracted aortic rings, both ACh and SNP produced concentration-dependent relaxations and the inactivation of endothelium by mechanical (by rubbing) or pharmacological (via incubation with 50 μ M of L-NAME) means selectively inhibited endothelium-dependent but not endothelium-independent vascular relaxations (Fig. 1A-C).

Pyrogallol regulated endothelium-dependent relaxant responses in aortic rings in a dose-dependent manner in that it improved and attenuated ACh-mediated relaxations at its lower (10 nM - 1 μ M) and higher (10 and 100 μ M) concentrations, respectively (Fig. 2A-B).

However, pyrogallol did not affect relaxant responses to SNP. It is of note that the pharmacological and mechanical inactivation of endothelium selectively abrogated ACh- but not SNP-mediated relaxant responses in the presence of pyrogallol (data not shown).

To investigate whether pyrogallol-induced effects are attributable to its $O_2^{\cdot-}$ -releasing function, the aortic rings were pretreated with one of the two structurally different $O_2^{\cdot-}$ scavengers, namely Tiron (4,5-dihydroxy-1,3-benzenedisulfonic acid, 1 μ M) or MPG (mercaptpropionylglycine, 1 μ M) for 20 min. Both agents prevented pyrogallol-mediated (10 μ M) endothelium-dependent relaxations without affecting the basal relaxations determined in the absence of pyrogallol (Fig. 3A-B). To support these findings, experiments with exogenous SOD (150 U/ml) or an endogenous inhibitor of SOD, namely diethyldithiocarbamate

(DETCA, 1-10 mM) were conducted. Pretreatments with either agent alone for 20 min failed to alter the ACh-mediated relaxant responses while the combination of DETCA and 10 μ M pyrogallol worsened these responses and the SOD and pyrogallol combination was found to be ineffective (Fig. 3C-D). To establish whether the effects of low concentrations of pyrogallol are also mediated by $O_2^{\cdot-}$, similar experiments were conducted in the presence of 10 nM of pyrogallol where Tiron and MPG suppressed low dose pyrogallol-induced increases in ACh-mediated relaxant responses by almost 22% while DETCA and exogenous SOD failed to alter these responses in a significant manner.

Experiments performed under identical conditions with Tiron, MPG, SOD or DETCA revealed that none of these treatments had an impact on endothelium-independent relaxant responses.

Elucidation of the putative involvement of other ROS in $O_2^{\cdot-}$ -mediated vascular relaxations

To examine the relative contributions of hydroxyl radical (\cdot OH), H_2O_2 or $OONO^{\cdot-}$ that may be generated during incubation with pyrogallol to relaxant changes, the aortic rings were treated with two different inhibitors of each agent in the absence or presence of a lower (100 nM) or a higher (10 μ M) concentration of pyrogallol. To this end, desferrioxamine (100 μ M) or DMSO (50 nM), catalase or glutathione peroxidase (100 U/ml for each agent) and Mn(III)TMPyP (10 μ M) or uric acid (100 μ M) were used as inhibitors of \cdot OH, H_2O_2 and $OONO^{\cdot-}$. It was revealed that none of these agents displayed significant effects on vasorelaxations. Figures 4A-C selectively show the effects of DMSO, catalase and Mn(III)TMPyP on ACh-mediated relaxations in the presence of 10 μ M of pyrogallol. To ensure whether pyrogallol treatments led to increases in intracellular levels of $OONO^{\cdot-}$ or H_2O_2 , the levels of these ROS were measured in aortic vessels which showed steady increases to incremental concentrations of pyrogallol (Table 2). Given that $OONO^{\cdot-}$ or H_2O_2 (10 nM - 100 μ M for each agent) *per se* produced dose-dependent increases in vascular relaxation in

vessels with or without endothelium (Fig. 5), it was safe to suggest that the pyrogallol-mediated increases in OONO^- or H_2O_2 did not reach the lower levels that were used in the latter experiments. Similar experiments with the listed inhibitors revealed their inefficacy in altering endothelium-independent relaxant responses to SNP.

Effects of O_2^- on vascular contractility

PE (0.003-10 μM) induced contractile responses in a dose-dependent manner in isolated aortic rings with or without endothelium (Fig. 6A). In endothelium-intact vascular rings, the lower concentrations of pyrogallol (10 nM - 1 μM) diminished PE-mediated contractile responses - significantly between 1 nM to 1 μM logarithmic concentrations of PE - without changing the E_{max} . However, the higher concentrations of pyrogallol (10 and 100 μM) significantly enhanced PE-mediated contractility (Fig. 6B). Pretreatment of vascular rings with L-NAME (50 μM) for 20 min significantly enhanced PE-mediated contractile responses regardless of the concentration of pyrogallol used (Fig. 6C). Similar to endothelium-intact rings, pyrogallol produced dose-dependent changes in endothelium-denuded aortic rings (data not shown).

The receptor-independent contractile responses of pyrogallol were determined through use of KCl (120 mM) that produced similar contractions in isolated aortic rings with or without endothelium in the absence or presence of pyrogallol (Fig. 7A-B). However, endothelial inactivation by L-NAME dramatically enhanced KCl-induced contractile responses in the presence of pyrogallol (Fig. 7C).

Elucidation of the putative involvement of other ROS in O_2^- -mediated vascular contractility

Tiron and MPG did not affect basal contractility to PE. However, they diminished pyrogallol-induced (10 μM) contractile responses to PE by almost 50% ($p < 0.05$). Inactivation of endothelium with L-NAME (50 μM) did not influence inhibitory effects of these components on pyrogallol-evoked contractions (Fig. 8). In contrast to O_2^- scavengers,

treatments with exogenous SOD, DETCA, catalase, DMSO and Mn(III)TMPyP in the presence of pyrogallol (100 nM or 10 μ M) did not significantly affect PE-mediated contractile responses and generated between 3 to 5% difference in these responses.

Discussion

Several factors including diminished production of NO due to inefficient utilization of substrate L-arginine or impaired activity of endothelial NO synthase have been implicated in the pathogenesis of endothelial dysfunction (Rodrigo et al., 1997; Liao et al., 1995; Goto et al., 2000). However, oxidative stress characterized by the excessive vascular presence of $O_2^{\cdot-}$, has recently been associated with the pathogenesis of this phenomenon in a variety of diseases with vascular complications such as hypertension and hypercholesterolemia (Luscher et al., 1986; Ohara et al., 1993). $O_2^{\cdot-}$ under pathological conditions may trigger the formation of other ROS such as $OONO^{\cdot-}$, $\cdot OH$ and H_2O_2 that exert opposing i.e. relaxant or contractile effects on endothelium through activating different pathways and/or modifying membrane permeability to macromolecules (Bayraktutan, 2002; Wei et al., 1996; Ellis et al., 2003; Gao et al., 2001; Shen et al., 2000; McQuaid et al., 1996).

Since the majority of the previous studies investigating the effects of ROS on vascular functions have focused mainly on exogenous H_2O_2 (Wei et al., 1996; Ellis et al., 2003; Gao et al., 2001; Shen et al., 2000; McQuaid et al., 1996; Lum et al., 2001), the current study aimed to investigate the direct effects of $O_2^{\cdot-}$, the foundation molecule of all ROS, on vasomotor functions using aortas from SD rats. This study primarily demonstrated that pyrogallol, a potent $O_2^{\cdot-}$ -generator, produced significant dose-dependent increases in intracellular levels of $O_2^{\cdot-}$ without affecting the tissue viability as determined by the measurements of LDH release. These findings were particularly important to ascertain that the putative differences that might be observed in the study would not be due to cellular damage. It also demonstrated that both ACh and SNP produced dose-dependent vasodilatations in aortic rings and that the

inactivation of endothelium by mechanical or pharmacological means selectively abolished relaxations to ACh thereby indicating the existence of a functional vascular smooth muscle layer following endothelial damage by either means. The investigation of the impact of O_2^- on vasomotor function revealed that the endothelium-derived vasodilations to ACh were improved and attenuated by lower (10 nM - 1 μ M) and higher (10 - 100 μ M) concentrations of pyrogallol, respectively and also that the inactivation of endothelium abrogated solely ACh-mediated responses regardless of the concentration of pyrogallol. Taken together these data implied a prerequisite for endothelial presence in modulation of beneficial activities of pyrogallol. This hypothesis was supported by the findings illustrating that the endothelial inactivation dramatically increased PE-induced contractile responses irrespective of the pyrogallol dose used despite initially recording significant decreases with lower concentrations (10 nM - 1 μ M) of pyrogallol in PE-evoked contractility.

It is noteworthy here that the combination of PE and pyrogallol generated similar effects in endothelium-intact and -denuded rings, findings were in agreement with previous ones suggesting that the smooth muscle cells or α 1-mediated responses may be the sites of action of O_2^- (Mizukawa et al., 1997; Gokce et al., 2005). Moreover, the additional contractility studies with KCl (120 mM) alone or with different doses of pyrogallol (10 nM - 100 μ M) produced similar contractions in vascular rings with or without endothelium. However, pretreatment of aortic rings with L-NAME significantly enhanced KCl-mediated contractile responses to pyrogallol selectively in endothelium-intact segments whereby indicated that the accentuation of pyrogallol-mediated contractility was due to endothelial inactivation but not to L-NAME itself. These data were rather surprising given that KCl induces smooth muscle contractions by depolarizing the cell membrane via a mechanism depending on calcium influx through the voltage-dependent channels (Karaki et al., 1997).

It has been well-documented that NO is involved in much of the endothelium-derived vasodilatation and is readily neutralized by $O_2^{\cdot-}$ (Bayraktutan, 2002; Gryglewski et al., 1986; Bellan et al., 1993). The direct correlation between $O_2^{\cdot-}$ and vascular reactivity was further assessed in this study using $O_2^{\cdot-}$ scavengers i.e. Tiron and MPG which prevented pyrogallol-induced impairments in ACh-mediated relaxant responses without altering the basal endothelium-dependent relaxations. Nonetheless, while the beneficial effects of $O_2^{\cdot-}$ scavengers were not mimicked by exogenous SOD, an irreversible inhibitor of endogenous SOD i.e. DETCA worsened relaxant responses when used in combination with pyrogallol. These findings suggest that an increase in endogenous $O_2^{\cdot-}$ stemmed from the inhibition of SOD alone is not sufficient to neutralize intracellular NO to deteriorate vascular relaxations.

It is inevitable that the incubation of aortic rings with pyrogallol would lead to generation of other ROS including $\cdot OH$, H_2O_2 and $OONO^{\cdot-}$. Indeed, exposure of aortas to incremental concentrations of pyrogallol produced steady increases in intracellular levels of both H_2O_2 and $OONO^{\cdot-}$ levels. To investigate the contributions of these free radicals to vascular relaxant and contractile responses, the relevant experiments were repeated in the presence of the specific inhibitors of $\cdot OH$ (DMSO; 50 nM or desferrioxamine; 100 μM), H_2O_2 (catalase or glutathione peroxidase; 100 U/ml for each agent) or $OONO^{\cdot-}$ (Mn(III)TMPyP; 10 μM or uric acid; 100 μM) that failed to alter the vascular responses when used alone or in the presence of pyrogallol (100 nM or 10 μM). In contrast, studies with $OONO^{\cdot-}$ and H_2O_2 on vasculature showed that both agents could produce relaxations in a dose-dependent manner in vessels with or without endothelium thereby indicating direct vasodilatory effects of these compounds on smooth muscle cells. In previous studies, $OONO^{\cdot-}$ -mediated relaxations have been assigned to elevation of cGMP levels, interference with Ca^{2+} movement and membrane hyperpolarization via K^+ -channel activation while those of H_2O_2 have been attributed to the NO synthesis and activation of smooth muscle K^+ channels (Li et al., 2005; Gil-Longo et al.,

2005). Although, the elucidation of the molecular mechanisms involved in $O_2^{\cdot-}$ -mediated vascular responses was out of the scope of the current study, it is possible that some of these mechanisms may be associated with $O_2^{\cdot-}$ -induced relaxant responses. In the current study, the inability of the ROS inhibitors to alter vascular responses may be explained by the existence of a delicate balance between their relaxant and contractile effects e.g. $OONO^{\cdot-}$, in addition to its vasodilator effects, also generates vascular contractions by reducing intracellular antioxidants including glutathione and oxidizing tetrahydrobiopterin, a critical cofactor for NO synthesis (Bayraktutan, 2002).

In summary, the major conclusions to be drawn from this study are firstly that the vascular relaxant and contractile functions are regulated by $O_2^{\cdot-}$ in a dose-dependent fashion. The similar bimodal regulation of vascular function has previously been reported with H_2O_2 in endothelium-intact and endothelium-denuded rat aortas (Gil-Longo and Gonzalez-Vazquez, 2005). Secondly, a functional endothelium is required for the regulation of $O_2^{\cdot-}$ -mediated dilator and contractile responses to ACh and PE, respectively. Finally, $O_2^{\cdot-}$ may exert a direct effect on smooth muscle cells via stimulation of α_1 -adrenergic receptors.

References

- Abebe, W., Harris, K.H. and MacLeod, K.M. (1991) Enhanced contractile responses of arteries from diabetic rats to α_1 -adrenoceptor stimulation in the absence and presence of extracellular calcium. *J. Cardiovasc. Pharmacol.* **16**, 239-248.
- Bayraktutan, U., Draper, N., Lang, D. and Shah, A.M. (1998) Expression of a functional neutrophil-type NADPH oxidase in cultured rat coronary microvascular endothelial cells. *Cardiovasc. Res.* **38**, 256-262.
- Bayraktutan, U. (2002). Free radicals, diabetes and endothelial dysfunction. *Diabetes Obes. Metab.* **4**, 224-238.

- Bayraktutan, U. (2004) Nitric Oxide Synthase and NAD(P)H Oxidase Enzymes Modulate Coronary Endothelial Cell Growth. *J. Mol. Cell. Cardiol.* **36**, 277-286.
- Bayraktutan, U. (2005) Coronary Microvascular Endothelial Cell Growth Regulates Expression of p22-phox Gene. *Free Rad. Biol. Med.* **39**, 1342-1352.
- Bellan, J.A., Longenecker, L.L., Kadowitz, P.J. and McNamara, D.B. (1993) Selective and complete blockade of acetylcholine-induced relaxation in rabbit aortic rings by Nw-Nitro-L-arginine but not glibenclamide. *Eur. J. Pharmacol.* **234**, 273-276.
- Bergmeyer, H. U. and Bernt, E. (1974) Lactate dehydrogenase UV-assay with pyruvate and NADH. In *Methods of Enzymatic Analysis*, pp. 574-578. Edited by H. U. Bergmeyer. New York, NY:Academic Press.
- Ellis, A., Pannirselvam, M., Anderson, T.J. and Triggle, C.R. (2003) Catalase has negligible inhibitory effects on endothelium-dependent relaxations in mouse isolated aorta and small mesenteric artery. *Br. J. Pharmacol.* **140**, 1193-1200.
- Gao, Y.J. and Lee, R.M. (2001) Hydrogen peroxide induces a greater contraction in mesenteric arteries of spontaneously hypertensive rats through thromboxane A₂ production. *Br. J. Pharmacol.* **134**, 1639-1646.
- Gil-Longo, J. and Gonzalez-Vazquez, C. (2005) Characterization of four different effects elicited by H₂O₂ in rat aorta. *Vascul. Pharmacol.* **43**, 128-138.
- Gokce, G. and Kerry, Z. (2005) Oxidative stress attenuates phenylephrine-induced contractile responses in rat aorta. *Hacettepe Uni. J. Fac. Pharmacy* **25**, 61-70.
- Goto, K., Fuji, K., Onaka, U., Abe, I. and Fujishama, M. (2000) ACE inhibitor prevents age-related endothelial dysfunction. *Hypertension* **36**, 581-587.
- Gryglewski, R.J., Palmer, R.M. and Moncada, S. (1986) O₂⁻ is involved in the breakdown of NO. *Nature* **320**, 454-456.

- Horie, S. and Kita, H. (1994) CD11b/CD18 is required for degranulation of human eosinophils induced by human recombinant granulocyte-macrophage colony-stimulating factor and platelet-activating factor. *J. Immunol.* **152**, 5457-5467.
- Karaki, H., Ozaki, H., Hori, M., Mitsui-Saito, M., Amano, K., Harada, K., Miyamoto, S., Nakazawa, H., Won, K.J. and Sato, K. (1997) Calcium movements, distribution, and function in smooth muscle. *Pharmacol. Rev.* **49**, 157-230.
- Kerr, S., Brosnan, M.J., McIntyre, M., Reid, J.L., Dominiczak, A.F. and Hamilton, C.A. (1999) Superoxide anion production is increased in a model of genetic hypertension: role of the endothelium. *Hypertension* **33**, 1353-1358.
- Liao, J.K., Shin, W.S., Lee, W.Y. and Clark, S.L. (1995) Ox-LDL decreases expression of eNOS. *J. Biol. Chem.* **270**, 319-324.
- Li, J., Li, W., Altura, B.T. and Altura, B.M. (2005) Peroxynitrite-induced relaxation in isolated rat aortic rings and mechanisms of action. *Toxicol. Appl. Pharmacol.* **209**, 269-276.
- López-Ongil, S., Hernandez-Perera, O., Navarro-Antolin, J., Perez de Lema, G., Rodriguez-Puyol, M., Lamas, S. and Rodriguez-Puyol, D. (1998) Role of reactive oxygen species in the role of signaling cascade of cyclosporine A-mediated upregulation of eNOS in vascular endothelial cells. *Br. J. Pharmacol.* **124**, 447-454.
- Lum, H. and Roebuck, K.A. (2001) Oxidant stress and endothelial cell dysfunction. *Am. J. Physiol., Cell Physiol.* **280**, C719-C741.
- Luscher, T.F. and Vanhoutte, P.M. (1986) Endothelium-dependent contractions to acetylcholine in the aorta of SHR. *Hypertension* **8**, 344-348.
- McQuaid K.E., Smyth, E.M. and Keenan, A.K. (1996) Evidence for modulation of hydrogen peroxide-induced endothelial barrier dysfunction by nitric oxide in vitro. *Eur. J. Pharmacol.* **307**, 233-241.

Mizukawa, H. and Okabe, E. (1997) Inhibition by singlet molecular oxygen of the vascular reactivity in rabbit mesenteric artery. *Br. J. Pharmacol.* **121**, 63-70.

Muijsers, R.B.R., van den Worm, E., Folkerts, G., Beukelman, C.J., Koster, A.S., Postma, D.S. and Nijkamp, F.P. (2000) Apocynin inhibits peroxynitrite formation by murine macrophages. *Br. J. Pharmacol.* **130**, 932-936.

Ohara, Y., Peterson, T.E. and Harrison, D.G. (1993) Hypercholesterolemia increases endothelial superoxide production. *J. Clin. Invest.* **91**, 2546-2551.

Panza, J.A., Casino, P.R., Kilcoyne, R.N. and Quyyumi, A.A. (1993) Role of endothelium-derived NO in the abnormal endothelium-dependent vascular relaxation of patients with essential hypertension. *Circulation* **87**, 1468-1474.

Rodrigo, E., Maeso, R., Garcia, R. and Lahera, V. (1997) Endothelial dysfunction in SHR: consequences of chronic treatment with losartan and captopril. *J. Hypertens.* **15**, 613-618.

Shen, J.Z., Zheng, X.F. and Kwan, C.Y. (2000) Differential contractile actions of reactive oxygen species on rat aorta: selective activation of ATP receptor by H₂O₂. *Life Sci.* **66**, 291-296.

Taddei, S., Viridis, A., Mattei, P. and Salvetti, A. (1993) Vasodilation to acetylcholine in primary and secondary forms of human hypertension. *Hypertension* **21**, 929-933.

Tesmafiarim, B., Halpern, W. (1988) Endothelium-dependent and endothelium-independent vasodilators in resistance arteries from hypertensive rats. *Hypertension* **11**, 440-444.

Wei, E.P., Kontos, H.A. and Beckman, J.S. (1996) Mechanisms of cerebral vasodilatation by superoxide, H₂O₂ and peroxynitrite. *Am. J. Physiol.* **271**, H1262-H1266.

Yu, B.P. (1994) Cellular defenses against damage from reactive oxygen species. *Physiol. Rev.* **74**, 139-162.




Fig. 1. (A) Acetylcholine (ACh)- and sodium nitroprusside (SNP)-induced relaxant responses in phenylephrine-precontracted aortic rings. Effects of mechanical and pharmacological (by L-NAME, 50 μ M) inactivation of endothelium on ACh-induced (B) and SNP-induced (C) relaxations. Results are expressed as mean \pm SEM from 8 separate experiments. * $P < 0.05$ compared to endothelium-intact rings.




Fig. 2. (A) Effects of low dose pyrogallol (Pyro) on acetylcholine (ACh)-induced aortic relaxations. (B) Effects of high dose pyrogallol on ACh-induced aortic relaxations. Results are expressed as mean±SEM from 8 separate experiments. * $P < 0.05$ compared to ACh-induced relaxations.




Fig. 3. Effects of O_2^- -scavengers, Tiron (**A**) and MPG (**B**) and exogenous superoxide dismutase (SOD, **C**) and the inhibitor DETCA (**D**) on acetylcholine (ACh)-induced aortic relaxations in the absence and presence of pyrogallol (Pyro). Results are mean \pm SEM from 8 separate experiments. * $P < 0.05$ compared to ACh-induced relaxations.




Fig. 4. Effects of DMSO (**A**), catalase (Cat, **B**) and Mn(III)TMPyP (**C**) on acetylcholine (ACh)-induced aortic relaxations in the absence and presence of pyrogallol (Pyro). Results are expressed as mean \pm SEM from 8 separate experiments. * P <0.05 compared to ACh-induced relaxations.




Fig. 5. Peroxynitrite (OONO⁻)- and hydrogen peroxide (H₂O₂)-induced relaxant responses in endothelium-intact [E(+)] and endothelium-denuded [E(-)] aortic rings. Results are expressed as mean±SEM from 8 separate experiments.




Fig. 6. (A) Phenylephrine (PE)-induced contractile responses in endothelium-intact [E(+)] and endothelium-denuded [E(-)] aortic rings. (B) Effects of different doses of pyrogallol (Pyro) on PE-induced contractile responses in E(+) aortic rings. (C) Effect of pharmacological inactivation of endothelium by L-NAME (50 μ M) on PE-induced contractility in the presence of different concentrations of pyrogallol in E(+) aortic rings. Results are expressed as mean \pm SEM from 8 separate experiments. * P <0.05 compared to PE-mediated contractile responses in E(+).




Fig. 7. (A) KCl-induced aortic contractile responses in endothelium-intact [E(+)] aortic rings in the absence and presence of pyrogallol. (B) KCl-induced aortic contractile responses in endothelium-denuded [E(-)] aortic rings in the absence and presence of pyrogallol. (C) KCl-induced aortic contractile responses in L-NAME-treated E(+) in the absence and presence of pyrogallol. * $P < 0.05$ compared to sole KCl-induced contractile response.




Fig. 8. Phenylephrine (PE)-induced aortic contraction in the absence and presence of Tiron and MPG alone or in combination with or pyrogallol in endothelium-intact [E(+)] aortic rings. Results are expressed as mean \pm SEM from 8 separate experiments. * P <0.05 compared to E(+) alone.

Table 1. Effect of different concentrations of pyrogallol on superoxide anion (O₂⁻) generation and LDH release

Pyrogallol	Lucigenin chemiluminescence (RLU/g.min)	Cytochrome C (pmol/mg tissue)	DHE oxidation (increase in % generation)	LDH release (~fold)
None	30 ± 6.5	9 ± 1.3	1.0 ± 0.0	1.00 ± 0.00
10 nM	100 ± 8.2*	36 ± 3.2*	24 ± 3.5	1.09 ± 0.11
100 nM	176 ± 13.9*	89 ± 9.3*	33 ± 4.6	1.13 ± 0.10
1 μM	340 ± 17.7*	183 ± 14.4*	39 ± 7.5	1.05 ± 0.09
10 μM	786 ± 24.1*	326 ± 22.1*	63 ± 9.1	0.91 ± 0.12
100 μM	978 ± 31.6*	541 ± 27.3*	79 ± 9.9	0.93 ± 0.11

The level of O₂⁻ was measured in thoracic aortic rings using lucigenin-enhanced chemiluminescence, dihydroethidium (DHE) oxidation and superoxide dismutase-inhibitable cytochrome C reduction assays while the level of cellular injury was measured by lactate dehydrogenase (LDH) leakage into culture medium. Results are expressed as mean ± SEM from three independent experiments. RLU; relative light unit. **P*<0.05, difference within each experimental group.

Table 2. Effect of different concentrations of pyrogallol on OONO⁻ and H₂O₂ generation

Pyrogallol	123-Rhodamine fluorescence (fold difference)	2',7'-dichlorodihydrofluorescein oxidation (arbitrary units)
None	1.0 ± 0.0	100 ± 0.0
10 nM	1.12 ± 0.06	135 ± 8*
100 nM	1.26 ± 0.05*	144 ± 10*
1 μM	1.37 ± 0.10*	166 ± 11*
10 μM	1.43 ± 0.12*	187 ± 13*
100 μM	1.51 ± 0.13*	205 ± 12*

The levels of OONO⁻ and H₂O₂ were measured in thoracic aortic rings through detections of 123-rhodamine and 2',7'-dichlorodihydrofluorescein (DCHF) oxidations. Results are expressed as mean ± SEM from three independent experiments.

**P*<0.05, difference to pyrogallol-untreated samples.