



Article (refereed) - postprint

Harris, Michael P.; Bogdanova, Maria I.; Daunt, Francis; Wanless, Sarah. 2012
Using GPS technology to assess feeding areas of Atlantic puffins *Fratercula arctica*.
Ringing & Migration, 27 (1). 43-49. [10.1080/03078698.2012.691247](https://doi.org/10.1080/03078698.2012.691247)

Copyright © 2012 British Trust for Ornithology

This version available <http://nora.nerc.ac.uk/18140/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This is an Author's Accepted Manuscript of an article published in *Ringing & Migration*, 27 (1). 43-49 [copyright British Trust for Ornithology], available online at: [http://www.tandfonline.com/ 10.1080/03078698.2012.691247](http://www.tandfonline.com/10.1080/03078698.2012.691247)

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **Feeding areas of Atlantic Puffins *Fratercula arctica* assessed by GPS**
2 **technology**

3 MICHAEL P. HARRIS*, MARIA I. BOGDANOVA, FRANCIS DAUNT and SARAH WANLESS

4 *Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK*

5

6 Large areas of sea around Britain have been identified as potential sites for renewable energy
7 development heightening the need for information about important areas for seabirds. The Atlantic
8 Puffin *Fratercula arctica* is one of the commonest seabirds in north-east Britain yet little is known
9 about where individuals feed. We used back-mounted GPS loggers to track Atlantic Puffins breeding
10 at a colony close to where wind farms are proposed. During chick rearing, birds made two types of
11 feeding trip: long absences that included an overnight stay at distant (38 – 66 km) feeding areas and
12 short daytime excursions to areas much nearer the colony (9 – 17 km). There was considerable
13 overlap of the distant feeding area with the proposed wind farms. However, Atlantic Puffins are known
14 to be sensitive to disturbance and comparison of the GPS birds and individuals without loggers
15 showed that the birds' behaviour had been disrupted by some aspect of the procedure. While the
16 areas used by the GPS birds accorded with expectations based on other methods, it is possible that
17 results from GPS birds represent a worst case scenario and overestimate the degree of overlap with
18 the proposed wind farms.

19

20

21

22 Running head: *Feeding areas of Puffins*

23 *Corresponding author.

24 mph@ceh.ac.uk

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

Large areas of coastal and offshore waters around Britain have been identified as potential sites for renewable energy development (The Crown Estate 2006). Many of these sites are in the vicinity of major breeding colonies of seabirds and have the potential to impact on these birds, for example where they overlap with foraging hotspots (Grecian *et al* 2010, Witt *et al* 2012). Thus information on where seabirds feed is essential for assessing the environmental risk of such developments. During the breeding season the foraging range of most species is constrained by the need to return to the colony at frequent intervals to relieve the mate and/or feed the brood. Switching to alternative feeding areas may not, therefore, be a viable option. Knowledge of where individuals from particular colonies feed is largely lacking because many seabirds fly fast (typical flight speeds of 30 – 70 km/h; Pennycuik 1987) and use areas well out of sight of a land-based observer. Until recently, Global Positioning System (GPS) loggers that provide detailed data on location have been too large to deploy on many of the seabird species breeding in the UK, but devices now available are potentially small enough to be used on birds weighing as little as 300-400 g. However, despite it having long been recognised that bird-borne devices can affect the flight and foraging behaviour of seabirds, particularly auks and other diving species (Wilson *et al* 1986, Barron *et al* 2010, Vandenabeele *et al* 2012), the urgency with which information is required risks results being used uncritically.

The Atlantic Puffin *Fratercula arctica* (hereafter referred to as Puffin) is a small, burrow-nesting, pursuit-diving seabird. It is one of the most abundant seabirds breeding in north-east Britain (Harris & Wanless 2004) yet a recent review of data relevant to the foraging ranges of British seabirds revealed that less is known of the feeding areas used by Puffins than most other species and more information is urgently needed (Thaxter *et al* 2012). One reason for this knowledge gap is that Puffins are sensitive to disturbance, particularly handling (Wernham 1993, Rodway *et al* 1996, Harris & Wanless 2011) and, like the Tufted Puffin *F. cirrhata* (Whidden *et al.* 2007), seem intolerant of back-mounted devices (personal observations). However, extensive areas within the likely foraging range of Puffins at many of the major colonies in Scotland have recently been designated as potential sites for the generation of wave, tidal or wind energy (The Crown Estate 2006). Given the heightened

54 imperative for information on populations in these areas, we carried out a small study to assess the
55 feasibility of using some of the smallest available GPS loggers to obtain information on the foraging
56 areas used. The original intention was to scale up the study in subsequent years but the results of
57 these first deployments were sufficiently worrying to be worth putting on record.

58

59 **METHODS**

60 The study was carried out, under the relevant licences, at the colony of c.40,000 pairs of Puffins on
61 the Isle of May (56° 11'N, 2° 34'W) in the Firth of Forth, south-east Scotland (Figure 1) during the
62 chick rearing period in 2010.

63 On 3 June, three Puffins with chicks were removed from their burrows and each fitted with a
64 dummy package of the same dimensions (40 x 22 x 11 mm) and weight (15 g) as the larger type of
65 GPS logger intended to be used for data collection. Each dummy was attached to a few central back
66 feathers using three strips of waterproof tape (Tesa Ltd; total mass 1.5 g). To minimise handling time
67 the birds were not weighed. However, the mean mass of a sample of Puffins feeding chicks caught
68 concurrently in the same area was 386 g \pm SE 3.3 (n = 75), which indicated that, on average, a
69 dummy plus attachment tape was 4.3% of the body mass. Birds were ringed and then returned to
70 their burrow. The whole procedure took 3-5 min. To check how long dummies stayed attached and if
71 there was any evidence that birds deserted their burrows and/or stopped feeding their chicks,
72 opportunistic observations were made from a hide c. 20 m away from the burrows over the following
73 two weeks.

74 Results from the dummy packages were encouraging (see Results). Six GPS loggers (i-gotU
75 GT-120, Mobile Action Technology) that had been repackaged to make them more streamlined and
76 which were of the same dimensions and weight as the dummies, were therefore attached using the
77 same method to Puffins with chicks on 19 June. Another of these loggers plus three smaller ones
78 (GiPSy-2, TechnoSmart; 48 x 20 x 9 mm, 11.5 g with tape attachment; 3.0% of adult mass) were
79 attached on 23 June. The larger loggers recorded one position (accurate to c. 20 m) every 1 min,
80 whereas the smaller ones recorded continuously every second for 1 min (same accuracy) but were
81 then off for 5 min. The larger loggers recorded data for 2-3 days and the smaller ones for 1-2 days.

82 One to two days after deployment, any instrumented bird seen entering its burrow was re-caught, the
83 device removed and the bird put back in the burrow. This procedure took 1-2 min. Deployments and
84 retrieval details are given in Table 1.

85 Data from both types of logger were used to identify when birds were in the vicinity of the
86 colony (assumed to correspond to periods when signals were constant and positions were within 500
87 m of the release location) or on foraging trips (all other periods). Neither logger type recorded a bird's
88 diving activity directly. However, the signal to the satellite is lost when the device is underwater and
89 Puffins typically spend c.78% time underwater when diving (Wanless *et al* 1988; Harris & Wanless
90 2011), so locations where the bird remained approximately stationary (distance between subsequent
91 locations < 500 m) and fixes were obtained intermittently instead of every minute were assumed to
92 correspond to diving. The approach was not feasible for the smaller loggers due to the lower
93 recording frequency.

94 Population distributions for all fixes and diving fixes were derived from fixed kernel analysis
95 performed in ArcGIS (Hawth's Analysis Tools 3.27; Beyer 2004) using a smoothing parameter (*h*) of
96 6000 m (identified using the Least-squares cross-validation method; Worton 1989) and a cell size of
97 1000 m. Maps with 50% (representing the core area of use), 70% and 90% (active area of use) kernel
98 density contours were produced in a Lambert equal-area azimuthal (North Pole) projection. To assess
99 the extent of overlap between population range and proposed offshore wind farms, the proportion of
100 50% and 90% kernels within the proposed sites was calculated. To assess the extent of overlap
101 between population range and proposed offshore wind farms, the proportion of 50% and 90% kernels
102 within the proposed sites was calculated using ArcGIS.

103 To check for effects of dummies and loggers on colony attendance and chick provisioning, the
104 burrows of these birds were watched for a total of 38.7 h (mean 3.5 ± 0.9 h per day; $n = 11$ days,
105 mainly during the daily peak of feeding (0500 – 0900 h; Harris & Wanless 2011)). During watches we
106 recorded visits to burrows by these birds and their mates, and whether or not they had fish.
107 Observations ceased on 1 July and the contents of burrows of dummy and logger birds were checked
108 the next day to establish whether a chick was still present. Some burrows were too long for chicks to
109 be caught but where possible chicks were weighed (to nearest g) and their wing length measured (to
110 nearest mm) as a proxy for age. These measurements were compared with those of 24 chicks in an

111 adjacent part of the colony so as not to increase disturbance of the focal birds, obtained on 30 June
112 and 3 July. Thirty-four chicks whose parents were not handled that were known to be alive at the start
113 of the logger work were checked on 1 - 3 July to see how many had survived.

114 To quantify chick provisioning rates of undisturbed Puffins, all-day watches of 60 and 68
115 burrows elsewhere in the colony were carried out on 24 and 28 June, respectively.

116

117 **RESULTS**

118 All three Puffins with dummy loggers continued to feed their chicks. One bird lost its dummy device
119 after c. 5 days but the other two dummies were still attached 13 and 20 days later when the birds
120 were re-caught and the dummies removed (Table 1). On the basis of these observations we
121 concluded that the behaviour of these Puffins had not been severely disrupted and there was a good
122 chance of obtaining data from loggers.

123 Of the ten GPS loggers that were subsequently deployed, seven (70%) were retrieved.
124 These provided information on 15 trips away from the colony over 14 bird-days (Table 2, Figure 1). It
125 was apparent that Puffins made two types of trips: long absences where individuals were away
126 overnight (12 cases lasting 15.0 – 41.0 h; maximum distance from the colony 37.8 – 65.5 km) and
127 shorter excursions (3 cases lasting 0.7 - 3.8 h; 9.3 – 17.5 km from the colony). Although some diving
128 occurred within 5 km of the Isle of May, most took place during the long trips when the birds were in
129 areas 30-60 km to the east and north-east of the island. Consequently, there was considerable
130 overlap between the Puffin kernel distributions and proposed offshore wind farms (Figure 2). For all
131 fixes, there was 84% overlap for the 50% kernel and 52% overlap for the 90% kernel. For diving fixes,
132 the overlaps were 73% and 50%, respectively.

133 However, observations indicated that the behaviour of the Puffins carrying loggers differed in
134 several important respects from those without loggers. Instrumented birds made an average of 1.07
135 trips per day (range 0.67 – 2.0; Table 1). Chicks in undisturbed burrows elsewhere in the colony at
136 this time received 4.1 ± 0.3 ($n = 68$ burrows) and 4.9 ± 0.3 ($n = 60$ burrows) feeds per day,
137 respectively. Both members of a Puffin pair feed their chick so to attain equivalent feeding rates as
138 undisturbed pairs, uninstrumented adults in logger pairs would have to deliver food at a rate 3-4 times
139 greater than their instrumented mates. The limited data on feeding rates of members of a pair suggest

140 that such a marked disparity is unlikely (personal observations). In addition, two of the seven logger
141 birds (29%) had returned without fish when they were re-caught (Table 2). In contrast, at the same
142 time Puffins without devices returned without fish significantly less frequently (2 (2%) of 81 returns;
143 Fisher Exact test $P = 0.03$).

144 Although the overall retrieval rate of dummy and logger devices was high (69%), observations
145 of the birds' burrows indicated that attendance patterns had been disrupted both after initial capture
146 and retrieval of the logger. Thus, three birds deserted after deployment while another was not seen
147 for two days despite frequent checks of the study area (Table 1). A further two birds deserted
148 following logger retrieval while another was never seen going down its burrow again even though its
149 chick was alive. On a more encouraging note, three logger birds and two dummy birds did then
150 behave more normally. For example, Bird 7 brought in three loads of fish within 3 h, and three loads
151 within 5 h, three and four days respectively after the removal of its logger. Overall, it returned with
152 more food loads than its mate (13 cf. 8 loads; $n = 34.5$ h of observation). Similarly, Bird 13 brought in
153 three loads in 5.5 h four days of observations after its logger was taken off, twice as many as its
154 partner (8 cf. 4 loads; $n = 22.7$ h).

155 The check of burrow contents on 2 July indicated that 3 (23%) no longer contained chicks.
156 This loss rate was markedly higher than that recorded over the same period at control burrows (3/34
157 i.e. 9%), although the difference was not statistically significant (Fisher Exact test, $P = 0.3$). Of the four
158 chicks that were weighed, only that of Bird 1 was of normal mass compared to the 24 control chicks
159 (Figure 3).

160

161 **DISCUSSION**

162 A standard criterion for assessing the impact of loggers is retrieval rate, with high rates assumed to
163 reflect low impact. On this basis, our overall retrieval rate of 69% of dummies and devices did not
164 cause undue concern. However, because the Puffin is well-documented as being less tolerant of
165 disturbance than many other seabirds, we were alert to additional effects. It became obvious that
166 some aspect of the deployments affected attendance behaviour sufficiently to impact on chick weights
167 and breeding success although small sample sizes made statistical evaluation difficult. These effects

168 need to be kept in mind when considering locational data for which there is no direct method to
169 compare behaviour of birds with and without devices.

170 Data from the GPS loggers indicated that Puffins were using areas many tens of kilometres
171 away from the Isle of May and these overlapped with the proposed wind farm sites. The only
172 previous direct information on where Puffins from this colony forage when they have chicks was a
173 single bird tracked using VHF radio-telemetry in 1987 (Wanless *et al* 1990). Eleven (73%) of 15 trips
174 made by this bird were to feeding sites within the c.10 km detection range of the receiving station on
175 the Isle of May but on four trips it flew out of range. Flight directions of both the close and more distant
176 trips were predominantly to the south-east of the island and thus there was little overlap with areas
177 used in 2010 that were mainly to the east and north-east. The only other GPS data for Puffins that we
178 are aware of are unpublished results of Richard Bevan from the Farne Islands, 100 km south of the
179 Isle of May, that suggest that the majority of foraging there occurs within 20 km of the colony (BBC
180 2010). Other information on foraging ranges of Puffins comes from ship-based surveys of Puffins in
181 the vicinity of breeding colonies. These suggest that on Skomer, south-west Wales, 85% of Puffins
182 carrying fish were within 15 km, 99% were within 25 km and the maximum recorded distance from the
183 colony was 35 km (Stone *et al* 1992). Similarly, Puffins seen feeding near St Kilda, north-west
184 Scotland were mostly within 40 km of the nearest colony (Leaper *et al* 1988). Around Newfoundland,
185 Puffins generally forage 3-5 km from the colony but sometimes up to 70 km away (Bradstreet &
186 Brown 1985, Schneider *et al.* 1990). However, when feeding conditions are poor, Puffins appear to
187 use more distant areas since they have been recorded carrying fish up to 137 km and 250 km from
188 the nearest colonies in Norway and the Faeroe Islands, respectively (Anker-Nilssen & Lorentsen
189 1990, M. L. Tasker, pers. comm.).

190 Whilst foraging range of individuals in our study accorded with previous work, the question
191 remains as to whether the positional information we obtained is likely to be representative of the
192 foraging distribution of the population as a whole. Work in 2008 using small leg-mounted time-depth
193 recorders (weight 2.7 g) deployed overwinter, showed that Puffins feeding young on the Isle of May
194 did not dive during the night, had peaks of diving late in the evening and early in the morning and the
195 mean start of diving by 10 individuals on 58 mornings in early July was 27 ± 4 min before sunrise
196 (Harris & Wanless 2011, unpublished data). These observations agree with the daily peak of Puffins

197 returning with fish that occurs a few hours after dawn, and support the hypothesis that the parents fly
198 to feeding grounds for self-feeding, perhaps on prey that are easily caught but too low food value to
199 carry back to a chick, late in the evening. They then stay on the feeding grounds overnight, feed
200 again soon after dawn and return to the colony with fish for the chick an hour or two later.

201 It is more problematic to use the dive data to evaluate what happens during the remainder of
202 the day. Chick provisioning rates were lower in GPS birds suggesting that shorter, and therefore quite
203 possibly closer, trips were under-represented in the daytime data. Such a bias would mean that the
204 true kernel distribution would be concentrated closer inshore (most likely between the current kernel
205 distribution and the colony) and the overlap with the proposed wind farm developments would be
206 reduced. Thus, our recorded distributions can be viewed as a worst case scenario in terms of
207 interactions of Puffins from the Isle of May with the proposed wind farm sites. However, it is likely that
208 even if Puffins make more use of closer areas than we recorded, most, if not all, will spend time in the
209 distant areas. Further insights could potentially be gained by combining kernel density distributions
210 with those from at-sea surveys although the latter would include nonbreeding Puffins and those from
211 other colonies. Concentrations of Puffins that coincide with those indicated by the kernel analysis
212 would provide reassurance that the distant locations for GPS birds are robust. By a similar argument,
213 hotspots of birds close to the colonies would provide support for the suggestion that shorter trips are
214 under-reported by the GPS deployments.

215 Device effects are species specific, and the Puffin is well documented as being intolerant of
216 disturbance and thus likely to be towards the upper end of the device effect scale. More generally,
217 even in species where effects are less pronounced, responses will probably vary according to
218 environmental conditions. In the case of Puffins on the Isle of May, 2010 appeared to be an average
219 season in terms of breeding success and provisioning activity so it is possible that disruption might
220 have been even more severe if conditions had been less favourable. To date, impact assessments
221 for proposed wind farms in UK waters have relied heavily on data from aerial and ship-based surveys.
222 These methods have several serious limitations notably that the provenance and breeding status of
223 the birds recorded is unknown. Bio-logging, in particular GPS telemetry provides a complementary
224 approach that overcomes these problems. However, recognizing that the behaviour of tagged
225 individuals will probably be affected to some extent needs to be taken into account and attempts

226 made to quantify the impact, particularly any implications this might have for interactions between
227 birds and marine renewables.

228

229 **ACKNOWLEDGEMENTS**

230 We thank Scottish Natural Heritage for permission to work on the Isle of May National Nature
231 Reserve, Mark Newell for help with the fieldwork and two reviewers for their helpful comments on the
232 manuscript.

233

234 **REFERENCES**

235 **Anker-Nilssen, T. & Lorentsen, S.-H.** (1990) Distribution of Puffins *Fratercula arctica* feeding off
236 Røst, northern Norway, during the breeding season, in relation to chick growth, prey and
237 oceanographical parameters. *Polar Research* **8**, 67-76.

238 **Barron, D.G., Brawn, J.D. & Weatherhead, P.J.** (2010) Meta-analysis of transmitter effects on avian
239 behaviour and ecology. *Methods in Ecology and Evolution* **1**, 180-187.

240 **BBC** (2010) Tags reveal puffin food 'hotspot'. Available online at
241 <http://www.bbc.co.uk/news/10528822> (accessed 10 April 2012).

242 **Beyer, H.L.** (2004) Hawth's Analysis Tools for ArcGIS. Available online at
243 <http://www.spatial ecology.com/htools> (accessed 7 April 2012).

244 **Bradstreet, M.S.W. & Brown, R.G.B.** (1985) Feeding ecology of the Atlantic Alcidae. In: *The Atlantic*
245 *Alcidae* (ed. by D.N. Nettleship & T.R. Birkhead), pp. 264-318. Academic Press, London.

246 **Grecian, W.J., Inger, R., Attrill, M.J., Bearhop, S., Godley, B.J., Witt, M.J. & Votier, S.C.** (2010)
247 Potential impacts of wave-powered marine renewable energy installations on marine birds.
248 *Ibis* **15**: 683-697.

249 **Harris, M. P. & Wanless, S.** (2004) The Atlantic Puffin *Fratercula arctica*. In *Seabird populations in*
250 *Britain and Ireland* (eds Mitchell, I. P., Newton, S. F., Ratcliffe, N. and Dunn, T. E.) pp 392-
251 406. T. & A.D. Poyser, London.

252 **Harris, M.P. & Wanless, S.** (2011) *The Puffin*. T. & A.D. Poyser, London.

253 **Leaper, G.M., Webb, A., Benn, S., Prendergast, H.D.V., Tasker, M.L. & Schofield, R.** (1988)
254 Seabird studies around St Kilda, June 1987. Nature Conservancy Council, CSD Report No.
255 804, Peterborough.

256 **Pennycuik, C. J.** (1987) Flight of auks (Alcidae) and other northern seabirds compared with
257 southern procellariiforms. *Journal of Experimental Biology* **128**, 335-347.

258 **Rodway, M. S., Montevecchi, W. A. & Chardine, J. W.** (1996) Effects of investigator disturbance on
259 breeding success of Atlantic puffins *Fratercula arctica*. *Biological Conservation*. **76**, 311-319.

260 **Schneider, D.C., Pierotti, R. & Threlfall, W.** (1990) Alcid patchiness and flight direction near a
261 colony in east Newfoundland. *Studies in Avian Biology*, **14**, 23-35.

262 **Stone, C.J., Webb, A., Barton, T.R. & Gordon, J.R.W.** (1992) Seabird distributions around Skomer
263 and Skokholm Islands, June 1962. JNCC Report No.152. Peterborough

264 **Thaxter, C.B., Lascelles, B., Sugar, K., Cook, A.S.C.P., Roos, S., Bolton, M., Langston, R.H.W. &**
265 **Burton, N.H.K.** (2012) Seabird foraging ranges as a preliminary tool for identifying candidate
266 Marine Protected Areas. *Biological Conservation* doi:10.1016/j.biocon.2011.12.009.

267 **The Crown Estate** (2006) Available online at http://www.thecrownestate.co.uk/offshore_wind_energy
268 (accessed 10 April 2012).

269 **Vandenabeele, S.P., Shepard, E.L., Grogan, A. & Wilson, R.P.** (2012) When three per cent may
270 not be three per cent: device-equipped seabirds experience variable flight constraints. *Marine*
271 *Biology* **159**, 1-14.

272 **Wanless, S., Harris, M. P. & Morris, J. A.** (1990). A comparison of feeding areas used by individual
273 Common Murres (*Uria aalge*), Razorbills (*Alca torda*) and an Atlantic Puffin (*Fratercula*
274 *arctica*) during the breeding season. *Colonial Waterbirds* **13**, 16-24.

275 **Wanless, S., Morris, J.A. & Harris, M.P.** (1988). Diving behaviour of Guillemot *Uria aalge*, Puffin
276 *Fratercula arctica* and Razorbill *Alca torda* as shown by radio-telemetry. *Journal of Zoology*,
277 *London* **216**, 73-81.

278 **Wernham, C. V.** (1993) *Ecology and energetics of breeding Puffins (Fratercula arctica); variation in*
279 *individual reproductive effort and success*. PhD Thesis, University of Stirling.

280 **Whidden, S. E., Williams, C. T., Breton, A. R. & Buck, C. L.** (2007) Effects of transmitters on the
281 reproductive success of Tufted Puffins. *Journal of Field Ornithology* **78**, 206-212.

- 282 **Wilson, R. P., Grant, W. S. & Duffy, D. C.** (1986) Recording devices on free-ranging marine animals:
283 does measurement affect foraging performance? *Ecology* **67**, 1091-1093.
- 284 **Witt, M.J., Sheehan, E.V., Bearhop, S., Broderick, A.C., Conley, D.C., Cotterell, S.P., Crow, E.,**
285 **Grecian, W.J., Halsband, C., Hodgson, D.J., Hosegood, P., Inger, R., Miller, P.I., Sims,**
286 **D.W., Thompson, R.C., Vanstaen, K., Votier, S.C., Attrill, M.J. & Godley, B.J.** (2012)
287 Assessing wave energy effects on biodiversity: the Wave Hub experience. *Philosophical*
288 *Transactions of the Royal Society A-Mathematical Physical and Engineering Science* **370**,
289 502-529.
- 290 **Worton, B. J.** (1989) Kernel methods for estimating the utilization distribution of home range studies.
291 *Ecology* **70**, 164-168.

Table 1. Details of deployment of dummy devices and GPS loggers on Atlantic Puffins on the Isle of May and their effect on burrow attendance and breeding success.

Bird No.	Logger type	Date attached	Hours deployed	Trips made	Subsequent sightings	Contents of burrow on 2 July
1	Dummy	3 June	488.0	-	Regular throughout	Chick healthy
2	Dummy	3 June	306.0	-	Regular throughout	Chick alive
3	Dummy	3 June	150-170*	-	Reluctant to go down burrow	Burrow empty
4	i-gotU	19 June	-	-	Not seen after deployment	Chick very thin
5	i-gotU	19 June	-	-	Not seen after deployment	Chick alive
6	i-gotU	19 June	48.1	3	Not seen after removal	Chick alive
7	i-gotU	19 June	47.1	2	Regular after removal	Chick healthy
8	i-gotU	19 June	70.7	3	Not seen after removal	Burrow empty
9	i-gotU	19 June	47.7	2	Delayed return, burrow dug-out	Burrow empty
10	GiPSy-2	23 June	-	-	Not seen after deployment	Chick thin
11	GiPSy-2	23 June	74.1	2	Never entered burrow after removal	Chick alive
12	i-gotU	23 June	24.9	1	Regular after removal	Chick alive
13	GiPSy-2	23 June	26.4	2	Regular after removal	Chick alive

Note: * Lost during this period

Table 2. Duration and maximum distance of trips made by Atlantic Puffins fitted with GPS loggers on the Isle of May. Where known, whether or not a bird was carrying a fish when it was recaptured is also shown.









Bird	Trip	Trip duration (hours)	Maximum distance from colony (km)	Carrying fish at recapture
6	1	3.8	17.2	
6	2	0.7	9.3	
6	3	41.0	37.6	yes
7	1	15.0	39.6	
7	2	24.7	63.3	yes
8	1	22.6	42.0	
8	2	23.1	47.4	
8	3	>12.5*	>39.7	no
9	1	23.3	43.1	
9	2	23.1	54.8	yes
11	1	24.4	49.7	
11	2	>17.5*	>37.7	no
12	1	24.5	48.2	yes
13	1	23.9	65.5	
13	2	1.2	17.5	yes

*Logger out of storage space




Figure 1. Foraging trips of Atlantic Puffins breeding on the Isle of May in 2010. The top left panel shows the location of the Isle of May in a British Isles context, the other panels show the GPS tracks of individual birds (bird number given in top left corner).

Figure 2. a) All fixes of Atlantic Puffins from the Isle of May obtained from i-gotU GT-120 GPS loggers overlaid on the areas (light shading) of sea of the Firth of Forth being considered for the location of wind turbines (The Crown Estate 2006); b) 50, 70 and 90% kernel density contours based on all locations obtained from i-gotU GT-120 loggers; c) 50, 70 and 90% kernel density contours based on diving locations obtained from i-gotU GT-120 loggers.

Figure 3. Mass in relation to wing length of Atlantic Puffin chicks where one adult had had a dummy or a GPS logger attached compared to chicks where neither adult had been handled.



Harris et al. Figure 1.



Harris et al. Figure 2.

Figure 3. Mass in relation to wing length of Puffin chicks where one adult had had a dummy or a GPS logger attached compared to chicks where the adult had not been handled.

