

## FRACTIONAL BOUNDARY VALUE PROBLEMS ON THE HALF LINE

Assia Frioui, Assia Guezane-Lakoud, and Rabah Khaldi

*Communicated by Jean Mawhin*

**Abstract.** In this paper, we focus on the solvability of a fractional boundary value problem at resonance on an unbounded interval. By constructing suitable operators, we establish an existence theorem upon the coincidence degree theory of Mawhin. The obtained results are illustrated by an example.

**Keywords:** boundary value problem at resonance, existence of solution, unbounded interval, coincidence degree of Mawhin, fractional differential equation.

**Mathematics Subject Classification:** 34B40, 34B15.

### 1. INTRODUCTION

In this paper, we are concerned with the existence of solutions for the following fractional differential equation

$$D_{0+}^q x(t) = f(t, x(t)) + e(t), \quad t \in (0, \infty), \quad (1.1)$$

subject to the nonlocal conditions

$$I_{0+}^{2-q} x(0) = 0, \quad D_{0+}^{q-1} x(\infty) = \frac{\Gamma(q)}{\eta^{q-1}} x(\eta), \quad (1.2)$$

where  $D_0^q$  denotes the Riemann Liouville fractional derivative of order  $q$ ,  $1 < q < 2$ ,  $\eta > 0$ ,  $f : [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$  and  $e : [0, \infty) \rightarrow \mathbb{R}$  are given functions satisfying some conditions.

The problem (1.1)–(1.2) happens to be at resonance in the sense that the dimension of the kernel of the linear operator  $Lx = D_{0+}^q x$  is not less than one under boundary conditions (1.2).

It should be noted that in recent years, there have been many works related to boundary value problems at resonance for ordinary differential equations. We refer the reader to [7–9, 11, 13, 14, 16] and the references therein.

However, the articles on the existence of solutions of fractional differential equations on the half-line are still few in number, and most of them deal with problems under nonresonance conditions. For some recent articles investigating resonant and nonresonant fractional problems on the unbounded interval, see [1–5, 10, 17].

Recently fractional differential equations have been investigated by many researchers and different methods have been used to obtain such fixed point theory, upper and lower solutions method, coincidence degree theory, etc.

The rest of this paper is organized as follows. In Section 2, we give some necessary notations, definitions and lemmas. In Section 3, we study the existence of solutions of (1.1)–(1.2) by the coincidence degree theory due to Mawhin. Finally, we give an example to illustrate the obtained results.

## 2. PRELIMINARIES

First recall some notation, definitions and theorems which will be used later.

Let  $X$  and  $Y$  be two real Banach spaces and let  $L : \text{dom } L \subset X \rightarrow Y$  be a linear operator which is a Fredholm map of index zero, define the continuous projections  $P$  and  $Q$ , respectively, by  $P : X \rightarrow X$ ,  $Q : Y \rightarrow Y$  such that  $\text{Im } P = \ker L$ ,  $\ker Q = \text{Im } L$ . Then  $X = \ker L \oplus \ker P$ ,  $Y = \text{Im } L \oplus \text{Im } Q$ , thus  $L|_{\text{dom } L \cap \ker P} : \text{dom } L \cap \ker P \rightarrow \text{Im } L$  is invertible, denote its inverse by  $K_P$ .

Let  $\Omega$  be an open bounded subset of  $X$  such that  $\text{dom } L \cap \Omega \neq \emptyset$ , the map  $N : X \rightarrow Y$  is said to be  $L$ -compact on  $\overline{\Omega}$  if  $QN(\overline{\Omega})$  is bounded and  $K_P(I - QN) : \overline{\Omega} \rightarrow X$  is compact. Since  $\text{Im } Q$  is isomorphic to  $\ker L$  then, there exists an isomorphism  $J : \text{Im } Q \rightarrow \ker L$ . It is known that the coincidence equation  $Lx = Nx$  is equivalent to  $x = (P + JQN)x + KP(I - Q)Nx$ .

**Theorem 2.1** ([15]). *Let  $L$  be a Fredholm operator of index zero and  $N$  be  $L$ -compact on  $\overline{\Omega}$ . Assume that the following conditions are satisfied:*

- (1)  $Lx \neq \lambda Nx$  for every  $(x, \lambda) \in [(\text{dom } L \setminus \ker L) \cap \partial\Omega] \times (0, 1)$ ,
- (2)  $Nx \notin \text{Im } L$  for every  $x \in \ker L \cap \partial\Omega$ ,
- (3)  $\deg(QN|_{\ker L}, \Omega \cap \ker L, 0) \neq 0$ , where  $Q : Y \rightarrow Y$  is a projection such that  $\text{Im } L = \ker Q$ .

*Then the equation  $Lx = Nx$  has at least one solution in  $\text{dom } L \cap \overline{\Omega}$ .*

**Theorem 2.2** ([6]). *Let  $C_\infty = \{y \in C([0, \infty)) : \lim_{t \rightarrow \infty} y(t) \text{ exists}\}$  equipped with the norm  $\|y\|_\infty = \sup_{t \in [0, \infty)} |y(t)|$ . Let  $F \subset C_\infty$ . Then  $F$  is relatively compact if the following conditions hold:*

- (1)  $F$  is bounded in  $C_\infty$ ,
- (2) the functions belonging to  $F$  are equicontinuous on any compact subinterval of  $[0, \infty)$ ,
- (3) The functions from  $F$  are equiconvergent at  $\infty$ .

Let  $X$  be the space

$$X = \left\{ x \in C([0, \infty)) : D_{0+}^{q-1}x \in C([0, \infty)), \right. \\ \left. \lim_{t \rightarrow \infty} e^{-t} |x(t)| \text{ and } \lim_{t \rightarrow \infty} e^{-t} \left| D_{0+}^{q-1}x(t) \right| \text{ exist} \right\}$$

endowed with the norm

$$\|x\| = \sup_{t \geq 0} e^{-t} |x(t)| + \sup_{t \geq 0} e^{-t} \left| D_{0+}^{q-1}x(t) \right|,$$

then  $X$  is a Banach space.

Let  $Y = L^1[0, \infty)$  equipped with the norm

$$\|y\|_1 = \int_0^\infty |y(t)| dt.$$

Define the operator  $L : \text{dom } L \subset X \rightarrow Y$  by  $Lx = D_{0+}^q x$ , where

$$\text{dom } L = \left\{ x \in X : D_{0+}^q x \in Y, I_{0+}^{2-q}x(0) = 0, D_{0+}^{q-1}x(\infty) = \frac{\Gamma(q)}{\eta^{q-1}}x(\eta) \right\} \subset X,$$

then  $L$  maps  $\text{dom } L$  into  $Y$ . Let  $N : X \rightarrow Y$  be the operator

$$Nx(t) = f(t, x(t)) + e(t), \quad t \in [0, \infty),$$

then the problem (1.1)–(1.2) can be written as  $Lx = Nx$ .

We recall the definition of the Riemann-Liouville fractional derivative and Riemann-Liouville fractional integral, we can find their properties in [12].

**Definition 2.3.** The Riemann-Liouville fractional integral of order  $p > 0$  is given by

$$I_{0+}^p g(t) = \frac{1}{\Gamma(p)} \int_0^t \frac{g(s)}{(t-s)^{1-p}} ds, \quad t > 0,$$

provided that the right-hand side exists.

**Definition 2.4.** The Riemann-Liouville fractional derivative  $D_{0+}^p g$  of order  $p > 0$  is given by

$$D_{0+}^p g(t) = \frac{1}{\Gamma(n-p)} \left( \frac{d}{dt} \right)^n \int_0^t \frac{g(s)}{(t-s)^{p-n+1}} ds, \quad t > 0,$$

where  $n = [p] + 1$  ( $[p]$  is the integer part of  $p$ ), provided that the right-hand side is point-wise defined on  $(0, \infty)$ .

**Lemma 2.5.** The homogenous fractional differential equation  $D_{a+}^q g(t) = 0$  has a solution

$$g(t) = c_1 t^{q-1} + c_2 t^{q-2} + \dots + c_n t^{q-n},$$

where  $c_i \in \mathbb{R}$ ,  $i = 1, \dots, n$  and  $n = [q] + 1$ .

**Lemma 2.6.** *Let  $p, q \geq 0$ ,  $f \in L_1[0, \infty)$ . Then the following assertions hold:*

1.  $I_{0+}^p I_{0+}^q f(t) = I_{0+}^{p+q} f(t) = I_{0+}^q I_{0+}^p f(t)$  and  $D_{0+}^q I_{0+}^q f(t) = f(t)$  for all  $t \in [0, \infty)$ .
2. If  $p > q > 0$ , then the formula  $D_{0+}^q I_{0+}^p f(t) = I_{0+}^{p-q} f(t)$  holds almost everywhere on  $t \in [0, \infty)$ , for  $f \in L_1[0, \infty)$  and it is valid at any point  $t \in [0, \infty)$  if  $f \in C[0, \infty)$ .
3. If  $q \geq 0$  and  $p > 0$ , then  $(I_{0+}^q t^{p-1})(x) = \frac{\Gamma(p)}{\Gamma(p+q)} x^{p+q-1}$ ,  $(D_{0+}^q t^{p-1})(x) = \frac{\Gamma(p)}{\Gamma(p-q)} x^{p-q-1}$ ,  $(D_{0+}^q t^{q-j})(x) = 0$ ,  $j = 1, 2, \dots, n$ , where  $n = [q] + 1$ .

### 3. MAIN RESULTS

We give the first result on the existence of a solution for the problem (1.1)–(1.2).

**Theorem 3.1.** *Assume that the following conditions are satisfied:*

- (H<sub>1</sub>) *There exists nonnegative functions  $\alpha, \beta \in L^1[0, \infty)$  such that for all  $x \in \mathbb{R}$   $t \in [0, \infty)$  we have*

$$|f(t, x) + e(t)| \leq e^{-t} \alpha(t) |x| + \beta(t). \quad (3.1)$$

- (H<sub>2</sub>) *There exists a constant  $M > 0$  such that for  $x \in \text{dom } L$  if  $|D_0^{q-1} x(t)| > M$  for all  $t \in [0, \infty)$*

$$\int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \neq 0. \quad (3.2)$$

- (H<sub>3</sub>) *There exists a constant  $M^* > 0$  such that for any  $x(t) = c_0 t^{q-1} \in \ker L$  with  $|c_0| > M^*$  either*

$$c_0 \left[ \int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \right] < 0 \quad (3.3)$$

or

$$c_0 \left[ \int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \right] > 0. \quad (3.4)$$

Then for every  $e \in Y$ , the problem (1.1)–(1.2) has at least one solution in  $X$ , provided  $(1 - 2A_1 \|\alpha\|_1) > 0$ , where

$$A_1 = M_1 + 1, \quad M_1 = \max_{t \geq 0} \left( \frac{e^{-t} t^{q-1}}{\Gamma(q)} \right) = \frac{[e^{-1}(q-1)]^{q-1}}{\Gamma(q)}.$$

In order to prove Theorem 3.1 we need some lemmas.

**Lemma 3.2.** *The operator  $L : \text{dom } L \subset X \rightarrow Y$  is a Fredholm operator of index zero. Furthermore, the linear projector operator  $Q : Y \rightarrow Y$  is defined by*

$$Qy(t) = Ae^{-t} \left( \int_0^\infty y(s)ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} y(s)ds \right) \tag{3.5}$$

and the linear operator  $K_P : \text{Im } L \rightarrow \text{dom } L \cap \ker P$  can be written as

$$K_P y(t) = I_{0+}^q y(t), \quad y \in \text{Im } L.$$

Moreover,

$$\|K_P y\| \leq A_1 \|y\|_1, \quad y \in \text{Im } L, \tag{3.6}$$

where

$$A^{-1} = 1 - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} e^{-s} ds.$$

*Proof.* Using Lemma 2.5, we get

$$x(t) = at^{q-1} + bt^{q-2}, \quad t > 0, \tag{3.7}$$

as a solution of  $Lx(t) = 0$ . Applying  $I_{0+}^{2-q}$  to both sides of the equality (3.7), then using condition  $I_{0+}^{2-q} x(0) = 0$ , we get  $b = 0$ . Thanks to condition  $D_{0+}^{q-1} x(\infty) = \frac{\Gamma(q)}{\eta^{q-1}} x(\eta)$ , we conclude

$$\ker L = \{x \in \text{dom } L : x(t) = at^{q-1}, a \in \mathbb{R}, t \in [0, \infty)\}.$$

Now the problem

$$D_{0+}^q x(t) = y(t) \tag{3.8}$$

has a solution  $x$  that satisfies the conditions  $I_{0+}^{2-q} x(0) = 0, D_{0+}^{q-1} x(\infty) = \frac{\Gamma(q)}{\eta^{q-1}} x(\eta)$  if and only if

$$\int_0^\infty y(s)ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} y(s)ds = 0. \tag{3.9}$$

In fact from (3.8) and together with the boundary condition  $I_{0+}^{2-q} x(0) = 0$  we get

$$x(t) = I_{0+}^q y(t) + at^{q-1}.$$

According to  $D_{0+}^{q-1} x(\infty) = \frac{\Gamma(q)}{\eta^{q-1}} x(\eta)$ , we obtain

$$\int_0^\infty y(s)ds = \frac{\Gamma(q)}{\eta^{q-1}} I_{0+}^q y(\eta).$$

On the other hand, if (3.9) holds, setting

$$x(t) = I_{0+}^q y(t) + at^{q-1},$$

where  $a$  is an arbitrary constant, then  $x(t)$  is a solution of (3.8). Hence

$$\text{Im } L = \left\{ y \in Y : \int_0^\infty y(s) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} y(s) ds = 0 \right\}.$$

Now taking (3.5) into account, it yields

$$\begin{aligned} Q^2 y = Q(Qy) &= A^2 e^{-t} \left( \int_0^\infty y(s) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} y(s) ds \right) \\ &\quad \times \left( \int_0^\infty e^{-s} ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} e^{-s} ds \right) = Qy, \end{aligned}$$

thus  $Q$  is a continuous projector and  $\text{Im } L = \ker Q$ . Rewrite  $y = (y - Qy) + Qy$ , then  $y - Qy \in \ker Q = \text{Im } L$ ,  $Qy \in \text{Im } Q$  and  $\text{Im } Q \cap \text{Im } L = \{0\}$ , then  $Y = \text{Im } L \oplus \text{Im } Q$ . Thus  $\dim \ker L = 1 = \dim \text{Im } Q = \text{co dim Im } L = 1$ , this means that  $L$  is a Fredholm operator of index zero. Now define a projector  $P$  from  $X$  to  $X$  by setting

$$Px(t) = \frac{D_{0+}^{q-1} x(0)}{\Gamma(q)} t^{q-1},$$

and the generalized inverse  $K_P: \text{Im } L \rightarrow \text{dom } L \cap \ker P$  of  $L$  as

$$K_P y(t) = I_{0+}^q y(t) = \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} y(s) ds.$$

Obviously,  $\text{Im } P = \ker L$  and  $P^2 x = Px$ . It follows from  $x = (x - Px) + Px$  that  $X = \ker P + \ker L$ . By simple calculation, we can get  $\ker L \cap \ker P = \{0\}$ . Hence  $X = \ker L \oplus \ker P$ .

Let us show that the generalized inverse of  $L$  is  $K_P$ . In fact, for  $y \in \text{Im } L$ , we have

$$(LK_P)y(t) = D_{0+}^q (K_P y(t)) = D_{0+}^q I_{0+}^q y(t) = y(t),$$

and for  $x \in \text{dom } L \cap \ker P$ , we obtain

$$\begin{aligned} (K_P L)x(t) &= (K_P) D_{0+}^q x(t) = I_{0+}^q D_{0+}^q x(t) \\ &= x(t) - \frac{D_{0+}^{q-1} x(0)}{\Gamma(q)} t^{q-1} - \frac{I_{0+}^{2-q} x(0)}{\Gamma(q-1)} t^{q-2}. \end{aligned}$$

In view of  $x \in \text{dom } L \cap \ker P$ , then  $I_{0+}^{2-q} x(0) = 0$  and  $Px = 0$ , thus

$$(K_P L)x(t) = x(t).$$

This shows that  $K_p = (L|_{\text{dom } L \cap \ker P})^{-1}$ . From the definition of  $K_p$ , we have

$$\begin{aligned} e^{-t} |(K_p y)(t)| &\leq \frac{e^{-t}}{\Gamma(q)} \int_0^t (t-s)^{q-1} |y(s)| ds \\ &\leq \frac{e^{-t} t^{q-1}}{\Gamma(q)} \int_0^\infty |y(s)| ds \leq \max_{t \geq 0} \left( \frac{e^{-t} t^{q-1}}{\Gamma(q)} \right) \|y\|_1 = M_1 \|y\|_1. \\ \left| D_{0+}^{q-1} (K_p y)(t) \right| &= |I_{0+}^1 y(t)| \leq \int_0^\infty |y(s)| ds = \|y\|_1, \end{aligned}$$

that leads to

$$\|K_p y\| \leq (M_1 + 1) \|y\|_1 = A_1 \|y\|_1.$$

This completes the proof. □

**Lemma 3.3.** *Let*

$$\Omega_1 = \{x \in \text{dom } L \setminus \ker L : Lx = \lambda Nx \text{ for some } \lambda \in [0, 1]\}.$$

*Then  $\Omega_1$  is bounded.*

*Proof.* Suppose that  $x \in \Omega_1$  and  $Lx = \lambda Nx$ . Thus  $\lambda \neq 0$  and  $QNx = 0$ , so

$$\int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds = 0.$$

Thus, by condition  $(H_2)$ , there exists  $t_0 \in \mathbb{R}_+$  such that  $|D_{0+}^{q-1} x(t_0)| \leq M$ . It follows from  $D_{0+}^q x(t) = DD_{0+}^{q-1} x(t)$  and the fact that  $D_{0+}^q x \in Y$ , that

$$\left| D_{0+}^{q-1} x(0) \right| = \left| D_{0+}^{q-1} x(t_0) - \int_0^{t_0} D_{0+}^q x(s) ds \right|.$$

Then, we have

$$\left| D_{0+}^{q-1} x(0) \right| \leq M + \int_0^\infty |Lx(s)| ds \leq M + \int_0^\infty |Nx(s)| ds = M + \|Nx\|_1. \tag{3.10}$$

On the other hand, since  $x \in \text{dom } L \setminus \ker L$ , then  $(I - P)x \in \text{dom } L \cap \ker P$  and  $LPx = 0$ , thus from Lemma 3.2, we get

$$\begin{aligned} \|(I - P)x\| &= \|K_p L(I - P)x\| \leq A_1 \|L(I - P)x\|_1 \\ &= A_1 \|Lx\|_1 \leq A_1 \|Nx\|_1. \end{aligned} \tag{3.11}$$

So

$$\|x\| \leq \|Px\| + \|(I - P)x\| = A_1 \left| D_{0+}^{q-1} x(0) \right| + A_1 \|Nx\|_1, \quad (3.12)$$

thanks to inequalities (3.10) and (3.11), (3.12) becomes

$$\|x\| \leq A_1 M + 2A_1 \|Nx\|_1. \quad (3.13)$$

On the other hand, by (3.1), we have

$$\|Nx\|_1 = \int_0^\infty |f(s, x(s)) + e(s)| ds \leq \|x\| \|\alpha\|_1 + \|\beta\|_1. \quad (3.14)$$

Thus

$$\|x\| \leq A_1 M + 2A_1 \|x\| \|\alpha\|_1 + 2A_1 \|\beta\|_1.$$

Since  $(1 - 2A_1 \|\alpha\|_1) > 0$ , we obtain

$$\|x\| \leq \frac{A_1 M}{1 - 2A_1 \|\alpha\|_1} + \frac{2A_1 \|\beta\|_1}{1 - 2A_1 \|\alpha\|_1} < \infty,$$

which proves that  $\Omega_1$  is bounded.  $\square$

**Lemma 3.4.** *The set  $\Omega_2 = \{x \in \ker L : Nx \in \text{Im } L\}$  is bounded.*

*Proof.* Let  $x \in \Omega_2$ , then  $x \in \ker L$  implies  $x(t) = ct^{q-1}$ ,  $c \in \mathbb{R}$  and  $Q Nx = 0$ , therefore

$$\int_0^\infty (f(s, cs^{q-1}) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, cs^{q-1}) + e(s)) ds = 0.$$

From condition  $(H_2)$  there exists  $t_1 \geq 0$  such that  $\left| D_{0+}^{q-1} x(t_1) \right| \leq M$  and so  $|c| \leq \frac{M}{\Gamma(q)}$ . On the other hand,

$$\begin{aligned} \|x\| &= \sup_{t \geq 0} e^{-t} |x(t)| + \sup_{t \geq 0} \left| D_{0+}^{q-1} x(t) \right| \leq |c| \left( \sup_{t \in [0, \infty)} e^{-t} t^{q-1} + \Gamma(q) \right) \\ &\leq |c| \Gamma(q) A_1 \leq M A_1 < \infty, \end{aligned}$$

so  $\Omega_2$  is bounded.  $\square$

**Lemma 3.5.** *Suppose that the first part of Condition  $(H_3)$  of Theorem 3.1 holds. Let*

$$\Omega_3 = \{x \in \ker L : -\lambda Jx + (1 - \lambda) Q Nx = 0, \lambda \in [0, 1]\},$$

where  $J : \ker L \rightarrow \text{Im } Q$  is the linear isomorphism given by  $J(ct^{q-1}) = ce^{-t}$  for all  $c \in \mathbb{R}$ ,  $t \geq 0$ . Then  $\Omega_3$  is bounded.



*Proof.* Let  $x_0 \in \Omega_3$ , then  $x_0(t) = c_0 t^{q-1}$  and  $\lambda Jx_0 = (1 - \lambda) QN x_0$  that is equivalently written as

$$\lambda c_0 = (1 - \lambda) A \left( \int_0^\infty (f(s, c_0 s^{q-1}) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, c_0 s^{q-1}) + e(s)) ds \right).$$

If  $\lambda = 1$ , then  $c_0 = 0$ . Otherwise, if  $|c_0| > M^*$ , in view of (3.3) one has

$$\lambda c_0^2 = (1 - \lambda) A c_0 \left( \int_0^\infty (f(s, c_0 s^{q-1}) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, c_0 s^{q-1}) + e(s)) ds \right) < 0,$$

which contradicts the fact that  $\lambda c_0^2 \geq 0$ . So  $|c_0| \leq M^*$ , moreover

$$\|x_0\| = |c_0| \left( \sup_{t \geq 0} e^{-t} t^{q-1} + \Gamma(q) \right) \leq |c_0| \Gamma(q) A_1 \leq M^* \Gamma(q) A_1,$$

thus  $\Omega_3$  is bounded. □

**Lemma 3.6.** *Under the second part of Condition  $(H_3)$  of Theorem 3.1, the set  $\Omega_3$  is bounded.*

*Proof.* The proof is similar to the one of Lemma 3.5. □

**Lemma 3.7.** *Suppose that  $\Omega$  is an open bounded subset of  $X$  such that  $\text{dom } L \cap \overline{\Omega} \neq \emptyset$ . Then  $N$  is  $L$ -compact on  $\overline{\Omega}$ .*

*Proof.* Suppose that  $\Omega \subset X$  is a bounded set. Without loss of generality, we may assume that  $\Omega = B(0, r)$ , then for any  $x \in \overline{\Omega}$ ,  $\|x\| \leq r$ . For  $x \in \overline{\Omega}$ , and by condition (3.1), we obtain

$$\begin{aligned}
|QNx| &\leq Ae^{-t} \left( \int_0^\infty |f(s, x(s)) + e(s)| ds \right. \\
&\quad \left. + \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} |f(s, x(s)) + e(s)| ds \right) \\
&\leq Ae^{-t} \left[ \int_0^\infty e^{-s} \alpha(s) |x(s)| + \beta(s) ds \right. \\
&\quad \left. + \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (e^{-s} \alpha(s) |x(s)| + \beta(s)) ds \right] \\
&\leq Ae^{-t} \left( r \int_0^\infty \alpha(s) ds + \int_0^\infty \beta(s) ds + r \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} \alpha(s) ds \right. \\
&\quad \left. + \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} \beta(s) ds \right) \\
&\leq Ae^{-t} (1 + \Gamma(q)) (r \|\alpha\|_1 + \|\beta\|_1).
\end{aligned}$$

Thus

$$\|QNx\|_1 \leq A(1 + \Gamma(q)) (r \|\alpha\|_1 + \|\beta\|_1), \quad (3.15)$$

which implies that  $QN(\overline{\Omega})$  is bounded. Next, we show that  $K_P(I - Q)N(\overline{\Omega})$  is compact. For  $x \in \overline{\Omega}$ , by (3.1) we have

$$\|Nx\|_1 = \int_0^\infty |f(s, x(s)) + e(s)| ds \leq r \|\alpha\|_1 + \|\beta\|_1. \quad (3.16)$$

On the other hand, from the definition of  $K_P$  and together with (3.6), (3.15) and (3.16) one gets

$$\begin{aligned}
\|K_P(I - Q)Nx\| &\leq A_1 \|(I - Q)Nx\|_1 \leq A_1 [\|Nx\|_1 + \|QNx\|_1] \\
&\leq A_1 (1 + A(1 + \Gamma(q))) (r \|\alpha\|_1 + \|\beta\|_1).
\end{aligned}$$

It follows that  $K_P(I - Q)N(\overline{\Omega})$  is uniformly bounded.

Let us prove that  $K_P(I - Q)N(\overline{\Omega})$  is equicontinuous. For any  $x \in \overline{\Omega}$ , and any  $t_1, t_2 \in [0, T]$ ,  $t_1 < t_2$  with  $T > 0$ , we have

$$\begin{aligned} & \left| e^{-t_1} (K_P(I - Q)Nx)(t_1) - e^{-t_2} (K_P(I - Q)Nx)(t_2) \right| = \\ & = \frac{1}{\Gamma(q)} \left| \int_0^{t_1} e^{-t_1} (t_1 - s)^{q-1} (I - Q)Nx(s) ds \right. \\ & \quad \left. - \int_0^{t_2} e^{-t_2} (t_2 - s)^{q-1} (I - Q)Nx(s) ds \right| \\ & \leq \frac{1}{\Gamma(q)} \left[ \int_0^{t_1} \left( e^{-t_2} (t_2 - s)^{q-1} - e^{-t_1} (t_1 - s)^{q-1} \right) |(I - Q)Nx(s)| ds \right. \\ & \quad \left. + \int_{t_1}^{t_2} e^{-t_2} (t_2 - s)^{q-1} |(I - Q)Nx(s)| ds \right] \\ & \leq \frac{1}{\Gamma(q)} \left[ e^{-t_1} \left( t_2^{q-1} - t_1^{q-1} \right) \int_0^{t_1} |(I - Q)Nx(s)| ds \right. \\ & \quad \left. + e^{-t_2} t_2^{q-1} \int_{t_1}^{t_2} |(I - Q)Nx(s)| ds \right] \rightarrow 0, \end{aligned}$$

as  $t_1 \rightarrow t_2$ . On the other hand, we have

$$\begin{aligned} & \left| e^{-t_1} D_{0+}^{q-1} (K_P(I - Q)Nx)(t_1) - e^{-t_2} D_{0+}^{q-1} (K_P(I - Q)Nx)(t_2) \right| \\ & = \left| \int_0^{t_1} e^{-t_1} (I - Q)Nx(s) ds - \int_0^{t_2} e^{-t_2} (I - Q)Nx(s) ds \right| \\ & \leq \int_0^{t_1} (e^{-t_1} - e^{-t_2}) |(I - Q)Nx(s)| ds + \int_{t_1}^{t_2} e^{-t_2} |(I - Q)Nx(s)| ds \\ & \leq (t_2 - t_1) \int_0^{t_1} |(I - Q)Nx(s)| ds + \int_{t_1}^{t_2} |(I - Q)Nx(s)| ds \rightarrow 0, \end{aligned}$$

as  $t_1 \rightarrow t_2$ . So  $K_P(I - Q)N(\overline{\Omega})$  is equicontinuous on every compact subinterval of  $[0, \infty)$ . In addition, we claim that  $K_P(I - Q)N(\overline{\Omega})$  is equiconvergent at infinity. In fact,

$$\left| e^{-t} (K_P(I - Q)Nx)(t) \right| \leq \frac{1}{\Gamma(q)} \int_0^t e^{-t} (t - s)^{q-1} |(I - Q)Nx(s)| ds$$

$$\begin{aligned} &\leq \frac{e^{-t}t^{q-1}}{\Gamma(q)} \int_0^t |(I - Q)Nx(s)| ds \leq \frac{e^{-t}t^{q-1}}{\Gamma(q)} \|(I - Q)Nx\|_1 \\ &\leq \frac{e^{-t}t^{q-1}}{\Gamma(q)} (\|Nx\|_1 + \|QNx\|_1) \leq \frac{e^{-t}t^{q-1}}{\Gamma(q)} (1 + 2A) [r\|\alpha\|_1 + \|\beta\|_1]. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} &\left| e^{-t}D_{0+}^{q-1}(K_P(I - Q)Nx)(t) \right| \leq e^{-t} \int_0^t |(I - Q)Nx(s)| ds \\ &\leq e^{-t} (\|Nx\|_1 + \|QNx\|_1) \leq e^{-t} (1 + A(1 + \Gamma(q))) (r\|\alpha\|_1 + \|\beta\|_1). \end{aligned}$$

Thus

$$\lim_{t \rightarrow \infty} e^{-t}(K_P(I - Q)Nx)(t) = 0, \lim_{t \rightarrow \infty} e^{-t}D_{0+}^{q-1}(K_P(I - Q)Nx)(t) = 0,$$

consequently  $K_P(I - Q)N(\overline{\Omega})$  is equiconvergent at  $\infty$ . □

Now we are able to give the proof of Theorem 3.1.

*Proof of Theorem 3.1.* In what follows, we shall prove that all conditions of Theorem 2.1 are satisfied. Let  $\Omega$  to be an open bounded subset of  $X$  such that  $\cup_{i=1}^3 \overline{\Omega}_i \subset \Omega$ . From Lemma 3.2, we know that  $L$  is a Fredholm operator of index zero. By Lemma 3.7, we have  $N$  is  $L$ -compact on  $\overline{\Omega}$ . By virtue of the definition of  $\Omega$ , it yields

- (i)  $Lx \neq \lambda Nx$  for all  $(x, \lambda) \in [(\text{dom } L \setminus \ker L) \cap \partial\Omega] \times (0, 1)$ ,
- (ii)  $Nx \notin \text{Im } L$  for all  $x \in \ker L \cap \partial\Omega$ .

Now we prove that condition (iii) of Theorem 2.1 is satisfied. Let  $H(x, \lambda) = \pm\lambda Jx + (1 - \lambda)QNx$ . Since  $\overline{\Omega}_3 \subset \Omega$ , then  $H(x, \lambda) \neq 0$  for every  $x \in \ker L \cap \partial\Omega$ . By the homotopy property of degree, we get

$$\begin{aligned} \deg(QN|_{\ker L}, \Omega \cap \ker L, 0) &= \deg(H(\cdot, 0), \Omega \cap \ker L, 0) \\ &= \deg(H(\cdot, 1), \Omega \cap \ker L, 0) \\ &= \deg(\pm J, \Omega \cap \ker L, 0) \neq 0. \end{aligned}$$

So, the third assumption of Theorem 2.1 is fulfilled and  $Lx = Nx$  has at least one solution in  $\text{dom } L \cap \overline{\Omega}$ , i.e. (1.1)–(1.2) has at least one solution in  $X$ . The proof is completed. □

**Example 3.8.** Consider the fractional boundary value problem

$$\begin{cases} D_{0+}^q x(t) = f(t, x(t)) + e(t), & t \in (0, \infty), \\ I_{0+}^{2-q} x(0) = 0, \quad D_{0+}^{q-1} x(\infty) = \frac{\Gamma(q)}{\eta^{q-1}} x(\eta), \end{cases} \tag{P}$$

where  $q = \frac{3}{2}$ ,  $f(t, x) = \frac{1}{4}e^{-t}(e^{-t}|x| - 8e^{-t})$ ,  $e(t) = 3e^{-t}$ . Choosing  $\alpha(t) = \frac{1}{4}e^{-t}$  and  $\beta(t) = 3e^{-t} - 2e^{-2t}$ , then  $\alpha, \beta$  are nonnegative and belong to  $L^1[0, \infty)$ . For all  $x \in \mathbb{R}$ ,  $t \in [0, \infty)$  we have

$$|f(t, x) + e(t)| \leq e^{-t}\alpha(t)|x| + \beta(t),$$

so hypothesis  $(H_1)$  of Theorem 3.1 is satisfied. We claim that condition  $(H_3)$  is satisfied, indeed, since

$$\begin{aligned} & \int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \\ & \geq \int_\eta^\infty (f(s, x(s)) + e(s)) ds, \end{aligned}$$

then for  $M^* = 8$ ,  $\eta = 1$  and any  $x(t) = c_0 t^{q-1} \in \ker L$  with  $|c_0| > M^*$ , we have

$$\begin{aligned} & \int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \\ & \geq \int_1^\infty \left( \frac{1}{4} e^{-s} \left( e^{-s} |c_0 s^{\frac{3}{2}-1}| - 8e^{-s} \right) + 3e^{-s} \right) ds \\ & \geq \int_1^\infty (2e^{-2s} \sqrt{s} - 2e^{-s} + 3e^{-s}) ds = 0.53173 > 0. \end{aligned}$$

As a result, if  $c_0 > 0$ , then

$$c_0 \left[ \int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \right] > 0,$$

or if  $c_0 < 0$ , then

$$c_0 \left[ \int_0^\infty (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^\eta (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \right] < 0.$$

Now, we have

$$\begin{aligned}
& \int_0^{\infty} (f(s, x(s)) + e(s)) ds - \frac{\Gamma(q)}{\eta^{q-1}} \int_0^{\eta} (\eta - s)^{q-1} (f(s, x(s)) + e(s)) ds \\
&= \int_0^{\infty} \left( \frac{1}{4} e^{-s} (e^{-s} |x(s)| - 8e^{-s}) + 3e^{-s} \right) ds \\
&\quad - \Gamma \left( \frac{3}{2} \right) \int_0^1 (1-s)^{\frac{3}{2}-1} \left( \frac{1}{4} e^{-s} (e^{-s} |x(s)| - 8e^{-s}) + 3e^{-s} \right) ds \\
&\geq \int_1^{\infty} \frac{1}{4} e^{-s} (e^{-s} |x(s)| - 8e^{-s}) + 3e^{-s} ds \geq \int_1^{\infty} (3e^{-s} - 2e^{-2s}) ds \\
&= 0.9683 \neq 0,
\end{aligned}$$

consequently,  $(H_2)$  is satisfied for any constant  $M > 0$ . Finally, a simple calculus gives

$$(1 - 2A_1 \|\alpha\|_1) = 0.35324 > 0,$$

where

$$A_1 = 1 + \frac{[e^{-1}(q-1)]^{q-1}}{\Gamma(q)} = 1.2935 \quad \text{and} \quad \|\alpha\|_1 = \int_0^{\infty} \frac{1}{4} e^{-t} ds = \frac{1}{4}.$$

We conclude from Theorem 3.1 that the problem (P) has at least one solution in  $X$ .

### Acknowledgments

The authors would like to thank the referee for his (her) helpful comments and suggestions.

### REFERENCES

- [1] R.P. Agarwal, M. Benchohra, S. Hamani, S. Pinelas, *Boundary value problems for differential equations involving Riemann-Liouville fractional derivative on the half line*, Dyn. Contin. Discrete Impuls. Syst. Ser. A. Math. Anal. **18** (2011) 2, 235–244.
- [2] R.P. Agarwal, D. O'Regan, *Infinity Interval Problems for Difference and Integral Equations*, Kluwer Academic Publisher Dordrecht, 2001.
- [3] R.P. Agarwal, D. O'Regan, *Infinite interval problems modeling phenomena which arise in the theory of plasma and electrical potential theory*, Stud. Appl. Math. **111** (2003) 3, 339–358.
- [4] A. Arara, M. Benchohra, N. Hamidi, J.J. Nieto, *Fractional order differential equations on an unbounded domain*, Nonlinear Anal. **72** (2010), 580–586.

- [5] Y. Chen, X. Tang, *Positive solutions of fractional differential equations at resonance on the half-line*, Boundary Value Problems (2012) 2012:64, 13 pp.
- [6] C. Corduneanu, *Integral Equations and Stability of Feedback Systems*, Academic Press, New York, 1973.
- [7] Y. Cui, *Solvability of second-order boundary-value problems at resonance involving integral conditions*, Electron. J. Differ. Equ. (2012) 2012:45, 9 pp.
- [8] W. Feng, J.R.L. Webb, *Solvability of three-point boundary value problems at resonance*, Nonlinear Anal. Theory, Methods and Appl. **30** (1997), 3227–3238.
- [9] D. Franco, G. Infante, M. Zima, *Second order nonlocal boundary value problems at resonance*, Math. Nachr. **284** (2011) 7, 875–884.
- [10] A. Guezane-Lakoud, A. Kilickman, *Unbounded solution for a fractional boundary value problem*, Advances in Difference Equations (2014) 2014:154.
- [11] C.P. Gupta, S.K. Ntouyas, P.Ch. Tsamatos, *On an  $m$ -point boundary-value problem for second-order ordinary differential equations*, Nonlinear Anal. **23** (1994), 1427–1436.
- [12] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier Science, Amsterdam, vol. 204, 2006.
- [13] N. Kosmatov, *Multi-point boundary value problems on an unbounded domain at resonance*, Nonlinear Anal. **68** (2008) 8, 2158–2171.
- [14] R. Ma, *Existence of positive solutions for second-order boundary value problems on infinity intervals*, Appl. Math. Lett. **16** (2003) 1, 33–39.
- [15] J. Mawhin, *Topological degree methods in nonlinear boundary value problems*, NSFCBMS Regional Conference Series in Mathematics, Am. Math. Soc, Providence, 1979.
- [16] D. O'Regan, B. Yan, R.P. Agarwal, *Solutions in weighted spaces of singular boundary value problems on the half-line*, J. Comput. Appl. Math. **205** (2007), 751–763.
- [17] X. Su, S. Zhang, *Unbounded solutions to a boundary value problem of fractional order on the half-line*, Comput. Math. Appl. **61** (2011), 1079–1087.

Assia Frioui  
frioui.assia@yahoo.fr

University Guelma  
Laboratory of Applied Mathematics and Modeling  
P.O. Box 401, Guelma 24000, Algeria

Assia Guezane-Lakoud  
a\_guezane@yahoo.fr

University Badji Mokhtar-Annaba  
Faculty of Sciences  
Laboratory of Advanced Materials  
P.O. Box 12, 23000, Annaba, Algeria

Rabah Khaldi  
rkhadi@yahoo.fr

University Badji Mokhtar-Annaba  
Faculty of Sciences  
Laboratory of Advanced Materials  
P.O. Box 12, 23000, Annaba, Algeria

*Received: March 11, 2016.*

*Revised: May 13, 2016.*

*Accepted: June 2, 2016.*