



## Characterization and Profiling of Liver microRNAs by RNA-sequencing in Cattle Divergently Selected for Residual Feed Intake

Wijdan Al-Husseini\*, Yizhou Chen<sup>1,\*</sup>, Cedric Gondro, Robert M. Herd<sup>2</sup>,  
John P. Gibson, and Paul F. Arthur<sup>1</sup>

The Centre for Genetics Analysis and Applications, School of Environmental and Rural Science,  
University of New England, Armidale, NSW 2351, Australia

**ABSTRACT:** MicroRNAs (miRNAs) are short non-coding RNAs that post-transcriptionally regulate expression of mRNAs in many biological pathways. Liver plays an important role in the feed efficiency of animals and high and low efficient cattle demonstrated different gene expression profiles by microarray. Here we report comprehensive miRNAs profiles by next-gen deep sequencing in Angus cattle divergently selected for residual feed intake (RFI) and identify miRNAs related to feed efficiency in beef cattle. Two microRNA libraries were constructed from pooled RNA extracted from livers of low and high RFI cattle, and sequenced by Illumina genome analyser. In total, 23,628,103 high quality short sequence reads were obtained and more than half of these reads were matched to the bovine genome (UMD 3.1). We identified 305 known bovine miRNAs. *Bta-miR-143*, *bta-miR-30*, *bta-miR-122*, *bta-miR-378*, and *bta-let-7* were the top five most abundant miRNAs families expressed in liver, representing more than 63% of expressed miRNAs. We also identified 52 homologous miRNAs and 10 novel putative bovine-specific miRNAs, based on precursor sequence and the secondary structure and utilizing the miRBase (v. 21). We compared the miRNAs profile between high and low RFI animals and ranked the most differentially expressed bovine known miRNAs. Bovine *miR-143* was the most abundant miRNA in the bovine liver and comprised 20% of total expressed mapped miRNAs. The most highly expressed miRNA in liver of mice and humans, *miR-122*, was the third most abundant in our cattle liver samples. We also identified 10 putative novel bovine-specific miRNA candidates. Differentially expressed miRNAs between high and low RFI cattle were identified with 18 miRNAs being up-regulated and 7 other miRNAs down-regulated in low RFI cattle. Our study has identified comprehensive miRNAs expressed in bovine liver. Some of the expressed miRNAs are novel in cattle. The differentially expressed miRNAs between high and low RFI give some insights into liver miRNAs regulating physiological pathways underlying variation in this measure of feed efficiency in bovines. (**Key Words:** Bovine, Feed Efficiency, Gene Expression, miRNAs, Next Generation Sequencing)

### INTRODUCTION

MicroRNAs (miRNAs) are small (~22 nucleotides) non-coding RNA that regulate gene expression by target

messenger RNA (mRNA) in a sequence-specific manner, leading to either translational repression or degradation of targeted transcript. In animals, miRNAs target the 3'untranslated regions of mRNA by an RNA-induced silencing complex (RISC), and subject to the accuracy of the sequence complementarities, either repression of translation or cleavage of the mRNA target is achieved (Yu et al., 2007). MicroRNAs are now known to repress thousands of target genes and regulate cellular processes, including cellular proliferation, differentiation and apoptosis. The aberrant expression or alteration of miRNAs also contributes to a range of human pathologies, including diabetes and cancer (Lu et al., 2005).

\* Corresponding Authors: Wijdan Al-Husseini. Tel: +61-2-6773-2155, E-mail: wabedaliune@gmail.com / Yizhou Chen. Tel: +61-2-4640-6453, Fax: +61-2-4640-6300, E-mail: yizhou.chen@dpi.nsw.gov.au

<sup>1</sup> NSW Department of Primary Industries, Elizabeth Macarthur Agricultural Institute, Menangle, NSW 2568, Australia.

<sup>2</sup> NSW Department of Primary Industries, Beef Industry Centre, Armidale, NSW 2351, Australia.

Submitted Jul. 19, 2015; Revised Oct. 10, 2015; Accepted Dec. 14, 2015

Feed efficiency is an economically important trait in beef production and can be measured as residual feed intake (RFI). This is the difference between an animal's actual feed intake recorded over a test period and its expected feed intake based on its size and growth rate (Koch et al., 1963). RFI takes into consideration feed required for daily weight gain of the animal as well as for maintenance of its metabolic weight; therefore, understanding of the molecular mechanism regulated RFI will help breeding sustainable and profitable animals in agriculture. Genome wide association studies have been carried out to identify gene markers associated with RFI in beef cattle (Sherman et al., 2009) and more than a hundred single nucleotide polymorphic (SNP) markers have been found associated with variation in RFI in beef cattle. However, a large proportion of SNPs markers are not located in annotated genes in bovine genome. Some of the most significant SNPs for RFI were in miRNAs motifs which suggest that these miRNAs could play an important role in regulation of RFI. Gene expression studies in cattle from high and low RFI divergent selection lines identified more than 160 differentially expressed genes (Chen et al., 2011). The variations of gene expression between high and low RFI cattle ranged from -1.5 to 0.8 fold change, and support the view that the phenotypic differences in RFI may due to level of gene expression instead of genes being switched on or off.

Profiling studies characterizing miRNAs encoded in livestock genomes in the last decade have found a wide and diverse range of conserved and species-specific miRNAs (Liu et al., 2009). In cattle, some initial characterizations of miRNAs have been carried out for various tissues including adipose tissue, mammary tissues (Gu et al., 2007), immune and embryonic tissues, pooled tissue, alveolar macrophages, ovarian, oocyte and testicular tissues (Miles et al., 2012), liver of dairy cows in the early postpartum period (Fatima et al., 2014). Liver is a central controller of metabolism and a major driver of whole animal oxygen consumption in mammals. However, little is known about miRNAs role in regulating key cellular and physiological pathways that may regulate RFI. In the current study, we profiled miRNAs abundance in liver tissue of Angus bulls from high and low RFI-selection lines by next-generation deep sequencing technology. We report liver miRNA-seq profiling study in beef cattle and their known and putative novel bovine miRNAs. We also report the differentially expressed miRNAs between high and low RFI-selection line cattle.

## MATERIALS AND METHODS

### Animal and liver biopsy sampling

Young Angus bulls resulting from approximately three generations of divergent selection for RFI were used in this

study. The selection lines were established in 1993 at the Agricultural Research Centre, Trangie, NSW, Australia (Arthur et al., 2001; Donoghue et al., 2011). The same animals were used for the microarray experiment reported by Chen et al. (2011), and were approved by the University of New England Animal Ethics Committee, certificate no. (AEC 06/123) and followed the University of New England Code of Conduct for Research in meeting the Australian Code of Practice for the Care and Use of Animals. In brief, bulls were born in 2005 and, when approximately one year-of-age, their growth and feed intake were measured. Post-weaning RFI for each animal in the test group was calculated using a linear regression of daily feed intake on mean metabolic mid-test weight and average daily gain (Arthur et al., 2001). Based on the RFI test results, liver biopsies were taken (7 days after the end of the RFI test) from 30 animals with the lowest RFI and 30 animals with the highest RFI (Chen et al., 2011). On average, cattle from the low RFI line consumed about 2 kg less feed per day than cattle from the high RFI line ( $p < 0.001$ ; as shown in Table 1). Total RNA from liver was isolated using TRI reagent (Ambion, Applied Biosystems, Foster City, CA, USA) following the manufacturer's protocol. The concentration and purity of the extracted RNA was checked by NanoDrop ND-1000 Spectrophotometer (NanoDrop Technologies Inc., Wilmington, DE, USA). The quality and integrity of RNA was further assessed with the RNA 6000 Nano Lab chip Kit using the Agilent 2100 Bioanalyzer (Agilent Technologies, USA). All RNA samples had a RNA integrity number (RIN) of 9.8 to 10) and were stored at  $-80^{\circ}\text{C}$  until further analysis.

### Small RNA library construction and sequencing

For small RNA library construction and deep sequencing, two pools of total RNA were constructed from 13 top high-RFI animals and 13 bottom low-RFI animals with equal quantities (1  $\mu\text{g}$  of RNA) from each animal. Libraries of small RNAs were prepared using the "Preparing Samples for Small RNA Sequencing Using the Alternative v1.5 Protocol" supplied by the manufacturer and using 10  $\mu\text{g}$  total RNA. The libraries were sequenced on the Illumina Genome Analyzer Iix using Single-Read

**Table 1.** Feed efficiency performance of young Angus bulls selected for and against residual feed intake (RFI)

Traits	Mean $\pm$ SE	
	Low RFI	High RFI
Residual feed intake (kg/d)	-1.34 $\pm$ 0.51	0.56 $\pm$ 0.28
Feed intake (kg/d)	10.45 $\pm$ 1.19	12.57 $\pm$ 0.97
Average daily gain (kg/d)	2.09 $\pm$ 0.30	1.94 $\pm$ 0.21
Start body weight (kg)	324 $\pm$ 29	357 $\pm$ 29
End body weight (kg)	471 $\pm$ 41	493 $\pm$ 40

SE, standard error.

Cluster Generation Kit v2 (Cat. no. FC-103–2001; Illumina) and 36 Cycle Sequencing Kit v4 (Cat. no. FC-104–4002). All sequencing data sets supporting the results in this study have been deposited in the publicly available NCBI's Gene Expression Omnibus Database (<http://www.ncbi.nlm.nih.gov/geo/>). The data are accessible through GEO Series accession number GSE63691 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE63691>).

#### Analysis of small RNA sequencing data

The raw sequence reads were processed with a Perl script to remove low quality reads and adaptor sequences and to count the reads of the Genome Analyzer (Illumina Inc., San Diego, CA, USA) into read-count files (read sequences and counts tab separated). Then the sequencing data were input into miRanalyzer (Hackenberg et al., 2011) to analyse the small RNAs. In brief, known bovine microRNAs were identified by mapping all sequence reads to the known bovine miRNAs database (miRBase v. 21) (Kozomara and Griffiths-Jones, 2011). Those reads matched to known bovine miRNAs were grouped and removed from the dataset so they could not erroneously be predicted as new miRNAs. Then, sequences reads mapped to non-bovine known miRNAs to identify homologous miRNAs for all other species in the miRBase. Genome sequences around the position of those mapped reads were extracted and the energetically best hairpin structures were retained as putative pre-miRNAs if they had: i) at least 19 base pairings in the secondary structure; and ii) at least 11 base pairings located in the read cluster region (number of pairings between putative mature and mature-star). Mapped reads were grouped as novel bovine homologue miRNAs and removed from data set. The remaining reads were further successively aligned to known transcriptome libraries such as RFam 10.1, GtRNAdb, RNAdb, and RefSeq (Gardner et al., 2009). To identify bovine-specific novel miRNAs, the remaining sequence reads were mapped to *Bos taurus* genome (BosTau6, UMD\_3.1) using Bowtie (Langmead et al., 2009). The mapped reads were first clustered into putative mature miRNAs and pre-miRNAs. Putative candidate miRNAs reported based on at least three out of five models from the five different Random Forest models (Hackenberg et al., 2011). The secondary structure of pre-miRNAs was determined by RNAfold using minimum free energy (MFE) algorithm.

#### Differential expression analysis of miRNAs

The raw read counts of known bovine miRNAs and homologous miRNAs were imported as a count table into the Bioconductor DESeq package (Anders and Huber, 2010), which uses the negative binomial distribution model to test for differential expression in deep sequencing datasets. The normalization and variance estimation were

based on the model of two conditions without replicate.

#### Prediction of putative miRNA targets

Potential gene targets for known miRNAs that differentially expressed between high and low RFI were identified using miRmap (Vejnar and Zdobnov, 2012). Only miRNAs target genes that highly scores (>80%) were selected for investigation as potential target genes. Also, all those genes expressed differentially between high and low RFI-selection line cattle which have been reported by Chen et al. (2011) were investigated for their potential miRNAs targets in the 3'UTR region.

## RESULTS AND DISCUSSION

#### Profiling bovine miRNAs


In order to profile and identify novel and differentially-expressed miRNAs in the bovine liver of low and high RFI cattle, two small RNA libraries were constructed by Solexa sequencing. After trimming the adaptor sequence and filtering low-quality sequences, a total of 23,628,109 sequences of short reads were obtained for low and high RFI libraries. As presented in Table 2, there were 10,820,087 sequences reads for the low RFI library (high feed efficiency cattle) and 12,808,022 sequence reads for the high RFI (low feed efficiency cattle). All sequence reads were aligned against the bovine miRNAs database (miRBase v.21). Importantly, high proportion 74.5% and 83.7% of sequence reads of the two RFI libraries have aligned to the bovine genome (UMD 3.1). Over half of each library reads (56% to 57%) have showed an exact match to the known bovine mature miRNAs, as shown in Table 2. These results demonstrated the high quality of our small RNAs sequencing read libraries. However, the presence of other small RNAs, such as ribosomal fragments, tRNA, snRNA, and mtRNA was approximately 12% and 17% of mapped sequence reads of low and high RFI libraries respectively. Sequence reads that matched to homologous miRNAs of other species reported in the miRBase

**Table 2.** Summary of miRNA sequences present in low and high RFI libraries

Parameter	Low RFI	High RFI
Initial high quality read count	10,820,087	12,808,022
Match to known bovine mature miRNAs	4,493,321	6,151,626
Matched to mature Homolog	114,603	103,465
Matched to known transcript libraries <sup>1</sup>	987,892	1,855,305
Matched to bovine genome	2,461,668	2,614,666
Not matched to bovine genome	2,762,603	2,082,960

RFI, residual feed intake.

<sup>1</sup> rRNA, tRNA, snRNA, and mtRNA.



**Figure 1.** Distribution of high quality reads that mapped to the bovine genome in the low RFI line (A) and high RFI line (B) cattle. RFI, residual feed intake.

constituted 1% of mapped reads (Figure 1).

### Unique known mature bovine miRNAs

In total 305 known mature miRNAs detected in two libraries and the full profile of unique bovine mature miRNAs expressed in liver tissue was listed in Additional files, Supplementary Table S1. Table 3 lists only the most abundant miRNAs (>100 read count per million mapped read [RCPM]) from 305 known mature miRNAs. About 40% (305 out of 807) of known bovine mature miRNAs were expressed in the liver tissue. The proportion of expressed miRNAs in bovine liver is similar to the level of expressed miRNAs in human and mouse liver (Kornfeld et al., 2013). In contrast, primary bovine mammary epithelial cells expressed only 20% of known bovine miRNA (Lawless et al., 2013).

A very broad range in expression levels of miRNAs was found ranged from 137,000 RCPM to less than one RCPM (Additional files, Supplementary Table S1). Only a small number of miRNAs were expressed at very high levels, such as *bta-miR-143*, *bta-miR-30a*, and *bta-miR-122*, and the majority of miRNAs being expressed at medium and low levels. However, the top 10 most abundantly expressed miRNAs families (*bta-miR-143*, *bta-miR-30*, *bta-miR-122*, *bta-miR-378*, *bta-let-7*, *bta-miR-148*, *bta-miR-192*, *bta-miR-101*, *bta-miR-140*, and *bta-miR-21*) constituted approximately 78.4% of the total mapped mature known bovine sequence reads. These findings are in agreement with studies of miRNAs profiling which revealed that small number of abundant miRNAs make up a high portion of the total miRNAs pool (Tripurani et al., 2010; Fatima et al., 2014).

Among the highly expressed known miRNAs detected in the bovine liver in our study, *miR-143* was the most abundant miRNA, constituting approximately 25% of the total expressed miRNAs. In human and mouse liver the most abundant miRNA is *miR-122* (Hu et al., 2012; Rottiers and Näär 2012), which makes up 70% of the total mouse

liver miRNA population (Wen and Friedman, 2012). Our results showed that bovine *bta-miR-122* was the third most expressed miRNA, comprising approximately 8% of the total known bovine miRNA population.

Previously, tissue-specific miRNAs were reported in dairy and beef cattle (Jin et al., 2009; Tripurani et al., 2010) using cloning and sequencing small-RNA libraries. One of those tissue-specific miRNAs was *bta-miR-9*. Bovine *miR-9* was identified and reported as brain-specific miRNA (Jin et al., 2009). Similarly, *bta-miR-1*, *bta-miR-133a*, and *bta-miR-206* have been reported as muscle-specific miRNAs in bovine (Jin et al., 2009). In the current study, these four miRNAs have expressed in bovine liver with sensible expression levels in the two high and low RFI libraries. The expression levels of *bta-miR-1*, *bta-miR-206*, and *bta-miR-133a* were 7,081, 1,450, and 689 RCPM respectively. These results suggest that previous reports on tissue specificity of these miRNAs might be limited by the tissues surveyed and/ or sensitivity of the techniques that used.

### Novel bovine miRNA


*Novel isomiRs*: variations with respect to the reference miRNA sequence (isomiRs) were named previously as mature-star (miRNA\*), and were considered as “minor” products from pre-microRNA sequence. In miRNAs biogenesis, it was often inferred to be degraded and only the dominant arm (sense strand) to be incorporated into the RISC complex. However, there is strong evidence that these isomiRs are heterogeneous variants of canonical miRNAs species and are of functional importance (Yang et al., 2011). We adopted the new nomenclature using -5'p/-3'p for the name of isomiRs miRNA in the bovine library (Figure 2). Fifty novel isomiRs were detected in the present study, which have not been observed previously in the bovine miRNA database. Four isomiRs, *bta-mir-143-3p*, *bta-mir-335-3p*, *bta-mir-136-3p*, and *bta-mir-2284w-3p* have identified to have higher read counts than the corresponding known mature miRNA reported in the bovine miRBase, as

**Table 3.** The most abundant bovine mature miRNAs in the high and low RFI sequence libraries from liver tissue

miRNA ID	Low RFI library		High RFI library		Average of two libraries	miRNA ID	Low RFI library		High RFI library		Average of two libraries
	RC	RCPM	RC	RCPM			RC	RCPM	RC	RCPM	
<i>bta-miR-143</i>	692,736	85,974.2	1,883,457	175,612.7	137,158.9	<i>bta-let-7e</i>	7,633	947.3	3,207	299	577.1
<i>bta-miR-30a-5p</i>	539,136	66,911.2	859,054	80,097.8	74,440.9	<i>bta-miR-99b</i>	7,465	926.5	3,224	300.6	569.1
<i>bta-miR-122</i>	535,909	66,510.7	287,465	26,803.1	43,837.2	<i>bta-miR-181a</i>	7,101	881.3	3,423	319.2	560.3
<i>bta-miR-378</i>	341,772	42,416.7	30,5081	28,445.6	34,439	<i>bta-miR-27a-3p</i>	5,065	628.6	5,040	469.9	538
<i>bta-let-7f</i>	277,021	34,380.6	285,630	26,632	29,956.1	<i>bta-miR-151-5p</i>	5,738	712.1	4,270	398.1	532.8
<i>bta-miR-148a</i>	159,891	19,843.8	346,010	32,261.8	26,934.6	<i>bta-miR-379</i>	4,125	511.9	4,922	458.9	481.7
<i>bta-let-7a-5p</i>	259,842	32,248.5	172,859	16,117.3	23,037.4	<i>bta-miR-22-3p</i>	4,695	582.7	3,987	371.7	462.2
<i>bta-miR-192</i>	106,160	13,175.3	309,352	28,843.8	22,122.2	<i>bta-miR-125a</i>	4,910	609.4	2,808	261.8	410.9
<i>bta-miR-101</i>	55,479	6,885.4	244,427	22,790.3	15,967.3	<i>bta-miR-146a</i>	2,299	285.3	4,752	443.1	375.4
<i>bta-miR-140</i>	98,925	12,277.4	81,434	7,592.9	9,602.5	<i>bta-miR-16a</i>	3,520	436.9	3,478	324.3	372.6
<i>bta-miR-21-5p</i>	39,273	4,874.1	139,116	12,971.1	9,497.6	<i>bta-miR-2904</i>	4,655	577.7	2,305	214.9	370.6
<i>bta-miR-30d</i>	97,949	12,156.3	63,107	5,884.1	8,574.8	<i>bta-miR-99a-5p</i>	3,502	434.6	2,995	279.3	345.9
<i>bta-miR-2340</i>	80,052	9,935.1	66,378	6,189.1	7,796.1	<i>bta-miR-331</i>	3,284	407.6	2,952	275.2	332
<i>bta-let-7b</i>	98,033	12,166.7	47,944	4,470.3	7,771.9	<i>bta-miR-2284x</i>	4,192	520.3	1,937	180.6	326.3
<i>bta-miR-30e-5p</i>	45,947	5,702.4	99,728	9,298.6	7,755.9	<i>bta-miR-455-3p</i>	5,401	670.3	417	38.9	309.8
<i>bta-miR-24-3p</i>	55,651	6,906.7	48,356	4,508.7	5,537.4	<i>bta-miR-199a-3p</i>	2,350	291.7	3,078	287	289
<i>bta-let-7g</i>	40,814	5,065.4	37,961	3,539.5	4,194.1	<i>bta-miR-28</i>	2,505	310.9	2,673	249.2	275.7
<i>bta-miR-26a</i>	34,314	4,258.6	38,649	3,603.6	3,884.6	<i>bta-miR-885</i>	3,531	438.2	1,513	141.1	268.5
<i>bta-miR-103</i>	40,051	4,970.7	31,093	2,899.1	3,787.8	<i>bta-miR-423-3p</i>	3,054	379	1,666	155.3	251.3
<i>bta-miR-27b</i>	44,864	5,568	26,279	2,450.2	3,787.7	<i>bta-miR-2285t</i>	1,275	158.2	3,252	303.2	241
<i>bta-miR-127</i>	39,439	4,894.7	23,104	2,154.2	3,329.8	<i>bta-miR-342</i>	2,666	330.9	1,764	164.5	235.9
<i>bta-miR-423-5p</i>	46,473	5,767.7	14,950	1,393.9	3,270.2	<i>bta-miR-2285i</i>	1,923	238.7	2,471	230.4	233.9
<i>bta-miR-30c</i>	31,144	3,865.2	30,057	2,802.5	3,258.4	<i>bta-miR-199a-5p</i>	2,595	322.1	1,623	151.3	224.6
<i>bta-miR-92a</i>	35,708	4,431.7	22,295	2,078.8	3,088.1	<i>bta-miR-10a</i>	2,006	249	1,869	174.3	206.3
<i>bta-miR-152</i>	24,291	3,014.7	30,852	2,876.6	2,935.9	<i>bta-miR-185</i>	1,727	214.3	2,007	187.1	198.8
<i>bta-miR-145</i>	30,560	3,792.7	22,396	2,088.2	2,819.4	<i>bta-miR-486</i>	2,406	298.6	1,280	119.3	196.2
<i>bta-miR-191</i>	30,755	3,816.9	20,972	1,955.4	2,754	<i>bta-miR-411a</i>	1,470	182.4	2,154	200.8	192.9
<i>bta-let-7c</i>	29,533	3,665.3	15,760	1,469.5	2,411.4	<i>bta-miR-139</i>	2,244	278.5	1,376	128.3	192.7
<i>bta-miR-194</i>	24,257	3,010.5	17,788	1,658.5	2,238.5	<i>bta-miR-505</i>	2,090	259.4	1,415	131.9	186.6
<i>bta-miR-23b-3p</i>	20,639	2,561.5	19,048	1,776	2,113	<i>bta-miR-17-5p</i>	1,630	202.3	1,864	173.8	186
<i>bta-miR-186</i>	20,776	2,578.5	17,689	1,649.3	2,047.9	<i>bta-miR-106b</i>	1,125	139.6	2,036	189.8	168.3
<i>bta-miR-151-3p</i>	20,077	2,491.7	16,933	1,578.8	1,970.4	<i>bta-miR-15a</i>	1,358	168.5	1,800	167.8	168.1
<i>bta-miR-451</i>	8,510	1,056.2	22,749	2,121.1	1,664.3	<i>bta-miR-32</i>	431	53.5	2,664	248.4	164.8
<i>bta-miR-193b</i>	17,868	2,217.6	11,363	1,059.5	1,556.3	<i>bta-miR-146b</i>	1,628	202	1,356	126.4	158.9
<i>bta-miR-29a</i>	14,498	1,799.3	14,405	1,343.1	1,538.8	<i>bta-miR-206</i>	974	120.9	1,926	179.6	154.4
<i>bta-miR-30f</i>	16,308	2,024	12,138	1,131.7	1,514.5	<i>bta-miR-200c</i>	1,623	201.4	1,168	108.9	148.6
<i>bta-miR-125b</i>	13,918	1,727.3	13,383	1,247.8	1,453.5	<i>bta-miR-708</i>	472	58.6	2,109	196.6	137.4
<i>bta-miR-26b</i>	11,314	1,404.2	9,777	911.6	1,122.9	<i>bta-miR-98</i>	960	119.1	1,533	142.9	132.7
<i>bta-miR-100</i>	10,433	1,294.8	9,971	929.7	1,086.3	<i>bta-miR-215</i>	426	52.9	2,015	187.9	130
<i>bta-miR-30b-5p</i>	6,914	858.1	13,348	1,244.6	1,078.8	<i>bta-miR-3432</i>	1,591	197.5	821	76.5	128.4
<i>bta-miR-365-3p</i>	9,859	1,223.6	10,292	959.6	1,072.9	<i>bta-miR-361</i>	1,539	191	854	79.6	127.4
<i>bta-miR-148b</i>	14,899	1,849.1	5,231	487.7	1,071.7	<i>bta-miR-1940</i>	943	117	1,394	130	124.4
<i>bta-miR-93</i>	11,603	1,440	7,282	679	1,005.5	<i>bta-miR-150</i>	1,435	178.1	830	77.4	120.6
<i>bta-let-7d</i>	9,114	1,131.1	8,230	767.4	923.4	<i>bta-miR-2484</i>	1,580	196.1	620	57.8	117.1
<i>bta-miR-23a</i>	7,554	937.5	8,017	747.5	829	<i>bta-miR-200b</i>	1,438	178.5	740	69	116
<i>bta-let-7i</i>	8,951	1,110.9	6,467	603	820.9	<i>bta-miR-195</i>	839	104.1	1,306	121.8	114.2
<i>bta-miR-126-5p</i>	3,028	375.8	10,647	992.7	728.1	<i>bta-miR-193a-3p</i>	405	50.3	1,725	160.8	113.4
<i>bta-miR-25</i>	6,956	863.3	6,179	576.1	699.3	<i>bta-miR-29b</i>	418	51.9	1,588	148.1	106.8
<i>bta-miR-126-3p</i>	7,816	970	5,250	489.5	695.6	<i>bta-miR-374b</i>	1,085	134.7	913	85.1	106.4
<i>bta-miR-339a</i>	6,292	780.9	5,246	489.1	614.3	<i>bta-miR-22-5p</i>	609	75.6	1314	122.5	102.4
<i>bta-miR-320a</i>	7,320	908.5	4,090	381.3	607.5						

RFI, residual feed intake; RC, read counts; RCPM, read count per million mapped reads.

illustrated in Table 4. For instance, *bta-mir-143-3p* was expressed by approximately 605 RCPM, while mature *bta-miR-143-5p* was expressed by only about 19.5 RCPM in the average of the two libraries. Therefore, more studies still



**Figure 2.** Precursor miRNA structures, the 5'p (mature sequence) is highlighted in red and 3'p (miRNA\* sequence) is highlighted in black bold fonts.

needed to evaluate which isomiRs is dominant and

functional to improve the bovine miRNA annotation.

**Putative homologous bovine miRNAs**

Bovine putative homologous miRNAs are those miRNAs that have been reported in other species but not in bovine miRNAs database yet. After removing all reads matched to known bovine miRNAs, the remaining reads were then aligned to non-bovine sets of known mature miRNAs from other species (miRBase v. 21). There were

**Table 4.** Novel isomiRs detected in low and high RFI libraries and their related expression (read counts)

miRNA	Read sequence	Low RFI library		High RFI library		Average of two libraries
		RC	RCPM	RC	RCPM	RCPM
bta-mir-30a-3p	CTTTCAGTCGGATGTTTGCAGC	22,799	2,829.54	23,959	2,233.93	2,489.44
bta-mir-30e-3p	CTTTCAGTCGGATGTTTACAGC	11,837	1,469.07	17,288	1,611.93	1,550.64
bta-mir-122-3p	AACGCCATTATCACACTAAATA	2,707	335.96	10,549	983.58	705.76
bta-mir-142-3p	AGTGTTCCTACTTTATGGA	2,743	340.43	8,622	803.91	605.08
bta-mir-30f-3p	CTGGGAGAAGGCTGTTACTCT	1,067	132.42	639	59.58	90.83
bta-let-7d-3p	TATACGACCTGCTGCCTTCT	1,096	136.02	527	49.14	86.41
bta-mir-148a-5p	AAAGTTCTGAGACACTCCGACT	703	87.25	796	74.22	79.81
bta-mir-125b-2-3p	ACAAGTCAGGCTCTTGGGACCT	823	102.14	661	61.63	79.01
bta-mir-140-5p	CAGTGGTTTTACCCTATGGTAGG	1,373	170.40	0	0.00	73.10
bta-mir-885-3p	AGGCAGCGGGGTGTAGTGGATA	851	105.62	312	29.09	61.92
bta-mir-194-2-3p	CAGTGGGGCTGCTGTTATCTG	476	59.08	210	19.58	36.52
bta-mir-374a-3p	CTTATCAGTTGTATTGTAATT	93	11.54	582	54.27	35.94
bta-mir-106b-3p	CCGCACTGTGGGTACTTGCTG	351	43.56	255	23.78	32.26
bta-mir-361-3p	CCCCAGGTGTGATTCTGATTTGC	319	39.59	182	16.97	26.67
bta-mir-148b-5p	GAAGTTCTGTTATACACTCAGGCT	0	0.00	369	34.41	19.65
bta-mir-139-3p	TGGAGACGCGGCCCTGTTGGAGT	346	42.94	0	0.00	18.42
bta-mir-335-3p	GTTTTTCATTATTGCTCCTGACC	126	15.64	116	10.82	12.88
bta-mir-532-3p	CCTCCCACACCCAAGGCTTGCA	172	21.35	62	5.78	12.46
bta-mir-379-3p	TATGTAACATGGTCCACTAAC	157	19.48	62	5.78	11.66
bta-mir-145-3p	GGATTCTGGAATACTGTCTTT	191	23.70	0	0.00	10.17
bta-mir-411a-3p	TATGTAACACGGTCCACTAACC	174	21.59	0	0.00	9.26
bta-mir-26b-3p	CCTGTTCTCCATTACTTGGCT	97	12.04	58	5.41	8.25
bta-mir-129-2-5p	CTTTTTGCGGTCTGGGCTTGC	61	7.57	46	4.29	5.70
bta-mir-15b-3p	CGAATCATTATTGCTGCTCTA	55	6.83	47	4.38	5.43
bta-mir-3432-2-3p	CAGCAACTAAAGATCCCTCAGG	71	8.81	27	2.52	5.22
bta-mir-7-2-3p	CAACAAATCACAGTCTGCCATA	28	3.48	63	5.87	4.84
bta-mir-185-3p	AGGGGCTGGCTTTCCTCCGGC	58	7.20	23	2.14	4.31
bta-mir-27b-5p	AGAGCTTAGCTGATTGGTGAACA	39	4.84	38	3.54	4.10
bta-mir-136-3p	CATCATCGTCTCAAATGAGTCT	0	0.00	74	6.90	3.94
bta-mir-493-5p	TTGTACATGGTAGGCTTTCATT	57	7.07	16	1.49	3.89
bta-mir-3613-3p	ACAAAAAAGGCAACCCCT	30	3.72	24	2.24	2.88
bta-let-7e-3p	TATACGGCCTCCTAGCTTCC	50	6.21	0	0.00	2.66
bta-mir-195-3p	CCAATATTGGCTGTGCTGCTCCA	30	3.72	18	1.68	2.56
bta-mir-338-5p	AACAATATCCTGGTGCTGAGT	34	4.22	11	1.03	2.40
bta-mir-33a-3p	CAATGTTCCACAGTGCATCA	0	0.00	40	3.73	2.13
bta-mir-2284w-3p	AAAACCTCAATGAACCTTTTGG	0	0.00	39	3.64	2.08
bta-mir-125b-1-3p	ACGGGTTAGGCTCTTGGGAGC	24	2.98	13	1.21	1.97
bta-mir-26a-2-3p	CCTATTCTTGATTACTTGTTTC	0	0.00	36	3.36	1.92
bta-mir-30d-3p	CTTTCAGTCAGATGTTTGTCTGC	6	0.74	27	2.52	1.76
bta-mir-21-3p	CAACAGCAGTCGATGGGCTGTC	0	0.00	29	2.70	1.54
bta-mir-16a-3p	CCAGTATTAACCTGTGCTGCTGAA	0	0.00	25	2.33	1.33
bta-mir-374b-3p	CTTATCAGGTTGATTATCATT	7	0.87	18	1.68	1.33
bta-mir-19b-5p	AGTTTTGCAGGTTTGCATCCAGC	10	1.24	14	1.31	1.28
bta-mir-210-5p	AGCCACTGCCACCGCACACTGC	15	1.86	8	0.75	1.22
bta-mir-365-2-5p	GAGGGACTTTCAGGGGCAGCTGT	19	2.36	0	0.00	1.01
bta-mir-130b-5p	ACTCTTCCCTGTGCACTACT	12	1.49	0	0.00	0.64
bta-mir-25-5p	AGGCGGAGACTTGGGCAATTGCT	11	1.37	0	0.00	0.59
bta-mir-204-3p	GCTGGGAAGGCAAAGGGACGT	8	0.99	0	0.00	0.43
bta-mir-380-5p	ATGGTTGACCATAGAACATGCG	8	0.99	0	0.00	0.43
bta-mir-369-5p	AGATCCACCGTGTATATTCG	6	0.74	0	0.00	0.32

RFI, residual feed intake; RC, read counts; RCPM, read count per million mapped read.

61 detected putative homologous miRNAs which have distinct hairpin loci in the bovine genome (Table 5). These putative novel homologous miRNAs sequences and related genomic locations are presented in Additional files, Supplementary Table S2.

### Detection of putative novel bovine microRNAs

More than quarter of the reads (25% from high and 31%

**Table 5.** Unique putative homolog miRNAs identified in bovine liver tissue

miRNA	Homolog mature miRNAs name	Sequence	Align length	Low RFI library		High RFI library		Average of two libraries CPM
				RC	CPM	RC	CPM	
miR-4448	<i>hsa-miR-4448</i>	GGCTCGTTGGTCTAGGGGTATGATTC	20	2,405	298.48	1,464	136.50	217.49
miR-203-3p	<i>mmu-miR-203-3p; hsa-miR-203a; rno-miR-203a-3p; mml-miR-203; ptr-miR-203; cfa-miR-203; ppy-miR-203</i>	TGAAATGTTTAGGACCAGTATCT	21	512	63.54	3,342	311.61	187.57
miR-6243	<i>mmu-miR-6243</i>	ACCATCTGTGGGATTATGACTGAACG	26	1,612	200.06	1,883	175.57	187.82
miR-574-3p	<i>hsa-miR-574-3p; mmu-miR-574-3p; cfa-miR-574; ssc-miR-574; ggo-miR-574</i>	CACGCTCATGCACACCCACATCTC	22	1,875	232.70	1,218	113.57	173.13
miR-802-5p	<i>hsa-miR-802-5p; mmu-miR-802-5p</i>	TCAGTAACAAGATTATCCTTGT	21	624	77.44	1,782	166.15	121.80
miR-1285	<i>cgr-miR-1285</i>	CTCCAGCCTGGGCAACATAGCGAGAC	20	1,485	184.30	603	56.22	120.26
miR-4532	<i>hsa-miR-4532</i>	CCCCGGGGAGCCCGCGGGCATCTCG	17	1,034	128.33	574	53.52	90.92
miR-3535	<i>gga-miR-3535</i>	GGATATGATGACTGATTATCTGAGAA	23	1,023	126.96	268	24.99	75.98
miR-664-3p	<i>ssc-miR-664-3p</i>	TATTCATTTATCTCCAGCCTACAAA	20	1,285	159.48	1	0.09	79.79
miR-5100	<i>mmu-miR-5100</i>	TCGAATCCCAGCGGTGCCTCCAATCT	20	737	91.47	375	34.96	63.22
miR-716b	<i>sha-miR-716b</i>	TCTTGGTGGTAGCAAAATATCAA	22	367	45.55	124	11.56	28.55
miR-5115	<i>mmu-miR-5115</i>	CTGGACCGGAGCCGGCCCTCCCGT	19	191	23.70	126	11.75	17.73
miR-6238	<i>mmu-miR-6238</i>	TATTAGTCAGCGGAGGAAAAAGAACT	19	170	21.10	86	8.02	14.56
miR6173	<i>hbr-miR6173</i>	CGTAAACGATGAATACTAGGTGTCGG	17	247	30.65	0	0.00	15.33
miR-6239	<i>mmu-miR-6239</i>	AGCGGTGGATCACTCGGCTCGTGCGT	17	114	14.15	44	4.10	9.13
miR-6240	<i>mmu-miR-6240</i>	CAAAGCATCGGAAGGCCCGCATCT	19	120	14.89	23	2.14	8.52
miR-6412	<i>mmu-miR-6412</i>	TCGAAACCATCTCTGCTACCAATCT	20	90	11.17	21	1.96	6.56
miR-1895	<i>mmu-miR-1895</i>	AGAGGAGGACGAGGAGGAAGAGGAGG	18	60	7.45	33	3.08	5.26
miR-320d	<i>hsa-miR-320d; ppy-miR-320d</i>	AAAAGCTGGTTGAGAGGATCTCGTA	19	56	6.95	30	2.80	4.87
miR-6129	<i>ptr-miR-6129; hsa-miR-6129</i>	TGAGGGAGTAGGGTGTATAGTATCTC	19	48	5.96	31	2.89	4.42
miR-2779	<i>bmo-miR-2779</i>	TTCCGGCTCGAAGGACCAATCTCGT	19	49	6.08	29	2.70	4.39
miR-124c-3p	<i>gga-miR-124c-3p</i>	TCAAGGTCCGCTGTGAACACGGATCT	0	25	3.10	50	4.66	3.88
miR-5109	<i>mmu-miR-5109</i>	TGGTGGCGACCAGGGGAATCCGACAT	23	41	5.09	30	2.80	3.94
miR-1230	<i>mml-miR-1230</i>	TGGTTCGGGGCATCTCGTATGCGTC	17	1	0.12	67	6.25	3.19
miR-6236	<i>mmu-miR-6236</i>	GCCGTGCCCGCAGTCGGAGAGATCT	18	45	5.58	8	0.75	3.17
miR-5097	<i>mmu-miR-5097</i>	TCATGTCCCTGTTCGGGCGCAATCT	22	22	2.73	19	1.77	2.25
miR-300-3p	<i>mmu-miR-300-3p; rno-miR-300-3p; cgr-miR-300</i>	TATGCAGGGCAAGCTCTCTGATCT	20	18	2.23	21	1.96	2.10
miR-24b	<i>xtr-miR-24b; xla-miR-24b</i>	TGGCTCAGTTCAGCAGGATCTCGT	18	17	2.11	19	1.77	1.94
miR-4485	<i>hsa-miR-4485</i>	AACGGCCGCGGTATCCTGACCGTGCA	17	7	0.87	23	2.14	1.51
miR-4497	<i>hsa-miR-4497</i>	CTCCGGGACGGCTGGGAAGCCGGCA	23	23	2.85	7	0.65	1.75
miR6300	<i>gma-miR6300</i>	GTCGTTGTAGTATAGTGGTAGTATT	18	28	3.48	0	0.00	1.74
miR-5108	<i>mmu-miR-5108</i>	GTAGAGCACTGGATGGATCTCGTATG	18	27	3.35	1	0.09	1.72
miR-5106	<i>mmu-miR-5106</i>	GGGTCTGTAGCTCAGTTGGTTAGAGC	19	25	3.10	1	0.09	1.60
miR-323c	<i>oar-miR-323c</i>	CACAATACACGGTCCGCCTATCTC	21	16	1.99	8	0.75	1.37
miR-5119	<i>mmu-miR-5119</i>	CATCACATCCTGGGGCTGTAGCCGGA	18	18	2.23	0	0.00	1.12
miR-3168	<i>hsa-miR-3168</i>	GAGTCTACAGTCCGACGATCGTATG	18	0	0.00	18	1.68	0.84
miR-4492	<i>hsa-miR-4492</i>	GGGGCTGGGCGCGCCGCGCATCT	17	12	1.49	6	0.56	1.02
miR-27e	<i>dre-miR-27e; fru-miR-27e; tni-miR-27e</i>	TTACAGTGGCTAAGTAGAATCTCGT	20	8	0.99	10	0.93	0.96
miR-6089	<i>hsa-miR-6089</i>	CGGGGTGGTTCGGGCGGGGCGGACT	18	16	1.99	0	0.00	0.99
miR-6327	<i>rno-miR-6327</i>	AGGACTGTAGATCCATCTCGTATGCC	18	0	0.00	12	1.12	0.56
miR-1957a	<i>mmu-miR-1957a</i>	AGTGGTAGAGCATTTGACTGATCTCG	18	1	0.12	11	1.03	0.57
miR159c-3p	<i>ath-miR159c; aly-miR159c-3p</i>	TTTGATTGAAGGGAGCATCTCGTAT	17	10	1.24	0	0.00	0.62
miR-3607-3p	<i>hsa-miR-3607-3p</i>	ACTGTAAACGCTTCTGATGATCTCG	20	10	1.24	0	0.00	0.62
miR-5124a	<i>mmu-miR-5124a</i>	GTCAAGTGACTAAGAGCATATGGTGG	19	7	0.87	1	0.09	0.48
miR-3591-5p	<i>hsa-miR-3591-5p</i>	TTTAGTGTGATAATGGCTTTATCTC	21	1	0.12	7	0.65	0.39
miR-378g	<i>hsa-miR-378g</i>	ACTGGGCTTGGAGTCGGAAGGCATCT	20	1	0.12	7	0.65	0.39
miR-4792	<i>hsa-miR-4792</i>	CGGTGAGCTCTCGTGGCATCTCGGA	18	1	0.12	7	0.65	0.39
miR-1261	<i>hsa-miR-1261</i>	ATGGATAAGGCATTGGCTTCTTAAGC	19	6	0.74	0	0.00	0.37
miR-535d	<i>mdm-miR535d</i>	TGACGACGAGAGAGACGACATCTC	21	6	0.74	0	0.00	0.37
miR-1949	<i>mmu-miR-1949</i>	CTATACCAGGATGCCAGCATAGTTAT	24	6	0.74	0	0.00	0.37
miR-378a	<i>ppy-miR-378a</i>	ACTGGACTTGGGTGAGGAGGAGCATCTC	21	6	0.74	0	0.00	0.37
miR-6516-3p	<i>gga-miR-6516-3p</i>	CATGTATGATACTGCAACAGAAATC	20	0	0.00	6	0.56	0.28

RFI, residual feed intake; RC, read counts; RCPM, read count per million mapped reads.



from low RFI libraries) that did not match to known bovine miRNAs matched to the bovine genome. To identify novel miRNAs, we extracted the candidate pre-miRNA structure, based on the location of clusters of mature miRNA on the genome, to select the energetically best candidate if they were having: i) at least 19 base pairings in the secondary structure, and ii) at least 11 base pairings located in the read cluster region (the number of pairings between 5'p and 3'p). Then five different models (Hackenberg et al., 2011) were used to predict whether a given candidate was likely to be a miRNA. We considered a candidate as a novel miRNA only if at least three out of five models were positive. Twelve putative new miRNAs were discovered in the present study. After realigning these putative novel consensus sequences to known mature miRNAs in other species, two candidates (*miR-664-3p* and *miR-802-5p*) were found to be homologous to other species. Ten novel bovine-specific candidate miRNAs with their sequence, genomic location, and the number of the models that predicted each novel miRNA candidate are presented in Table 6. These ten novel bovine miRNAs represented by 5,437 and 1,097 reads in the low RFI and high RFI libraries, respectively. The sequences and secondary structure of those 10 novel bovine-specific miRNAs candidates and related pre-miRNAs with the MFE, which have been predicted by RNAfold program, are illustrated in Additional files, Supplementary Figure S1. Novel candidate 1 and 2 have identical mature sequences but resulted from distinct precursor sequences located in different genomic region.

#### Differentially expressed miRNA between high and low RFI libraries and their target genes

To understand the role of miRNA in regulation of residual intake, we first identified differentially expressed miRNAs from high and low RFI selection lines. As miRNAs play important roles in the regulation of almost every biological process in eukaryotes; these differentially

expressed miRNAs could play important roles in regulation of the physiological processes and pathways involved in variation in RFI in beef cattle. To define the differentially expressed miRNAs, we consider only those miRNAs with high to modest expression with at least 100 RCPM and folds changes >2. This approach was adapted in earlier microarray experiment and was less stringent than using  $p < 0.05$  due to lack of replicates. A total of 49 unique miRNAs were identified to be differentially expressed between the high RFI line and low RFI line cattle (Table 7). More than two thirds (33 out of 49) of differentially expressed miRNAs were up-regulated in high RFI animals. Specifically, six miRNA (*miR-203-3p*, *bta-miR-32*, *bta-miR-215*, *bta-miR-708*, and *bta-miR-101*) that reached  $p < 0.05$  were all up-regulated in high RFI animals.

Importantly, many up-regulated miRNAs in high RFI animals played important roles in metabolic homeostasis including glucose and lipid metabolism. These overexpression patterns were similar to the patterns observed in obesity mouse models and human subject. For example *bta-miR-143*, the most expressed miRNA in bovine liver, was up-regulated in high RFI cattle. It has been reported that hepatic *miR-143* were up-regulated in obesity mouse models whether they are genetic or dietary induced. The overexpression of hepatic *miR-143* impaired insulin-stimulated AKT activation and glucose homeostasis by targeting insulin signalling and its regulation (Jordan et al., 2011). *Bta-mir-122-3p* was highly expressed in bovine liver and is up-regulated in high RFI cattle. *MiR-122* was the first miRNA to be linked to metabolic control and affect hepatic cholesterol, lipid metabolism and the maintenance of liver cell differentiation (Lewis and Jopling, 2010).

A bovine homologous of *miR-802-5p* was identified in our study and was up-regulated in high RFI cattle. *MiR-802* has been reported up-regulated in the liver of two obese mouse models and obese human subjects (Kornfeld et al., 2013). Overexpression of *miR-802* in mice causes impaired

**Table 6.** Novel bovine miRNAs detected in the bovine liver tissue, with sequence, genomic location and their related expression (read counts)

Predicted miRNA ID	Chr.	Position		Strand	No.mod. Pred.	Read cluster sequence (5' - 3')	RC in two pools <sup>1</sup>
		Start	End				
Candidate_1	chr11	76,942,835	76,942,919	+	3	GAGAGAACGTAATCTGAGTGGTTTC	2,102
Candidate_2	chr19	8,794,560	8,794,644	-	4	GAGAGAACGTAATCTGAGTGGTTTC	2,102
Candidate_3	chr21	65,541,217	65,541,301	-	4	TTCCTGGGCATCCTCTGCTTTAT	1,022
Candidate_4	chr27	7,419,033	7,419,126	+	5	GTTCCGGGGGGAGTATGGTTGCAAAG	416
Candidate_5	chr28	32,693,899	32,693,979	-	3	GTCTGGTGGGAAGGAAGGGACACATC	217
Candidate_6	chr29	49,019,853	49,019,926	-	3	GGAATACCGGGTGCTGCAGGCTTT	63
Candidate_7	chr21	173,761	173,839	+	4	TTGTCCTACTTCTCAGCTGTCTT	53
Candidate_8	chr27	7,419,033	7,419,119	+	4	TTATTCCCATGACCCGCTGGCAGC	22
Candidate_9	chr3	67,781,210	67,781,312	-	3	CTGCGGGATGAACCGAACGCCGGGTTAAG	339
Candidate_10	chr12	36,358,265	36,358,343	-	5	TCCACATCCCTCACAGTTTGGTG	198

Chr., chromosome number; No.mod. Pred., the number of models that predict the specific novel microRNA; RFI, residual feed intake.

<sup>1</sup> RC in two pools, total reads count in low and high RFI libraries.



**Table 7.** miRNAs differentially expressed between the high and low RFI libraries

miRNA	Base-mean		Fold-change	p-value
	Low RFI	High RFI		
bta-miR-32	392.5	2,925.6	7.45	0.02
bta-miR-215	387.9	2,212.8	5.7	0.04
bta-miR-708	429.8	2,316.1	5.39	0.05
bta-miR-101	50,519	268,425.1	5.31	0.03
bta-miR-193a-3p	368.8	1,894.4	5.14	0.06
bta-miR-29b	380.6	1,743.9	4.58	0.08
bta-miR-1	2,820.1	12,150.3	4.31	0.06
bta-miR-21-5p	35,761.9	152,774.6	4.27	0.06
bta-miR-126-5p	2,757.3	11,692.3	4.24	0.06
bta-miR-424-5p	220.4	932.4	4.23	0.12
bta-miR-192	96,668.9	339,724.6	3.51	0.1
bta-miR-6119-5p	176.7	600.7	3.4	0.21
bta-miR-143	630,803	2,068,377.1	3.28	0.12
bta-miR-451	7,749.2	24,982.5	3.22	0.13
bta-miR-19b	364.2	1,166.3	3.2	0.19
bta-miR-7	123.8	394.2	3.18	0.29
bta-miR-2285t	1,161	3,571.3	3.08	0.16
bta-miR-29c	117.5	355.8	3.03	0.32
bta-miR-374a	359.7	1,089.4	3.03	0.21
bta-miR-30e-5p	41,839.2	109,519.4	2.62	0.2
bta-miR-148a	145,596.2	379,981.7	2.61	0.2
bta-miR-22-5p	554.6	1,443	2.6	0.26
bta-miR-146a	2,093.5	5,218.6	2.49	0.24
bta-miR-6120-3p	269.5	649	2.41	0.35
bta-miR-206	886.9	2,115.1	2.38	0.28
bta-miR-30b-5p	6,295.9	1,4658.5	2.33	0.27
bta-miR-106b	1,024.4	2,235.9	2.18	0.33
bta-miR-2285f	592.8	1,278.3	2.16	0.36
bta-miR-130a	621	1,247.5	2.01	0.41
bta-miR-31	584.6	289.9	0.5	0.46
bta-miR-874	642.9	318.5	0.5	0.45
bta-miR-1307	1,068.1	519.4	0.49	0.4
bta-miR-15b	1,160.1	551.3	0.48	0.38
bta-miR-2484	1,438.7	680.9	0.47	0.37
bta-miR-148b	13,567	5,744.6	0.42	0.26
miR-716b	334.2	136.2	0.41	0.42
bta-miR-380-3p	284.1	113.1	0.4	0.43
bta-miR-197	1,249.3	488.7	0.39	0.27
bta-miR-423-5p	42,318.2	16,417.8	0.39	0.21
bta-miR-375	312.3	114.2	0.37	0.38
bta-miR-6529	356	124.1	0.35	0.34
bta-miR-455-3p	4,918.1	457.9	0.09	0.01
isomiRs differentially expressed between high and low RFI				
bta-mir-122-3p	2,465	11,584.7	4.7	0.05
bta-mir-142-3p	2,497.8	9,468.5	3.79	0.09
bta-mir-885-3p	774.9	342.6	0.44	0.37
Homologous differentially expressed between high and low RFI				
miR-203-3p	466.2	3,670.1	7.87	0.01
miR-802-5p	568.2	1,957	3.44	0.14
miR-1285	1,352.2	662.2	0.49	0.39
miR-3535	931.5	294.3	0.32	0.2

RFI, residual feed intake.

glucose tolerance and attenuates insulin sensitivity by suppressing its target gene HNF1 Homeobox B (Hnf1b).

*Bta-miR-29b* was up-regulated in high RFI. The function of *miR-29* has been reported to regulate glucose transport in liver, muscle and adipose (Pandey et al., 2011). *Bta-miR-19b*, *bta-miR-101*, *bta-miR-106b*, and *bta-miR-142-3p* were up-regulated in high RFI cattle. Recently studies showed their expression in steers liver were highly influenced by energy density of the diet (mainly lipid levels in the diet) (Romao et al., 2012).

To understand miRNAs expression and their target genes, we searched all the potential miRNAs target sites in the 3'UTR region of 161 differentially expressed genes identified by Chen et al. (2011) by using miRmap (Vejnar and Zdobnov, 2012). Then we examined if these potential miRNAs were expressed differentially in liver between high and low RFI cattle.

We found 36 differentially-expressed genes in liver (Chen et al., 2011) containing putative miRNAs target sites (Table 8). Many genes have multiple miRNAs target sites such as helicase with zinc finger (*HELZ*), espin (*ESPN*), cytochrome P450 family 2 subfamily C member 18 (*CYP2C18*), snail family transcriptional repressor 2 (*SNAIL2*), and superoxide dismutase 3 (*SOD3*). Five down-regulated miRNAs (*bta-miR-424-5P*, *bta-miR-19b*, *bta-miR-29b*, *bta-miR-30b*, and *bta-miR-2285*) having sites binding to the mRNA of 12 distinct genes calponin 1 (*CNN1*), atypical chemokine receptor 3 (*CXCR7*), endothelin receptor type (*BEDNRB*), fibrinogen alpha chain (*FGA*), insulin like growth factor binding protein 3 (*IGFBP3*), regulator of G-protein signaling 2 (*RGS2*), periostin, osteoblast specific factor (*POSTN*), monoamine oxidase (*AMAOA*), collagen type IV alpha 6 (*COL4A6*), dehydrogenase/reductase (*SDR family member 3*) (*DHRS3*), solute carrier family 22 (organic anion transporter), member 7 (*SLC22A7*). These genes were up-regulated in low RFI animals (Chen et al., 2011). Four genes, *BEDNRB*, *IGFBP3*, *POSTN*, and *DHRS3* were up-regulated in low-RFI cattle (Chen et al., 2011) and have *bta-miR-19b* putative target sites in their 3'UTR region. Take together we believe these differentially expressed genes play important roles regulating RFI in beef cattle.

However, we acknowledge the differentially expressed genes detected in this study are based on pools with limited statistic power. Follow up studies are need for validating the differentially expressed miRNAs and their function.

In conclusion, RFI as a measure of feed efficiency is influenced by several physiological systems including basal metabolic rate, energy balance, the regulation of growth and development, regulation of feed intake and homeostatic control of body mass. Our study revealed a comprehensive miRNA population in bovine liver. We identified 305 known bovine miRNAs, 50 novel isomiRs, 52 homologous miRNAs, and 10 novel miRNAs. We further revealed that

**Table 8.** Differentially expressed genes and their potential miRNA expression in liver; up and down- regulations were based on contrast of low RFI<sup>1</sup>

Target gene	miRNA	Seed match	Expression of miRNA	Expression of mRNA	Network ID	Top functions
<i>AVPRIA</i>	<i>bta-miR-885</i>	8mer	UP	Down	1	Cellular Growth and Proliferation, Cancer, Cardiovascular System Development and Function
<i>CNN1</i>	<i>bta-miR-424-5P</i>	7mer-m8	Down	Up	1	
<i>CXCR7</i>	<i>bta-miR-29b</i>	7mer-m8	Down	Up	1	
<i>EDNRB</i>	<i>bta-miR-19b</i>	8mer	Down	Up	1	
<i>FGA</i>	<i>bta-miR-29b</i>	7mer	Down	Up	1	
<i>GHR</i>	<i>bta-miR-101</i>	8mer	Down	Down	1	
<i>IGFBP3</i>	<i>bta-miR-19b</i>	8mer	Down	Up	1	
<i>NKIRAS1</i>	<i>bta-miR-19b</i>	7mer-m8	Down	Down	1	
<i>RGS2</i>	<i>bta-miR-30b</i>	7mer-m9	Down	Up	1	
<i>AHR</i>	<i>bta-miR-29b</i>	7mer-m8	Down	Down	2	Hepatic System Disease, Dermatological Disease and Conditions, Cellular Growth and Proliferation
<i>CD4</i>	<i>bta-miR-143</i>	7mer	Down	Up	2	
<i>GSTM1</i>	<i>bta-miR-30b-3p</i>	7mer-m8	Down	Down	2	
<i>S100A10</i>	<i>bta-miR-21</i>	7mer-m8	Down	Down	2	
<i>HELZ</i>	<i>bta-miR-15</i>	8mer	Up	Down	2	
	<i>bta-miR-424</i>	8mer	Down	Down	2	
<i>HLA-DRB1</i>	<i>bta-miR-197</i>	7mer-m8	Up	Up	2	
<i>POSTN</i>	<i>bta-miR-19b</i>	8mer	Down	Up	2	
<i>AP3B2</i>	<i>bta-miR-30b</i>	7mer-m8	Up	Down	3	Cellular Assembly and Organization, Cancer, Cellular Movement
<i>ESPN</i>	<i>bta-miR-424</i>	7mer-m8	Down	Down	3	
	<i>bta-miR-15b</i>	7mer-m8	Up	Down	3	
<i>MAOA</i>	<i>bta-miR-2285</i>	8mer	Down	Up	3	
<i>CPEB1</i>	<i>bta-miR-19b</i>	7mer-m8	Down	Down	4	Protein Synthesis, Development Disorder, Neurological disease
<i>AHSG</i>	<i>bta-miR-31</i>	8mer	Up	Up	5	
<i>COL3A1</i>	<i>bta-miR-122</i>	8mer	Down	Up	5	Drug Metabolism, Endocrine System Development, Lipid Metabolism
<i>CYP2C18</i>	<i>bta-miR-424</i>	7mer-m8	Down	Down	5	
	<i>bta-miR-143</i>	7mer	Down	Down	5	
<i>MEP1B</i>	<i>bta-miR-32</i>	7mer	Down	Down	5	
<i>SLC27A6</i>	<i>bta-miR-424</i>	7mer-m8	Down	Down	5	
<i>ABCC4</i>	<i>bta-miR-19b</i>	7mer-m8	Down	Down	6	Carbohydrate Metabolism, Drug Metabolism, Small Molecular Biochemistry
<i>ABHD5</i>	<i>bta-miR-19b</i>	7mer-m8	Down	Down	6	
<i>COL4A6</i>	<i>bta-miR-29b</i>	8mer	Down	Up	6	
<i>MAP2K6</i>	<i>bta-miR-29b</i>	7mer	Down	Down	6	
<i>SNAI2</i>	<i>bta-miR-30b</i>	7mer-m8	Down	Down	6	
	<i>bta-miR-30d</i>	7mer-m8	Down	Down	6	
<i>AVPRIA</i>	<i>bta-miR-885</i>	8mer	Up	Down	7	Cell Death, Cell Signaling, Molecular Transport
<i>DDC</i>	<i>bta-miR-708</i>	7mer-m8	Down	Down	7	
<i>DHRS3</i>	<i>bta-miR-19b</i>	8mer	Down	Up	7	
<i>SLC22A7</i>	<i>bta-miR-29b</i>	8mer	Down	Up	7	
<i>SOD3</i>	<i>bta-miR-1</i>	7mer-m8	Down	Down	7	
	<i>bta-miR-423-5p</i>	7mer	Up	Down	7	
	<i>bta-miR-708</i>	7mer-m8	Down	Down	7	
	<i>bta-miR-19a</i>	7mer	Down	Down	7	
	<i>bta-miR-19b</i>	7mer	Down	Down	7	

RFI, residual feed intake.

<sup>1</sup> The differentially expressed genes and gene networks were from previous study (Chen et al., 2011).

many up-regulated miRNAs in high RFI cattle showed a similar expression pattern as found in a mouse obesity model and have functions related to glucose and lipid metabolism. We demonstrated the expression of miRNAs have effects on their target genes expression. Combining the patterns of miRNA and mRNA expression will provide

further power to understanding the molecular mechanisms that regulate feed efficiency in beef cattle.

#### CONFLICT OF INTEREST

We certify that there is no conflict of interest with any

financial organization regarding the material discussed in the manuscript.

## ACKNOWLEDGMENTS

This work was funded by the former Cooperative Research Centre for Beef Genetic Technologies and the University of New England. Wijdan Al-Husseini was supported by the University of Babylon/ Ministry of Higher Education and Scientific Research (MOHESR)/Iraq.

## REFERENCES

- Anders, S. and W. Huber. 2010. Differential expression analysis for sequence count data. *Genome Biol.* 11:R106.
- Arthur, P. F., J. A. Archer, D. J. Johnston, R. M. Herd, E. C. Richardson, and P. F. Parnell. 2001. Genetic and phenotypic variance and covariance components for feed intake, feed efficiency, and other postweaning traits in Angus cattle. *J. Anim. Sci.* 79:2805-2811.
- Chen, Y., C. Gondro, K. Quinn, R. M. Herd, P. F. Parnell, and B. Vanselow. 2011. Global gene expression profiling reveals genes expressed differentially in cattle with high and low residual feed intake. *Anim. Genet.* 42:475-490.
- Donoghue, K. A., P. F. Arthur, J. F. Wilkins, and R. M. Herd. 2011. Onset of puberty and early-life reproduction in Angus females divergently selected for post-weaning residual feed intake. *Anim. Prod. Sci.* 51:183-190.
- Fatima, A., D. J. Lynn, P. O'Boyle, C. Seoighe, and D. Morris. 2014. The miRNAome of the postpartum dairy cow liver in negative energy balance. *BMC Genomics* 15:279.
- Gardner, P. P., J. Daub, J. G. Tate, E. P. Nawrocki, D. L. Kolbe, S. Lindgreen, A. C. Wilkinson, R. D. Finn, S. Griffiths-Jones, S. R. Eddy, and A. Bateman. 2009. Rfam: updates to the RNA families database. *Nucl. Acids Res.* 37:D136-D140.
- Kozomara, A. and S. Griffiths-Jones. 2014. miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucl. Acids. Res.* 42:D68-D73.
- Gu, Z., S. Eleswarapu, and H. Jiang. 2007. Identification and characterization of microRNAs from the bovine adipose tissue and mammary gland. *FEBS Lett.* 581:981-988.
- Hackenbarg, M., N. Rodriguez-Ezpeleta, and A. M. Aransay. 2011. miRanalyzer: An update on the detection and analysis of microRNAs in high-throughput sequencing experiments. *Nucl. Acids Res.* 39:W132-W138.
- Hu, J., Y. Xu, J. Hao, S. Wang, C. Li, and S. Meng. 2012. MiR-122 in hepatic function and liver diseases. *Protein Cell.* 3:364-371.
- Jin, W., J. R. Grant, P. Stothard, S. S. Moore, and L. L. Guan. 2009. Characterization of bovine miRNAs by sequencing and bioinformatics analysis. *BMC Mol. Biol.* 10:90.
- Jordan, S. D., M. Kruger, D. M. Willmes, N. Redemann, F. T. Wunderlich, H. S. Bronneke, C. Merkwirth, H. Kashkar, V. M. Olkkonen, T. Bottger, T. Braun, J. Seibler, and J. C. Bruning. 2011. Obesity-induced overexpression of miRNA-143 inhibits insulin-stimulated AKT activation and impairs glucose metabolism. *Nat. Cell Biol.* 13:434-446.
- Koch, R. M., L. A. Swiger, D. Chambers, and K. E. Gregory. 1963. Efficiency of feed use in beef cattle. *J. Anim. Sci.* 22:486-494.
- Kornfeld, J. W., C. Baitzel, A. C. Konner, H. T. Nicholls, M. C. Vogt, K. Herrmanns, L. Scheja, C. Haumaitre, A. M. Wolf, U. Knippschild, J. Seibler, S. Cereghini, J. Heeren, M. Stoffel, and J. C. Brüning. 2013. Obesity-induced overexpression of miR-802 impairs glucose metabolism through silencing of Hnf1b. *Nature* 494:111-115.
- Kozomara, A. and S. Griffiths-Jones. 2011. miRBase: integrating microRNA annotation and deep-sequencing data. *Nucl. Acids Res.* 39:D152-D157.
- Langmead, B., C. Trapnell, M. Pop, and S. L. Salzberg. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.
- Lawless, N., A. B. Foroushani, M. S. McCabe, C. O'Farrelly, and D. J. Lynn. 2013. Next Generation sequencing reveals the expression of a unique miRNA profile in response to a gram-positive bacterial infection. *PLoS One.* 8:e57543.
- Lewis, A. P. and C. L. Jopling. 2010. Regulation and biological function of the liver-specific miR-122. *Biochem. Soc. Trans.* 38:1553-1557.
- Liu, H. C., J. A. Hicks, N. Trakooljul, and S. H. Zhao. 2010. Current knowledge of microRNA characterization in agricultural animals. *Anim. Genet.* 41:225-231.
- Lu, J., G. Getz, E. A. Miska, E. Alvarez-Saavedra, J. Lamb, D. Peck, A. Sweet-Cordero, B. L. Ebert, R. H. Mak, A. A. Ferrando, J. R. Downing, T. Jacks, H. R. Horvitz, and T. R. Golub. 2005. MicroRNA expression profiles classify human cancers. *Nature* 435:834-838.
- Miles J. R., T. G. McDaneld, R. T. Wiedmann, R. A. Cushman, S. E. Echterkamp, J. L. Vallet, and T. P. L. Smith. 2012. MicroRNA expression profile in bovine cumulus-oocyte complexes: Possible role of let-7 and miR-106a in the development of bovine oocytes. *Anim. Reprod. Sci.* 130:16-26.
- Pandey, A. K., G. Verma, S. Vig, S. Srivastava, A. K. Srivastava, and M. Datta. 2011. miR-29a levels are elevated in the db/db mice liver and its overexpression leads to attenuation of insulin action on *PEPCK* gene expression in HepG2 cells. *Mol. Cell. Endocrinol.* 332:125-133.
- Romao, J. M., W. Jin, M. He, T. McAllister, and L. L. Guan. 2012. Altered microRNA expression in bovine subcutaneous and visceral adipose tissues from cattle under different diet. *PLoS One.* 7:e40605.
- Rottiers, V. and A. M. Näär. 2012. MicroRNAs in metabolism and metabolic disorders. *Nat. Rev. Mol. Cell Biol.* 13:239-250.
- Sherman, E. L., J. D. Nkrumah, C. Li, R. Bartusiak, B. Murdoch, and S. S. Moore. 2009. Fine mapping quantitative trait loci for feed intake and feed efficiency in beef cattle. *J. Anim. Sci.* 87:37-45.
- Tripurani, S. K., C. Xiao, M. Salem, and J. Yao. 2010. Cloning and analysis of fetal ovary microRNAs in cattle. *Anim. Reprod. Sci.* 120:16-22.
- Vejnar, C. E. and E. M. Zdobnov. 2012. miRmap: Comprehensive prediction of microRNA target repression strength. *Nucl. Acids Res.* 40:11673-11683.
- Wen, J. and J. R. Friedman. 2012. miR-122 regulates hepatic lipid metabolism and tumor suppression. *J. Clin. Invest.* 122:2773-2776.

- Yang, J. S., M. D. Phillips, D. Betel, P. Mu, A. Ventura, A. C. Siepel, K. C. Chen, and E. C. Lai. 2011. Widespread regulatory activity of vertebrate microRNA\* species. *RNA*. 17:312-326.
- Yu, Z., Z. Jian, S. H. Shen, E. Purisima, and E. Wang. 2007. Global analysis of microRNA target gene expression reveals that miRNA targets are lower expressed in mature mouse and *Drosophila* tissues than in the embryos. *Nucl. Acids Res.* 35:152-164.