

## Autoionization of very-high- $n$ strontium Rydberg atoms

X. Zhang\*, G. Fields\*, F.B. Dunning\*<sup>1</sup>, S. Yoshida†, and J. Burgdörfer†

\* Department of Physics & Astronomy, Rice University, Houston, Texas, USA

† Institute for Theoretical Physics, Vienna University of Technology, Vienna, Austria, EU


**Synopsis** The autoionization of high  $n$ ,  $n \sim 280$ -430, strontium Rydberg states through excitation of the  $5s \ ^2S_{1/2} \rightarrow 5p \ ^2P_{1/2}$  transition in the core ion is investigated. The autoionization rates decrease rapidly as  $L$  is increased paving the way for production of long-lived two-electron-excited planetary atoms.

The ability to create localized wave packets in Rydberg atoms whose behavior mimics that of a classical particle suggests the possible engineering of atoms that contain two excited electrons, each moving along quasi-classical orbits. While such atoms, termed planetary atoms, have been the subject of much theoretical interest,[1] their practical realization is a challenge because interactions between the excited electrons can lead to rapid autoionization. In the present work we explore autoionization using very-high  $n$  strontium Rydberg atoms, and the techniques that can be used to control it.

Strontium atoms in a collimated beam are excited to selected  $n^1P_1$  or  $n^1F_3$  states and autoionization is induced by exciting the  $5s \ ^2S_{1/2} \rightarrow 5p \ ^2P_{1/2}$  transition in the core ion using a 100ns-long pulse of 422 nm laser radiation. (Use of a short pulse minimizes population of the long-lived  $4d \ ^2D_{3/2}$  state which can also induce autoionization) The number of surviving Rydberg atoms is determined by field ionization.

Measurements with both  $nP$  and  $nF$  states in which the frequency of the 422 nm laser is scanned reveal broad autoionization features whose widths decrease with increasing  $n$  and, in the limit of very high  $n$ , approach that of the transition in the  $Sr^+$  core ion.

The effect of increasing the angular momentum,  $L$ , of the excited Rydberg electron is examined either by initially exciting selected extreme Stark states in a dc electric field or by sudden application of a “pump field” to  $n^1P_1$  atoms which creates a Stark wavepacket that undergoes Stark precession resulting in periodic changes in  $|L|$ . [2] Even modest increases in  $L$  are found to result in a dramatic decrease in the autoionization loss. This is illustrated in Fig. 1 which shows the fraction of  $n=320 \ ^1P_1$  atoms that survive autoionization versus the duration of the pump pulse.



**Figure 1.** Rydberg atom survival probability versus pump pulse duration.

Pulse durations of  $\sim 10$ -15 ns are sufficient to populate states with  $|L| \sim 20$ -30.

The present work indicates that by placing one electron in an outer localized very-high- $n$ , high- $\ell$ ,  $\ell \sim n$ , near-circular Bohr-like orbit, it should be possible, with further excitation of the inner electron, to produce long-lived two-electron-excited planetary atoms.

Research supported by the NSF, The Robert A. Welch Foundation, the FWF (Austria), and the SFB-NextLite.

### References

- [1] G. Tanner, K. Richter, and J.-M. Rost 2000 *Rev. Mod. Phys.* **72**, 497
- [2] B. Wyker, S. Ye, T.J. McKinney, F.B. Dunning, S. Yoshida, C.O. Reinhold, and J. Burgdörfer 2011 *J. Phys. B :At. Mol. Opt. Phys* **44**, 184002

E-mail: [fbd@rice.edu](mailto:fbd@rice.edu)



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.