



Strathprints Institutional Repository

Davydov, O. and Schumaker, L.L. (2002) *On stable local bases for bivariate polynomial spline spaces*. Constructive Approximation, 18 (1). pp. 87-116. ISSN 0176-4276

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: <mailto:strathprints@strath.ac.uk>



Davydov, O. and Schumaker, L.L. (2001) On stable local bases for bivariate polynomial spline spaces. *Constructive Approximation*, 18 (1). pp. 87-116. ISSN 0176-4276

<http://eprints.cdlr.strath.ac.uk/4542/>

This is an author-produced version of a paper published in *Constructive Approximation*, ISSN 0176-4276.

This version has been peer-reviewed, but does not include the final publisher proof corrections, published layout, or pagination.

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in Strathprints to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profitmaking activities or any commercial gain. You may freely distribute the url (<http://eprints.cdlr.strath.ac.uk>) of the Strathprints website.

Any correspondence concerning this service should be sent to The Strathprints Administrator: eprints@cis.strath.ac.uk

On Stable Local Bases for Bivariate Polynomial Spline Spaces

Oleg Davydov ¹⁾ and Larry L. Schumaker ²⁾

Abstract. Stable locally supported bases are constructed for the spaces $\mathcal{S}_d^r(\Delta)$ of polynomial splines of degree $d \geq 3r + 2$ and smoothness r defined on triangulations Δ , as well as for various superspline subspaces. In addition, we show that for $r \geq 1$, it is impossible to construct bases which are simultaneously stable and locally linearly independent.

§1. Introduction

This paper deals with the classical space of polynomial splines

$$\mathcal{S}_d^r(\Delta) := \{s \in C^r(\Omega) : s|_T \in \mathcal{P}_d \text{ for all triangles } T \in \Delta\},$$

where \mathcal{P}_d is the space of polynomials of degree d , and Δ is a regular triangulation of a polygonal set Ω . We also discuss superspline subspaces of the form

$$\mathcal{S}_d^{r,\rho}(\Delta) := \{s \in \mathcal{S}_d^r(\Delta) : s \in C^{\rho_v}(v) \text{ for all } v \in \mathcal{V}\}, \quad (1.1)$$

with $\rho := \{\rho_v\}_{v \in \mathcal{V}}$, where ρ_v are given integers such that $r \leq \rho_v \leq d$, and \mathcal{V} is the set of all vertices of Δ .

Our aim is to describe algorithms for constructing locally supported bases $\{B_i\}_{i \in \mathcal{I}}$ for these spaces which are stable in the following sense:

$$K_1 \|c\|_\infty \leq \left\| \sum_{i \in \mathcal{I}} c_i B_i \right\|_\infty \leq K_2 \|c\|_\infty \quad (1.2)$$

for all choices of the coefficient vector $c = (c_i)_{i \in \mathcal{I}}$. We are interested in a construction for which (1.2) holds with constants K_1 and K_2 which depend only on d and the smallest angle θ_Δ in the triangulation, and not on the number of triangles or any other property of Δ . Stable bases are of critical importance for both theoretical and practical purposes.

¹⁾ Mathematical Institute, Justus Liebig University, D-35392 Giessen, Germany, oleg.davydov@math.uni-giessen.de

²⁾ Department of Mathematics, Vanderbilt University, Nashville, TN 37240, s@mars.cas.vanderbilt.edu. Supported by the National Science Foundation under grant DMS-9803340 and by the Army Research Office under grant DAAD-19-99-1-0160.

To date, algorithms which produce stable local bases have been described only for certain very special spline spaces, see Remarks 13.2–13.5. In this paper we give algorithms to construct locally supported stable bases for general superspline spaces $\mathcal{S}_d^{r,\rho}(\Delta)$ with $d \geq 3r + 2$. Such bases for the full spline spaces $\mathcal{S}_d^r(\Delta)$ (which correspond to the choice $\rho_v = r$ for all $v \in \mathcal{V}$) are of special importance due to the fact that these spaces are nested for nested triangulations, while most superspline spaces are not, see Remark 13.6.

The paper is organized as follows. In Sects. 2–4 we review the Bernstein-Bézier framework for dealing with bivariate splines and the minimal determining set approach to constructing bases for spline spaces. In Sect. 5 we describe how to construct corresponding dual basis splines. Sect. 6 contains a special construction of minimal determining sets for superspline spaces defined on near-singular cells. Sects. 7 and 8 deal with near-degenerate edges and near-singular vertices. In Sect. 9 we construct stable local bases for the superspline space $\mathcal{S}_d^{r,\mu}(\Delta)$ defined by choosing $\rho_v = \mu$ for all $v \in \mathcal{V}$, where

$$\mu := r + \left\lfloor \frac{r+1}{2} \right\rfloor. \quad (1.3)$$

In Sect. 10 we consider superspline spaces on arbitrary cells. The main result for general superspline spaces $\mathcal{S}_d^{r,\rho}(\Delta)$ is established in Sect. 11, and the connection between stability and local linear independence is explored in Sect. 12. We conclude the paper with several remarks in Sect. 13.

§2. The B-form of polynomials

In this section we briefly recall some well-known notation. Let $T := \langle u, v, w \rangle$ be a triangle in the plane. Throughout the paper, whenever we write a triangle T in terms of its vertices, we assume that u, v, w appear in counterclockwise order.

Given an integer d , we write $\{B_{ijk}^d\}_{i+j+k=d}$ for the usual Bernstein polynomials of degree d associated with T . Then it is well known (cf. [12]) that every polynomial $p \in \mathcal{P}_d$ can be written uniquely in the form

$$p = \sum_{i+j+k=d} c_{ijk} B_{ijk}^d. \quad (2.1)$$

The expansion (2.1) is called the Bernstein Bézier (B-) representation of p , and the c_{ijk} are called the B-coefficients. It is known that the B_{ijk}^d form a stable basis for \mathcal{P}_d , cf. Lemma 4.1 of [25].

We follow the standard convention of associating the B-coefficient c_{ijk} with the domain point

$$\xi_{ijk}^T := (iu + jv + kw)/d.$$

Thus, the coefficients of p can be indexed with the set

$$\mathcal{D}_{d,T} := \{\xi_{ijk}^T\}_{i+j+k=d}. \quad (2.2)$$

§3. The B-form of splines

Given a triangulation Δ , it is well known that there is a one–one correspondence between the spline space $\mathcal{S}_d^0(\Delta)$ and the set of coefficients $\{c_\xi\}_{\xi \in \mathcal{D}_{d,\Delta}}$, where

$$\mathcal{D}_{d,\Delta} := \bigcup_{T \in \Delta} \mathcal{D}_{d,T}. \quad (3.1)$$

In particular, $\{c_\xi\}_{\xi \in T \cap \mathcal{D}_{d,\Delta}}$ are the B-coefficients of $s|_T$.

This means that the set $\mathcal{D}_{d,\Delta}$ can be used to parametrize $\mathcal{S}_d^0(\Delta)$. It is common to write λ_ξ for the linear functional which picks off the B-coefficient c_ξ of $s \in \mathcal{S}_d^0(\Delta)$.

We recall some additional standard notation. Given $T = \langle u, v, w \rangle$, we define the distance of the domain point ξ_{ijk}^T from the vertex u to be $\text{dist}(\xi_{ijk}^T, u) := d - i$, with similar definitions for the other two vertices. We define the distance of ξ_{ijk}^T from the edge $\langle v, w \rangle$ to be i , with similar definitions for the other two edges.

Given a vertex u , the ring of radius m around u is the set $R_m(u) := \{\eta : \text{dist}(\eta, u) = m\}$. The disk of radius m around u is $D_m(u) := \{\eta : \text{dist}(\eta, u) \leq m\}$. We also define the arc $a_{m,e}^r(u)$ around u associated with an edge $e := \langle u, w \rangle$ to be the set of domain points in the ring $R_m(u)$ whose distance to $\langle u, w \rangle$ is at most r .

§4. Minimal determining sets

Suppose \mathcal{S} is a linear subspace of $\mathcal{S}_d^0(\Delta)$, and that $\mathcal{M} \subseteq \mathcal{D}_{d,\Delta}$. Then \mathcal{M} is said to be a determining set for \mathcal{S} if setting the coefficients of $s \in \mathcal{S}$ associated with the domain points in \mathcal{M} to zero implies that all coefficients of s corresponding to domain points in $\mathcal{D}_{d,\Delta}$ are zero. \mathcal{M} is called a minimal determining set (MDS) for \mathcal{S} if no proper subset $\mathcal{M}' \subset \mathcal{M}$ is a determining set. It is well known that \mathcal{M} is an MDS if and only if, given any real numbers $\{c_\xi\}_{\xi \in \mathcal{M}}$, there exists a unique spline $s \in \mathcal{S}$ whose B-coefficients in \mathcal{M} are $\{c_\xi\}_{\xi \in \mathcal{M}}$.

In this paper we are interested in subspaces of $\mathcal{S}_d^0(\Delta)$ which are defined by requiring that the B-coefficients $\{c_\xi\}_{\xi \in \mathcal{D}_{d,\Delta}}$ satisfy smoothness conditions across edges or at vertices. If \mathcal{M} is a MDS for such a spline space \mathcal{S} , then setting the B-coefficients c_ξ for all $\xi \in \mathcal{M}$, we can compute all remaining B-coefficients using the smoothness conditions. Indeed, suppose s is a spline in $\mathcal{S}_d^0(\Delta)$, and let $T := \langle v_1, v_2, v_3 \rangle$ and $\tilde{T} := \langle v_4, v_3, v_2 \rangle$ be a pair of adjoining triangles in Δ sharing the edge $e = \langle v_2, v_3 \rangle$. Let c_{ijk} and \tilde{c}_{ijk} be the B-coefficients of $s|_T$ and $s|_{\tilde{T}}$, respectively. Then it is well known (cf. [12,22]) that s is C^r continuous across the edge e if and only if

$$\tilde{c}_{n,m-n,d-m} = \sum_{i+j+k=n} c_{i,j+d-m,k+m-n} B_{ijk}^n(v_4), \quad (4.1)$$

for $m = n, \dots, d$ and $n = 1, \dots, r$. Here B_{ijk}^n are the Bernstein polynomials of degree n on the triangle T . We recall the following two lemmas from [25].

Lemma 4.1. *Suppose s is a spline in $\mathcal{S}_d^r(\Delta)$, and that for fixed $n \leq m \leq d$ and $1 \leq n \leq r$ the coefficients appearing on the right-hand side of (4.1) are known, and that C is the maximum of their absolute values. Then $|c_{n,m-n,d-m}| \leq KC$, where K is a constant depending only on d and the smallest angle in the triangulation.*

Lemma 4.2. *Suppose T and \tilde{T} are two triangles as above, and suppose*

$$v_4 = \alpha v_1 + \beta v_2 + \gamma v_3. \quad (4.2)$$

with $\gamma \neq 0$. Suppose we know the coefficients of a spline $s \in \mathcal{S}_d^r(\Delta)$ for all domain points in the disk $D_{m-1}(v_2)$ with $m \geq r+1$. Let $c_i := c_{i,d-m,m-i}^T$ be the coefficients of $p := s|_T$ on the arc $a_{m,e}^r(v_2)$ associated with the edge $e := \langle v_2, v_3 \rangle$, and let $\tilde{c}_i := c_{i,m-i,d-m}^{\tilde{T}}$ be those of $\tilde{p} := s|_{\tilde{T}}$ on the same arc. Suppose that the coefficients c_i and \tilde{c}_i are known for $i \in \{r-q+1, \dots, r\}$ and that the coefficients c_0, \dots, c_{r-2q} are also known for some q with $r+1 \geq 2q$. Then the coefficients c_i and \tilde{c}_i are uniquely determined for all $0 \leq i \leq r$. Moreover, if C is the maximum of the known coefficients, then the computed coefficients are bounded by KC , where K is a constant depending only on d , the smallest angle in the triangulation, and the size of γ^{-1} .

Lemma 4.2 cannot be applied in the situation when $\gamma = 0$. In this case the edge $e := \langle v_2, v_3 \rangle$ is called *degenerate*. The stability of the computation is also compromised whenever e is near-degenerate in the sense that γ is near zero. We discuss near-degenerate edges in more detail in Sect. 7.

§5. Dual bases

Our construction of stable local bases will be based on the following

Lemma 5.1. [6] *Suppose \mathcal{M} is a MDS for a space of splines \mathcal{S} which is a subset of $\mathcal{S}_d^0(\Delta)$ defined by smoothness conditions. For each $\xi \in \mathcal{M}$, let B_ξ be the unique spline in \mathcal{S} satisfying*

$$\lambda_\eta B_\xi = \delta_{\xi,\eta}, \quad \text{all } \eta \in \mathcal{M}, \quad (5.1)$$

where λ_η is the linear functional which picks off the B -coefficient c_η . Then the set $\{B_\xi\}_{\xi \in \mathcal{M}}$ is a basis for \mathcal{S} . We call it the dual basis corresponding to \mathcal{M} .

Proof: To construct the spline B_ξ , choose $c_\xi = 1$, and set all other coefficients corresponding to $\eta \in \mathcal{M}$ to zero. Then since \mathcal{M} is a MDS, all remaining coefficients are uniquely determined. Property (5.1) is obvious. \square

For a given spline space \mathcal{S} , there are generally many different minimal determining sets \mathcal{M} . Our aim in this paper is to design algorithms for *choosing minimal determining sets* which produce stable local bases when applied to a space $\mathcal{S} \subseteq \mathcal{S}_d^0(\Delta)$.

Definition 5.2. Suppose $\mathcal{B} := \{B_\xi\}_{\xi \in \mathcal{M}}$ is a basis for a spline space $\mathcal{S} \subseteq \mathcal{S}_d^0(\Delta)$. Then we call \mathcal{B} a stable local basis provided that there exist constants ℓ and K depending only on d and the smallest angle θ_Δ in Δ such that

$$\text{supp}(B_\xi) \subseteq \text{star}^\ell(v_\xi) \text{ for some vertex } v_\xi, \quad (5.2)$$

$$\|B_\xi\|_\infty \leq K \quad (5.3)$$

for all $\xi \in \mathcal{M}$. A minimal determining set \mathcal{M} for a spline space \mathcal{S} is called a stable local MDS provided that the corresponding dual basis $\mathcal{B} := \{B_\xi\}_{\xi \in \mathcal{M}}$ is a stable local basis for \mathcal{S} .

For (5.2), we recall that given a vertex v of Δ , $\text{star}(v) = \text{star}^1(v)$ is the union of triangles sharing v , and $\text{star}^\ell(v)$, $\ell \geq 2$, is defined recursively as the union of the stars of the vertices in $\mathcal{V} \cap \text{star}^{\ell-1}(v)$.

Theorem 5.3. Suppose that \mathcal{M} is a stable local minimal determining set for a spline space $\mathcal{S} \subseteq \mathcal{S}_d^0(\Delta)$. Then the dual basis \mathcal{B} satisfies (1.2).

Proof: Let $s \in \mathcal{S}$. Then for any $\xi \in \mathcal{M}$, the corresponding coefficient c_ξ is the B-coefficient of the polynomial $s_T := s|_T$, where T is a triangle containing ξ . But then by the stability of the Bernstein-Bézier basis for polynomials (cf. Lemma 4.1 of [25]), we have $|c_\xi| \leq C \|s_T\|_{\infty, T} \leq C \|s\|_\infty$, where C is a constant which depends only on d . This establishes the left-hand side of (1.2) with the constant $K_1 := 1/C$ which depends only on d . For the right-hand side, we note that by Lemma 3.1 of [25], for any triangle T , the number of basis splines B_ξ whose supports contain T is bounded by a constant depending only on d and θ_Δ (cf. the proof of Theorem 9.2 in [25]). Combining this with the boundedness of the basis functions completes the proof. \square

With an appropriate renorming, it can be shown that the dual splines also provide a basis which is L_p -stable for $1 \leq p < \infty$, see Remark 13.7.

§6. Four-cells

In this section we construct minimal determining sets for spline spaces defined on triangulations which consist of exactly four triangles surrounding one interior vertex v . We call such a triangulation Δ_v a four-cell. We are particularly interested in the case where the vertex is singular (formed by the intersection of two straight lines) or near-singular.

Suppose v_1, \dots, v_4 are the boundary vertices of Δ_v in counterclockwise order. Let $T_i := \langle v, v_i, v_{i+1} \rangle$, for $i = 1, \dots, 4$, where $v_5 = v_1$. For $\mu + 1 \leq \ell \leq 2r$, we introduce some simplified notation for certain domain points on the ring $R_\ell(v)$. Let

$$\begin{aligned} a_{\ell,j}^i &:= \xi_{d-\ell, \ell-r+j-1, r-j+1}^{T_i}, \quad 1 \leq j \leq n_\ell, \\ g_{\ell,j}^i &:= \xi_{d-\ell, \ell-r+n_\ell+j-1, r-n_\ell-j+1}^{T_i}, \quad 1 \leq j \leq n_\ell, \\ d_{\ell,j}^i &:= \xi_{d-\ell, \ell-r+2n_\ell+j-1, r-2n_\ell-j+1}^{T_i}, \quad 1 \leq j \leq r - 2n_\ell + 1, \end{aligned} \quad (6.1)$$




Fig. 1. The points in (6.1) for $r = 4$, $\mu = 6$, $d = 14$.

where

$$n_\ell := 2r + 1 - \ell. \quad (6.2)$$

Note that $n_\ell \geq 1$ and $r - 2n_\ell + 1 \geq 1$. We illustrate this notation for $r = 4$, $\mu = 6$, and $d = 14$ in Fig. 1.

Let $1 \leq k \leq 2r$. We say that $\mathcal{M} \subseteq D_k(v)$ is a MDS for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$ if setting the coefficients $\{c_\xi : \xi \in \mathcal{M}\}$ to arbitrary real numbers, the coefficients $\{c_\xi : \xi \in D_k(v) \setminus \mathcal{M}\}$ can be uniquely computed by using those smoothness conditions defining $\mathcal{S}_d^{r,\mu}(\Delta_v)$ that do not involve c_ξ for $\xi \notin D_k(v)$. Such an MDS will be called stable if

$$\max_{\xi \in D_k(v) \setminus \mathcal{M}} |c_\xi| \leq K \max_{\xi \in \mathcal{M}} |c_\xi|,$$

where K is a constant depending only on d and the smallest angle θ_Δ in Δ_v . By Lemma 4.1 it is easy to see that $D_k^{T_1}(v)$ is a stable MDS for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$ if

$k \leq \mu$.

It is not difficult to describe a stable MDS for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$ for $\mu + 1 \leq k \leq 2r$ if v is singular.

Theorem 6.1. *Suppose Δ_v is a four-cell associated with a singular vertex v . For each $\ell = \mu + 1, \dots, 2r$, let*

$$\begin{aligned} \mathcal{M}_{v,\ell} &:= \{a_{\ell,1}^1, \dots, a_{\ell,n_\ell}^1\} \cup \bigcup_{i=1}^4 \{g_{\ell,1}^i, \dots, g_{\ell,n_\ell}^i\}, \\ O_{v,\ell} &:= \bigcup_{i=1}^4 \{d_{\ell,1}^i, \dots, d_{\ell,r-2n_\ell+1}^i\}. \end{aligned} \tag{6.3}$$

Then for each $k = \mu, \dots, 2r$,

$$\Gamma_k := D_\mu^{T_1}(v) \cup \bigcup_{\ell=\mu+1}^k [\mathcal{M}_{v,\ell} \cup O_{v,\ell}] \tag{6.4}$$

is a stable MDS for the space $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$.

Proof: We proceed by induction on k . The result is clear for $k = \mu$. Suppose we set the coefficients c_ξ of $s \in \mathcal{S}_d^{r,\mu}(\Delta_v)$ for $\xi \in \Gamma_k$. Then by the inductive hypothesis, all coefficients c_ξ with $\xi \in D_{k-1}(v)$ are uniquely determined by those with $\xi \in \Gamma_{k-1} \subseteq \Gamma_k$. We then compute the coefficients associated with domain points on the ring $R_k(v)$ using the standard smoothness conditions as in Lemma 4.1. Namely, for each $i = 1, 2, 3, 4$, we first use the coefficients associated with the domain points $g_{k,1}^{i+1}, \dots, g_{k,n_k}^{i+1}, d_{k,1}^{i+1}, \dots, d_{k,r-2n_k+1}^{i+1}$ and those in $D_{k-1}(v)$ to compute the coefficients corresponding to $\{\xi_{d-k,0,k}^{T_i}, \dots, \xi_{d-k,r-n_k,k-r+n_k}^{T_i}\}$. After this only the coefficients associated with domain points $a_{k,1}^i, \dots, a_{k,n_k}^i$, $i = 2, 3, 4$, remain undetermined in $R_k(v)$. We therefore use the coefficients corresponding to $R_k^{T_1}(v)$ to compute the coefficients c_ξ with $\xi \in \{a_{k,1}^2, \dots, a_{k,n_k}^2\}$, and, proceeding counter-clockwise around v , successively compute the coefficients with $\xi \in \{a_{k,1}^3, \dots, a_{k,n_k}^3\}$ and $\xi \in \{a_{k,1}^4, \dots, a_{k,n_k}^4\}$. Note that here we have not used a portion of the smoothness conditions across the edge $e_1 := \langle v, v_1 \rangle$ which involve the coefficients c_ξ for $\xi \in \{a_{k,1}^1, \dots, a_{k,n_k}^1\}$. Nevertheless, these conditions must be satisfied since the number of free parameters c_ξ , $\xi \in \Gamma_k \setminus \Gamma_{k-1}$, used in the above computation on ring $R_k(v)$, is equal to

$$\dim \mathcal{S}_k^{r,\mu}(\Delta_v) - \dim \mathcal{S}_{k-1}^{r,\mu}(\Delta_v) = 4(k-r) + n_k$$

(cf. Theorem 2.2 of [30]). Thus, we are able to compute all coefficients c_ξ , $\xi \in D_k(v) \setminus \Gamma_k$, by applying Lemma 4.1 several times. By that lemma, the maximum of the computed coefficients is bounded by a constant K times the maximum of the set coefficients, where K depends only on d and the smallest angle in Δ_v . \square

For later use in building stable local minimal determining sets for general spline spaces, it is critical that the stable MDS in Theorem 6.1 contains the sets $O_{v,\ell}$. We now extend this result to arbitrary four-cells, although in fact we will use it only for cells which are near-singular. In Sect. 10 we construct stable minimal determining sets for supersplines on general cells. The construction there is simpler, but does not guarantee that the resulting MDS contains the needed sets $O_{v,\ell}$.

Theorem 6.2. *Suppose Δ_v is a four-cell associated with a nonsingular vertex v . For each $\mu + 1 \leq \ell \leq 2r$, let $O_{v,\ell}$ be the sets in (6.3), and let n_ℓ be the integer defined in (6.2). Then there exists a set of $4n_\ell$ domain points*

$$\mathcal{M}_{v,\ell} \subseteq \mathcal{A}_{v,\ell} := \bigcup_{i=1}^4 \left[\{a_{\ell,j}^i\}_{j=1}^{n_\ell} \cup \{g_{\ell,j}^i\}_{j=1}^{n_\ell} \right]. \quad (6.5)$$

such that for each $k = \mu, \dots, 2r$,

$$\Gamma_k := D_\mu^{T_1}(v) \cup \bigcup_{\ell=\mu+1}^k [\mathcal{M}_{v,\ell} \cup O_{v,\ell}] \quad (6.6)$$

is a stable MDS for the space $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$.

Proof: We proceed by induction on k . The statement of the theorem holds for $k = \mu$ since $\Gamma_\mu = D_\mu^{T_1}(v)$ is trivially a stable MDS for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_\mu(v)$.

Fix $\mu + 1 \leq k \leq 2r$, and suppose that Γ_{k-1} is a stable MDS for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_{k-1}(v)$. To construct Γ_k which is a stable MDS for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$, we need to supplement Γ_{k-1} with an appropriate subset of the domain points on the ring $R_k(v)$. Using the fact that v is not a singular vertex, it is easy to see that the number of edges attached to v with different slopes is at least three. Then Theorem 2.2 of [30] implies

$$m := \dim \mathcal{S}_k^{r,\mu}(\Delta_v) - \dim \mathcal{S}_{k-1}^{r,\mu}(\Delta_v) = 4(k - r). \quad (6.7)$$

Thus, to get a minimal determining set Γ_k for $\mathcal{S}_d^{r,\mu}(\Delta_v)$ on $D_k(v)$, we need to add to Γ_{k-1} exactly m points on the ring $R_k(v)$.

To simplify the discussion of how to choose these m points, we first reduce the problem to one of considering splines whose coefficients are zero for all points in the disk $D_{k-1}(v)$. Given $s \in \mathcal{S}_d^{r,\mu}(\Delta_v)$, let $\mathcal{T}_{k-1}s$ be the spline in $\mathcal{S}_d^r(\Delta_v)$ constructed in Lemma 6.4 below such that for each triangle attached to v , $g_T := \mathcal{T}_{k-1}s|_T$ interpolates the derivatives up to order $k-1$ of $s|_T$ at v . Note that since $s \in C^\mu(v)$, $\mathcal{T}_{k-1}s$ is also in $C^\mu(v)$. Then the spline $\hat{s} := s - \mathcal{T}_{k-1}s \in \mathcal{S}_d^{r,\mu}(\Delta_v)$ has all zero coefficients in $D_{k-1}(v)$. Computing its coefficients on the ring $R_k(v)$ will stably and uniquely determine the coefficients of s on $R_k(v)$, since by Lemma 6.4 the size of the coefficients of $\mathcal{T}_{k-1}s$ on this ring is bounded by the size of the coefficients of s in $D_{k-1}(v)$.

$$H_i^d := \begin{pmatrix} \binom{r-n+1}{r-2n} r_i^{r-2n} t_i^{n+1} & \cdots & \binom{r-n+1}{1} r_i t_i^{r-n} & t_i^{r-n+1} \\ \vdots & & \vdots & \vdots \\ \binom{r}{r-2n} r_i^{r-2n} t_i^{2n} & \cdots & \binom{r}{1} r_i t_i^{r-1} & t_i^r \end{pmatrix},$$

and I is the $n \times n$ identity matrix. We call a column of H a d -column when it passes through one of the matrices H_i^d . We define a -columns and g -columns similarly.

The matrix H has $4n$ rows and $4(r+1)$ columns where $n < r+1$. We claim that it has full rank $4n$. Indeed, the number of independent solutions $4(r+1) - \text{rank}(H)$ of the homogeneous system (6.8) must be equal to m , which implies $\text{rank}(H) = 4n$. This means that there is a choice of $4n$ indices $1 \leq i_1 < \cdots < i_{4n} \leq 4r+4$ such that the determinant of the corresponding square submatrix $H(i_1, \dots, i_{4n})$ is not zero. Moreover, it follows from Lemma 6.3 below that we can choose i_1, \dots, i_{4n} such that no column of $H(i_1, \dots, i_{4n})$ is a d -column.

We are ready to describe the set $\mathcal{M}_{v,k}$. Denote by Σ the set of indices of all a - and g -columns of H , and let $\{i_1^*, \dots, i_{4n}^*\} \subseteq \Sigma$ be such that

$$|\det H(i_1^*, \dots, i_{4n}^*)| = \max_{i_1, \dots, i_{4n} \in \Sigma} |\det H(i_1, \dots, i_{4n})|. \quad (6.9)$$

We take $\mathcal{M}_{v,k}$ to be the set of domain points in $\mathcal{A}_{v,k}$ which correspond to the columns with indices in the set $\Sigma \setminus \{i_1^*, \dots, i_{4n}^*\}$. Then $\mathcal{M}_{v,k} \cup \mathcal{O}_{v,k}$ is the set of domain points on $R_k(v)$ which correspond to the columns of H with indices in the set $J^* := \{1, \dots, 4r+4\} \setminus \{i_1^*, \dots, i_{4n}^*\}$.

Now assuming that the coefficients $\{z_j\}_{j \in J^*}$ of \hat{s} corresponding to points in $\mathcal{M}_{v,k} \cup \mathcal{O}_{v,k}$ have been set, we may compute the remaining coefficients corresponding to points in $\mathcal{A}_{v,k} \cup \mathcal{O}_{v,k}$ from the nonsingular system

$$H(i_1^*, \dots, i_{4n}^*) \begin{pmatrix} z_{i_1^*} \\ \vdots \\ z_{i_{4n}^*} \end{pmatrix} = - \sum_{j \in J^*} z_j H(j), \quad (6.10)$$

where $H(j)$ is the j -th column of H . Using Cramer's rule and taking account of (6.9) and Lemma 6.3, we conclude that

$$|z_{i_\nu^*}| \leq \frac{\sum_{j \in J^*} |z_j| |\det H(i_1^*, \dots, i_{\nu-1}^*, j, i_{\nu+1}^*, \dots, i_{4n}^*)|}{|\det H(i_1^*, \dots, i_{4n}^*)|} \leq K \max_{j \in J^*} |z_j|,$$

for $\nu = 1, \dots, 4n$, where K is a constant depending only on d and the smallest angle in Δ_v . This shows that the computation of $z_{i_1^*}, \dots, z_{i_{4n}^*}$ is stable. \square

We now state and prove two lemmas which were used in the proof of Theorem 6.2. The first result concerns determinants formed from $4n \times 4n$ submatrices of H . Let

$$R := \begin{pmatrix} \binom{r-n+1}{r-n} & \cdots & \binom{r-n+1}{r-2n+1} \\ \vdots & \ddots & \vdots \\ \binom{r}{r-n} & \cdots & \binom{r}{r-2n+1} \end{pmatrix}.$$

A simple computation shows that

$$\det R = C \det \begin{pmatrix} \frac{1}{1!} & \cdots & \frac{1}{n!} \\ \vdots & \cdots & \vdots \\ \frac{1}{n!} & \cdots & \frac{1}{(2n-1)!} \end{pmatrix},$$

where C is a positive constant depending only on r and n . It is well-known that this determinant is nonzero for all choices of n , and thus the matrix R is nonsingular.

Lemma 6.3. *Let $H(i_1, \dots, i_{4n})$ be a $4n \times 4n$ submatrix of H containing a nontrivial d -column. Then there exists another submatrix $H(j_1, \dots, j_{4n})$ with one less d -column such that*

$$|\det H(i_1, \dots, i_{4n})| \leq C |\det H(j_1, \dots, j_{4n})|, \quad (6.11)$$

where $C > 0$ is a constant depending only on d and the smallest angle of Δ_v .

Proof: Suppose $H(i_1, \dots, i_{4n})$ includes a nontrivial d -column $i_p = (r+1)(i-1) + 2n+j$ with $1 \leq i \leq 4$ and $1 \leq j \leq r-2n+1$. Note that the column is nontrivial if and only if the corresponding t_i is nonzero. For any $1 \leq j \leq r-2n+1$, it is not difficult to see that

$$H_i^d(j) = \sum_{\kappa=1}^n x_\kappa^{[j]} \left(\frac{t_i}{r_i} \right)^{j+n-\kappa} H_i^g(\kappa),$$

where the numbers $x_\kappa^{[j]}$ are determined from the nonsingular linear system

$$R \begin{pmatrix} x_1^{[j]} \\ \vdots \\ x_n^{[j]} \end{pmatrix} = \begin{pmatrix} \binom{r-n+1}{r-2n+1-j} \\ \vdots \\ \binom{r}{r-2n+1-j} \end{pmatrix}. \quad (6.12)$$

Since the κ -th column of H_i^g corresponds to the $(r+1)(i-1) + n + \kappa$ -th column of H , this implies that

$$\det H(i_1, \dots, i_{4n}) = \sum_{\kappa=1}^n x_\kappa^{[j]} \left(\frac{t_i}{r_i} \right)^{j+n-\kappa} \det H_\kappa,$$

where

$$H_\kappa := H(i_1, \dots, i_{p-1}, (r+1)(i-1) + n + \kappa, i_{p+1}, \dots, i_{4n}).$$

Since $|r_i|$ is the quotient of the areas of two neighboring triangles T_{i-1} and T_i , we have by Lemma 3.2 of [25],

$$0 < K_1 \leq |r_i| \leq K_2, \quad (6.13)$$

where K_1, K_2 depend only on θ_Δ . Therefore,

$$|\det H(i_1, \dots, i_{4n})| \leq K_3 |t_i|^j \max_{\kappa} |\det H_\kappa|, \quad (6.14)$$

where K_3 depends only on d and θ_Δ . The result follows since $|t_i| \leq K_4$ where K_4 is a constant depending only on θ_Δ . (In fact, $|t_i|$ is quite small if v is near-singular.)

□

The following lemma was used in the proof of Theorem 6.2 above, and will also be useful in Sect. 12 below.

Lemma 6.4. Let Δ_v be a cell, and let $0 \leq r < k \leq d$ be integers. Given a spline $s \in \mathcal{S}_d^r(\Delta_v)$, let $\mathcal{T}_{k-1}s$ be such that for each triangle T attached to v ,

$\mathcal{T}_{k-1}s|_T :=$ the unique polynomial of degree $k-1$ which matches
the derivatives of $s|_T$ at v up to order $k-1$.

Then $\mathcal{T}_{k-1}s \in \mathcal{S}_{k-1}^r(\Delta_v) \subseteq \mathcal{S}_d^r(\Delta_v)$. Moreover, if

$$s|_T = \sum c_\xi^T B_\xi^T, \quad \mathcal{T}_{k-1}s|_T = \sum \hat{c}_\xi^T B_\xi^T,$$

where B_ξ^T are the Bernstein polynomials of degree d associated with a triangle T , then $\hat{c}_\xi^T = c_\xi^T$ for all $\xi \in D_{k-1}^T(v)$, and

$$\max_{\xi \in R_k^T(v)} |\hat{c}_\xi^T| \leq K \max_{\xi \in D_{k-1}^T(v)} |c_\xi^T|, \quad (6.15)$$

where K is a constant depending only on d .

Proof: Comparing cross derivatives of neighboring pieces of $\mathcal{T}_{k-1}s$, it is easy to see that it satisfies C^r smoothness conditions across the interior edges of Δ_v , and thus is a spline in $\mathcal{S}_{k-1}^r(\Delta_v) \subseteq \mathcal{S}_d^r(\Delta)$. Now fix a triangle $T := \langle v, v_i, v_{i+1} \rangle$ in Δ_v . Then by the well-known connection between derivatives and coefficients of a polynomial written in Bernstein-Bézier form, it follows that $\hat{c}_\xi^T = c_\xi^T$ for all $\xi \in D_{k-1}^T(v)$. Finally, to establish (6.15), we observe that since $\mathcal{T}_{k-1}s$ is a polynomial of degree $k-1$, its k -th derivatives are identically zero, and thus for all $\nu = 0, \dots, k$,

$$\begin{aligned} 0 &= D_{v_i-v}^\nu D_{v_{i+1}-v}^{k-\nu} \mathcal{T}_{k-1}s|_T(v) \\ &= \frac{d!}{(d-k)!} \sum_{j_1=0}^\nu \sum_{j_2=0}^{k-\nu} \binom{\nu}{j_1} \binom{k-\nu}{j_2} (-1)^{k-j_1-j_2} \hat{c}_{d-j_1-j_2, j_1, j_2}^T. \end{aligned}$$

It follows that

$$\hat{c}_{d-k, \nu, k-\nu}^T = -\frac{d!}{(d-k)!} \sum_{\substack{0 \leq j_1 \leq \nu, 0 \leq j_2 \leq k-\nu \\ j_1+j_2 \leq k-1}} \binom{\nu}{j_1} \binom{k-\nu}{j_2} (-1)^{k-j_1-j_2} \hat{c}_{d-j_1-j_2, j_1, j_2}^T,$$

which immediately implies (6.15). \square

Example 6.5. Let $r = 5$, $\mu = 8$, $d = 17$ in Theorem 6.2.

Discussion: Fig. 2 shows the domain points in this case, where the D_{2r} disks are shaded light gray and the D_μ disks are shaded dark gray. We concentrate on the ring $R_9(v)$, where $n_9 = 2$. The points in $O_{v,9}$ are marked with the symbol \odot (except for the point $d_{9,2}^3$ which we have marked with a \oplus for a later discussion). In this case the set $\mathcal{M}_{v,9}$ must contain eight of the sixteen points in the set $\mathcal{A}_{v,9}$ described in (6.5). These eight points are chosen by the method of maximization of the determinant in (6.9), and therefore depend on the exact geometry of the cell. In Fig. 2 we show a possible constellation where we mark the eight points in $\mathcal{M}_{v,9} \cap \mathcal{A}_{v,9}$ with the symbol \blacksquare . This leaves eight points which are computed by the linear system (6.8). They are marked with boxed numbers 1 through 8. \square




Fig. 2. The points in $\mathcal{M} \cap R_9(v)$ for $r = 5$, $\mu = 8$, $d = 17$.

§7. Near-degenerate edges

We recall that an edge $e = \langle v_2, v_3 \rangle$ shared by two triangles $T := \langle v_1, v_2, v_3 \rangle$ and $\tilde{T} := \langle v_4, v_3, v_2 \rangle$ is called *degenerate* at v_2 provided that the points v_1, v_2, v_4 lie on a straight line. An edge which is nearly degenerate is called *near-degenerate*, cf. [25]. We shall introduce the following quantitative form, where the choice of the constant $2\theta^2/\pi$ is motivated by Lemma 8.2 below.

Definition 7.1. *Suppose $T := \langle v_1, v_2, v_3 \rangle$ and $\tilde{T} := \langle v_4, v_3, v_2 \rangle$ are two triangles which share the edge $e = \langle v_2, v_3 \rangle$. We say that e is θ -near-degenerate at v_2 provided that the smaller of the two angles between the edges $\langle v_2, v_1 \rangle$ and $\langle v_2, v_4 \rangle$ is greater than $\pi - 2\theta^2/\pi$.*

Let (α, β, γ) be the barycentric coordinates of v_4 in terms of the triangle T , i.e., $v_4 = \alpha v_1 + \beta v_2 + \gamma v_3$. The following lemma gives a lower bound on the size of γ

in the case where e is not θ -near-degenerate. This is important since it shows that for such edges e , the process of computing coefficients by smoothness conditions across e as in Lemma 4.2 is stable.

Lemma 7.2. *Suppose the smallest angle in the triangles T and \tilde{T} is at least θ , and that $e_2 := \langle v_2, v_3 \rangle$ is not θ -near-degenerate at v_2 . Then $|\gamma| \geq \sin(\theta) \sin(2\theta^2/\pi)$.*

Proof: Let $e_1 := \langle v_2, v_1 \rangle$ and $e_3 := \langle v_2, v_4 \rangle$. Then

$$|\gamma| = \frac{|e_1||e_3|\sin(a)}{|e_1||e_2|\sin(\theta_1)},$$

where a is the smaller of the two angles between e_1 and e_3 , and θ_1 is the angle between the edges e_1 and e_2 . It was shown in the proof of Lemma 3.2 of [25] that under the above hypotheses, $|e_3|/|e_2| \geq \sin(\theta)$. But then the result follows from the fact that $|\sin(a)| \geq \sin(2\theta^2/\pi)$ while $|\sin(\theta_1)| \leq 1$. \square

§8. Near-singular vertices

A singular vertex is an interior vertex of a triangulation which is formed by the intersection of exactly two lines. It is well known that singular and near-singular vertices complicate the construction of stable bases for spline spaces.

Definition 8.1. *Suppose v is a vertex where four edges meet. If all four edges are θ -near-degenerate at v , then we call v a θ -near-singular vertex.*

The following lemma will be used later.

Lemma 8.2. *Fix $\theta \leq \pi/4$, and let Δ be a triangulation with smallest angle at least θ . Then for any interior vertex v which is not θ -near-singular, there is at least one edge attached to v which is not θ -near-degenerate at either end.*

Proof: Let v_1, \dots, v_n be the vertices attached to v in counter-clockwise order. We set $T_i := \langle v, v_i, v_{i+1} \rangle$ and denote by $\theta_i, \phi_i, \omega_i$ the angles of T_i at v, v_i, v_{i+1} , respectively. We distinguish three cases.

Case 1: $n = 3$. Consider the edge $e_1 := \langle v, v_1 \rangle$, and let $\alpha := \phi_1 + \omega_3$ and $\beta := \theta_1 + \theta_3$. Then it is clear that $\beta \geq \pi + 2\theta$. Now $\alpha + \beta + \omega_1 + \phi_3 = 2\pi$. This implies $\alpha \leq \pi - 4\theta$ since $\omega_1, \phi_3 \geq \theta$.

Case 2: $n = 4$. Since v is not θ -near-singular, there is at least one edge attached to v which is not θ -near-degenerate at v . Without loss of generality we can assume it is the edge $e_1 := \langle v, v_1 \rangle$, and that the angle $\beta := \theta_1 + \theta_4$ is at least $\pi + 2\theta^2/\pi$. Then arguing as in Case 1, we see that $\alpha := \phi_1 + \omega_4 \leq \pi - 2\theta - 2\theta^2/\pi \leq \pi - 2\theta$.

Case 3: $n \geq 5$. Consider the edge $e_1 := \langle v, v_1 \rangle$. Let $\alpha_i := \phi_i + \omega_{i-1}$ and $\beta_i := \theta_i + \theta_{i-1}$, for $i = 1, \dots, n$ where we identify $\theta_{n+i} = \theta_i, \omega_{n+i} = \omega_i$. We claim

that at least three of the α_i satisfy $\alpha_i \leq \pi - 4\theta/(n-2)$. Indeed, if this were not the case, then

$$(n-2)\pi = \sum_{i=1}^n \alpha_i > (n-2)\left(\pi - \frac{4\theta}{n-2}\right) + 4\theta = (n-2)\pi.$$

On the other hand, we claim that at most two of the β_i satisfy $\beta_i \geq \pi - \theta/2$. Suppose to the contrary that there are three, say $\beta_k, \beta_l, \beta_m$. Then at least two of these do not overlap, say β_k, β_l . But then there are $n-4$ of the angles θ_i which are not covered by β_k or β_l , which would lead to the contradiction

$$2\pi = \beta_k + \beta_l + (n-4)\theta > 2\pi - \theta + (n-4)\theta \geq 2\pi.$$

Now $n\theta \leq 2\pi$ implies $4/(n-2) \geq 4/n \geq 2\theta/\pi$. We conclude that for one of the edges, $\alpha_i \leq \pi - 4\theta/(n-2) \leq \pi - 2\theta^2/\pi$ and $\beta_i \leq \pi - \theta/2 \leq \pi - 2\theta^2/\pi$. It follows that this edge is not θ -near-degenerate at either end. \square

§9. Stable local bases for $\mathcal{S}_d^{r,\mu}(\Delta)$

Our ultimate aim is to give stable local bases for the general superspline spaces (1.1) defined on arbitrary triangulations Δ of a polygonal set Ω . However, in order to illustrate the construction in a somewhat simpler setting, in this section we consider the superspline space

$$\mathcal{S}_d^{r,\mu}(\Delta) := \{s \in \mathcal{S}_d^r(\Delta) : s \in C^\mu(v) \text{ for all } v \in \mathcal{V}\},$$

for $d \geq 3r+2$, where μ is defined in (1.3). This is the special case of (1.1) with $\rho_v = \mu$ for all $v \in \mathcal{V}$. The analogous construction for general superspline spaces requires further analysis of cells (see Sect. 10), and is given in Section 11.

To describe a minimal determining set for $\mathcal{S}_d^r(\Delta)$ we need some additional notation. Let \mathcal{V}_S and \mathcal{V}_{NS} be the sets of vertices of Δ which are singular and θ_Δ -near-singular, respectively, where θ_Δ is the smallest angle in Δ . Given a triangle T , we divide the set $\mathcal{D}_{d,T}$ into various subsets. Let

$$C^T := \{\xi_{ijk}^T : i > r, j > r, k > r\}.$$

and

$$D_\mu^T(u) := \{\xi_{ijk}^T : i \geq d - \mu\}$$

$$A^T(u) := \bigcup_{i=1}^{\lfloor \frac{r}{2} \rfloor} \bigcup_{j=0}^{i-1} \{\xi_{d-2r+i-1, r-j, r-i+j+1}^T\},$$




Fig. 3. Domain points for $r = 4$, $\mu = 6$, $d = 15$.

with similar definitions for the other two vertices of T . Associated with the edge $e := \langle u, v \rangle$, let

$$\begin{aligned}
 F^T(e) &:= \{\xi_{ijk}^T : k \leq r\} \\
 G_L^T(e) &:= \bigcup_{i=1}^{\lfloor \frac{r}{2} \rfloor} \bigcup_{j=0}^{i-1} \{\xi_{d-2r+i-1, r+1+j, r-i-j}^T\} \\
 G_R^T(e) &:= \bigcup_{i=1}^{\lfloor \frac{r}{2} \rfloor} \bigcup_{j=0}^{i-1} \{\xi_{r+1+j, d-2r+i-1, r-i-j}^T\} \\
 E^T(e) &:= F^T(e) \setminus \left[D_\mu^T(u) \cup D_\mu^T(v) \cup A^T(u) \cup A^T(v) \cup G_L^T(e) \cup G_R^T(e) \right],
 \end{aligned} \tag{9.1}$$

with similar definitions for the other two edges of T . Note that $G_L^T(\langle u, v \rangle) = G_R^T(\langle v, u \rangle)$.

As an aid to visualizing where these points are located, in Fig. 3 we show them for the case where $r = 4$, $\mu = 6$, and $d = 15$. The points in the three disks $D_6(u)$, $D_6(v)$, and $D_6(w)$ are marked with the symbol \oplus . The points in the three sets $A^T(u)$, $A^T(v)$ and $A^T(w)$ are marked with stars, and the point in C^T is marked with a dark circle. The points in $E^T(e)$ for edge $e := \langle u, v \rangle$ are marked with triangles, and the points in $G_L^T(e)$ and $G_R^T(e)$ are marked with the symbols \otimes and open circle, respectively. For clarity, we have not marked points associated with the other two edges.

Theorem 9.1. *Let \mathcal{M} be the following set of domain points:*

- 1) *for each triangle T , include the set C^T ,*
- 2) *for each edge e , include the set $E^T(e)$, where T is some triangle sharing e ,*
- 3) *for each edge of a triangle T such that e lies on the boundary of Ω , include the sets $G_L^T(e)$ and $G_R^T(e)$,*
- 4) *for each vertex $v \in \mathcal{V}$, include $D_\mu^T(v)$ for some triangle T attached to v ,*
- 5) *suppose the vertex $v \notin \mathcal{V}_{NS}$ is connected to v_1, \dots, v_n in counterclockwise order. Let $T_i := \langle v, v_i, v_{i+1} \rangle$ and set $T_0 := T_n = \langle v, v_n, v_1 \rangle$ if v is an interior vertex. Let $1 \leq i_1 < \dots < i_k < n$ be such that e_{i_j} is θ_Δ -near-degenerate at either end, where $e_i := \langle v, v_i \rangle$ for $i = 1, \dots, n$. Let $J_v := \{i_1, \dots, i_k\}$. Then*
 - a) *include $G_L^{T_i}(e_i)$ for all $i \in J_v$,*
 - b) *include $A^{T_i}(v)$ for all $1 \leq i \leq n - 1$ such that $i \notin J_v$,*
 - c) *include $A^{T_n}(v)$ if v is an interior vertex,*
- 6) *for each vertex $v \in \mathcal{V}_S$, include the sets $\mathcal{M}_{v, \mu+1}, \dots, \mathcal{M}_{v, 2r}$ constructed in Theorem 6.1,*
- 7) *for each $v \in \mathcal{V}_{NS} \setminus \mathcal{V}_S$ include the sets $\mathcal{M}_{v, \mu+1}, \dots, \mathcal{M}_{v, 2r}$ constructed in Theorem 6.2.*

Then \mathcal{M} is a stable local minimal determining set for $\mathcal{S}_d^{r, \mu}(\Delta)$.

Proof: We claim that \mathcal{M} is well defined. In particular, if $v \notin \mathcal{V}_{NS}$, then by Lemma 8.2 there exists at least one edge attached to v which is not θ_Δ -near-degenerate at either end. In the numbering of the edges in item 5) above, we can choose this edge to be $\langle v, v_n \rangle$, and the construction insures that for each interior vertex $v \notin \mathcal{V}_{NS}$ and edge $e_i := \langle v, v_i \rangle$ attached to it, if $v_i \notin \mathcal{V}_{NS}$, then \mathcal{M} includes exactly one of the two sets $A^{T_i}(v)$ or $G_L^{T_i}(e_i)$. The construction also guarantees that for all vertices $v \notin \mathcal{V}_{NS}$, there is at least one triangle T with vertex at v such that \mathcal{M} contains the set $A^T(v)$.

To see that \mathcal{M} is a determining set for $\mathcal{S}_d^{r, \mu}(\Delta)$, we show that setting $c_\xi = 0$ for all $\xi \in \mathcal{M}$ implies s is identically zero. Since for every vertex v of Δ the set \mathcal{M} contains $D_\mu^T(v)$ for some triangle attached to v , using the smoothness conditions and Lemma 4.1 we immediately see that all coefficients of s associated with domain points in the disks $D_\mu(v)$ vanish.

Next we compute coefficients on the rings $R_{\mu+1}(v)$ for all v . First we do the vertices v which are not in \mathcal{V}_{NS} . As in [25], we process arcs in a *counterclockwise* direction around v , starting with an edge e such that the preceding triangle T contains the set $A^T(v)$. These computations are based on the smoothness conditions of Lemma 4.1, or (only if the corresponding edge is not θ_Δ -near-degenerate) those of Lemma 4.2. Next we use Theorem 6.1 for each vertex $v \in \mathcal{V}_S$, and Theorem 6.2 for each vertex in $\mathcal{V}_{NS} \setminus \mathcal{V}_S$. To do this, we need the coefficients corresponding to the sets $O_{v, \mu+1}$, but these will all have been set to zero or computed at this point. We now repeat this entire process one ring at a time until we have completed all of the rings up to $R_{2r}(v)$ for all v .

At this point we have shown that all coefficients of s corresponding to domain points in the disks $D_{2r}(v)$ are zero. Since \mathcal{M} contains the sets C^T , the only coefficients remaining correspond to points in sets of the form

$$E^T(e) \setminus \left[D_{2r}(v) \cup D_{2r}(u) \right], \quad (9.2)$$

where $e = \langle v, u \rangle$ is an interior edge. These coefficients can be computed from the associated coefficients in the neighboring triangle (which will have been set to zero) using smoothness conditions as in Lemma 4.1.

We have shown that \mathcal{M} is a determining set for $\mathcal{S}_d^r(\Delta)$. To see that it is minimal, we simply check that its cardinality is equal to the dimension of $\mathcal{S}_d^r(\Delta)$ as given in (2.9) of [24]. Let

$$\begin{aligned} n_a &:= \#A^T(v) = \#G_L^T(e) = \#G_R^T(e) = \binom{2r - \mu + 1}{2}, \\ n_c &:= \#C^T = \binom{d - 3r - 1}{2}, \\ n_d &:= \#D_\mu^T(v) = \binom{\mu + 2}{2}, \\ n_e &:= \#E^T(e) = n_f - 4n_a, \\ n_f &:= \frac{(r + 1)(2d - 4\mu + r - 2)}{2}. \end{aligned}$$

It is easy to check that the number of points chosen in item 6) is $5n_a$ and in item 7) is $4n_a$. This is n_a points for each edge attached to v , and an additional n_a points when v is singular. It follows that

$$\#\mathcal{M} = n_d V + n_a(2E + S + E_B) + n_e E + n_c N, \quad (9.3)$$

where

$$\begin{aligned} E &= \text{number of edges of } \Delta, \\ E_B &= \text{number of boundary edges of } \Delta, \\ N &= \text{number of triangles of } \Delta, \\ S &= \text{number of singular vertices of } \Delta, \\ V &= \text{number of vertices of } \Delta. \end{aligned}$$

Using the fact that $3N = 2E_I + E_B$, (9.3) reduces to

$$\#\mathcal{M} = n_d V + n_a(S - 3N) + n_f E + n_c N, \quad (9.4)$$

which is the formula (2.9) in [24].

This completes the proof that \mathcal{M} is a MDS for $\mathcal{S}_d^r(\Delta)$. We now claim that the corresponding dual splines B_ξ satisfy (5.3) for some constant K depending only on θ_Δ , *i.e.*, \mathcal{M} is stable. To see this, fix $\xi \in \mathcal{M}$, and choose c_η satisfying (5.1). The remaining coefficients of B_ξ can now be computed from smoothness conditions proceeding in the same order as described above in proving that \mathcal{M} was a determining set. But the computations in Lemma 4.1 are always stable, and since we only apply Lemma 4.2 to edges which are not θ_Δ -near-degenerate, those computations are also stable. The computations in the rings $R_{\mu+1}(v), \dots, R_{2r}(v)$ for a singular or θ_Δ -near-singular vertex v are stable by Theorems 6.1 and 6.2.

Finally, we claim that for each $\xi \in \mathcal{M}$, the support of the dual spline B_ξ is at most $\text{star}^3(v_\xi)$ for some vertex v_ξ depending on ξ . We divide the discussion into four cases. Let $\mathcal{M}_0 := \mathcal{M} \setminus \{\xi\}$.

Case 1: Suppose $\xi \in C^T$, where $T := \langle v_1, v_2, v_3 \rangle$ with edges $e_i := \langle v_i, v_{i+1} \rangle$. Then by Lemma 9.2 below, the coefficients of B_ξ must be zero on all three disks $D_{2r}(v_i)$. But since $E^T(e_i) \setminus [D_{2r}(v_i) \cup D_{2r}(v_{i+1})] \subseteq \mathcal{M}_0$, the coefficients of B_ξ are also zero on these sets, and we conclude that the support of B_ξ is just the triangle T .

Case 2: Suppose $\xi \in E^T(e_1) \setminus [D_{2r}(v_1) \cup D_{2r}(v_2)]$, where $T := \langle v_1, v_2, v_3 \rangle$ and $e_1 := \langle v_1, v_2 \rangle$ is a boundary edge of Δ . In this case B_ξ has zero coefficients on the disks $D_{2r}(v_i)$ and on $E^T(e_j) \setminus [D_{2r}(v_j) \cup D_{2r}(v_{j+1})]$ for the other two edges $e_j := \langle v_j, v_{j+1} \rangle$ of T . It follows that the support of B_ξ is just the triangle T .

Case 3: Suppose $\xi \in E^T(e) \setminus [D_{2r}(v) \cup D_{2r}(u)]$, where $e = \langle v, u \rangle$ is an interior edge shared by T and a neighboring triangle \tilde{T} . Then arguing as in Case 2, we see that the support of B_ξ is $T \cup \tilde{T}$.

The situation is more complicated when ξ lies in some disk $D_{2r}(v)$. This is due to the fact that when $d < 4r + 1$ the $2r$ -disks overlap, and nonzero coefficients in one such disk can propagate to a neighboring disk as discussed in [25].

Case 4: Suppose $\xi \in \mathcal{M} \cap D_{2r}(v)$. Suppose z_1, \dots, z_n are the points on the boundary of $\text{star}^3(v)$ in counterclockwise order. Then Lemma 9.2 below shows that the coefficients of B_ξ are zero on the disks $D_{2r}(z_i)$. Now for each $e_i := \langle z_i, z_{i+1} \rangle$, $E^{T_i}(e_i) \setminus [D_{2r}(z_i) \cup D_{2r}(z_{i+1})] \subseteq \mathcal{M}_0$ for some triangle T_i sharing the edge e_i . It follows that the corresponding coefficients are also zero, and we conclude that the support of B_ξ is a subset of $\text{star}^3(v)$. \square

The following lemma is used in the proof of the previous theorem.

Lemma 9.2. *Let \mathcal{M} be the MDS in Theorem 9.1, and let B_ξ be the dual spline corresponding to a $\xi \in \mathcal{M}$. Then B_ξ can have a nonzero coefficient corresponding to a domain point in a disk $D_{2r}(w)$ only if $\xi \in D_{2r}(v)$ for some v , and either $v = w$, or w is connected directly to v with an edge $\langle w, v \rangle$ or by a pair of edges $\langle w, u \rangle, \langle u, v \rangle$, where u is a θ_Δ -near-singular vertex.*

Proof: It is clear from the first part of the proof of Theorem 9.1 that the coefficients in a disk $D_{2r}(w)$ are computed from smoothness conditions which involve only

coefficients in such disks. Hence, if ξ is not in any disk $D_{2r}(v)$, then B_ξ has zero coefficients on all disks $D_{2r}(w)$.

Suppose now that $\xi \in D_{2r}(v)$ and that $w \neq v$ is not connected directly to v with an edge $\langle w, v \rangle$ or by a pair of edges $\langle w, u \rangle, \langle u, v \rangle$ where u is a θ_Δ -near-singular vertex. Let w_1, \dots, w_n be the vertices attached to w and let $e_i := \langle w, w_i \rangle$ and $T_i := \langle w, w_i, w_{i+1} \rangle$. Clearly, $D_\mu(w) \subseteq \mathcal{M}_0$ and $E(e_i) \subseteq \mathcal{M}_0$ for each $1 \leq i \leq n$, where $E(e_i)$ is one of the sets $E^{T_i}(e_i)$ and $E^{T_{i-1}}(e_i)$. There are two cases.

Case 1: $w \in \mathcal{V}_{NS}$. In this case the four vertices w_1, \dots, w_4 are all different from v , which insures $D_\mu(w_i) \subseteq \mathcal{M}_0$ for $i = 1, \dots, 4$. Moreover, none of the w_i is θ_Δ -near-singular since two θ_Δ -near-singular vertices cannot be neighbors. Since the edges $\langle w, w_i \rangle$ are all θ_Δ -near-degenerate at w , we conclude that \mathcal{M}_0 also contains the sets $G_R^{T_{i-1}}(e_i)$, $1 \leq i \leq 4$. This implies that all coefficients of B_ξ in $D_{2r}(w)$ must be zero. To see this, we first calculate the coefficients on the ring $R_{\mu+1}(w)$ from the nonsingular system (6.10) in the proof of Theorem 6.2. We will get zero coefficients if the right-hand side is zero, which happens as soon as the coefficients associated with domain points in $D_\mu^{T_1}(w)$ and the sets $\mathcal{M}_{w, \mu+1}$ and $O_{w, \mu+1}$ in the theorem are zero. Since $D_\mu^{T_1}(w) \cup \mathcal{M}_{w, \mu+1} \subseteq \mathcal{M}_0$, we only have to check $O_{w, \mu+1}$. It is easy to see that $O_{w, \mu+1} \subseteq \bigcup_{i=1}^4 E(e_i)$, and it follows that the coefficients in the disk $D_{\mu+1}(w)$ are zero. Repeating this process for each of the rings $R_{\mu+2}(w), \dots, R_{2r}(w)$, we note that for each $k = \mu + 2, \dots, 2r$,

$$O_{w, k} \subseteq \bigcup_{i=1}^4 [D_\mu(w_i) \cup G_R^{T_{i-1}}(e_i) \cup G_R^{T_i}(e_i) \cup E^{T_{i-1}}(e_i) \cup E^{T_i}(e_i)].$$

Since the coefficients corresponding to \mathcal{M}_0 are zero, and the coefficients corresponding to the disk $D_{k-1}(w)$ are also zero by the induction hypothesis, it is easy to see that the coefficients associated with points in $O_{w, k}$ must be zero, which in turn implies that we have only zero coefficients associated with $R_k(w)$.

Case 2: $w \notin \mathcal{V}_{NS}$. By Lemma 8.2, there is at least one edge attached to w which is not θ_Δ -near-degenerate at either end. Without loss of generality we may assume it is e_n . Then $A^{T_n}(w) \subseteq \mathcal{M}_0$, and the corresponding coefficients of B_ξ must be zero. We now compute coefficients on the ring $R_{\mu+1}(w)$ proceeding in counterclockwise order around w . For $i = 1, \dots, n$, we show that coefficients on the arc $a_{\mu+1, e_i}^r(w)$ are all zero. Assuming this holds for for all $i \leq k - 1$, we now show it for $i = k$.

- a) If e_k is not θ_Δ -near-degenerate at either end, then \mathcal{M}_0 contains $A^{T_k}(w)$ and either $E^{T_k}(e_k)$ or $E^{T_{k-1}}(e_k)$. Hence the coefficients for points in $a_{\mu+1, e_k}^r(w) \setminus (G_L^{T_{k-1}}(e_k) \cup G_L^{T_k}(e_k))$ are zero, and we get all zero coefficients for points on the arc $a_{\mu+1, e_k}^r(w)$ by Lemma 4.2.
- b) If e_k is θ_Δ -near-degenerate at one end and $w_k \notin \mathcal{V}_{NS}$, then \mathcal{M}_0 contains all of the sets $D_\mu(w)$, $D_\mu(w_k)$, $G_R^{T_{k-1}}(e_k)$, $G_L^{T_k}(e_k)$ and either $E^{T_k}(e_i)$ or $E^{T_{k-1}}(e_i)$. Moreover, by the induction hypothesis, the coefficients for points in




Fig. 4. Blocking propagation.

$a_{\mu+1, e_k}^r(w) \cap a_{\mu+1, e_{k-1}}^r(w)$ are zero, too. Then using the smoothness condition, it follows that all coefficients associated with points on the arc $a_{\mu+1, e_k}^r(w)$ must be zero (cf. Lemma 8.2 of [25]).

- c) If $w_k \in \mathcal{V}_{NS}$, then w_k is not connected directly to v with an edge $\langle w_k, v \rangle$ or by a pair of edges $\langle w_k, u \rangle, \langle u, v \rangle$ with u a θ_Δ -near-singular vertex. Then, by Case 1, all coefficients of B_ξ associated with points in $D_{2r}(w_k)$ must be zero, and the same argument as in b) shows that all coefficients associated with points on the arc $a_{\mu+1, e_k}^r(w)$ must be zero.

To complete the proof, we now repeat this process for each of the rings $R_{\mu+2}(w), \dots, R_{2r}(w)$. \square

Fig. 4 illustrates Case 2b of Lemma 9.2 for $r = 4$, $\mu = 6$, and $d = 14$. Suppose e_2 is θ_Δ -near-degenerate at either w_2 or w . Then the coefficients corresponding to points in the sets $G_L^{T_2}(e_2)$ and $G_R^{T_1}(e_2)$ are zero. We have marked those points with the symbol \otimes . The coefficients associated with the points in $E^T(e_2)$ and $A^{T_1}(w)$ are also zero. They are marked with \blacksquare and \odot , respectively. Then using smoothness conditions, we see that all of the coefficients corresponding to points marked with \odot along with those in $A^{T_2}(w)$ (marked with the number 4) must be zero. \square

We conclude this section with an example to illustrate that propagation to $\text{star}^3(v)$ can actually happen.

Example 9.3. Let Δ be the triangulation shown in Fig. 5, and let $r = 5$, $\mu = 8$, $d = 17$.

Discussion: For ease of understanding, we shade the disks D_μ and D_{2r} in dark and light gray, respectively. Suppose \mathcal{M} contains the set $D_\mu^T(v)$ where T is the




Fig. 5. Propagation to $\text{star}^3(v)$.

top triangle in the figure, and suppose ξ is the point at the vertex v . Then B_ξ has support on all of the triangles surrounding v , and in particular, it has a nonzero coefficient corresponding to the point $R_\mu(v) \cap \langle v, u \rangle$. This point is numbered 1 in the figure, and can be identified with the point marked with a \oplus in Fig. 2. As seen from that figure, the nonzero coefficient at point number 1 can propagate to a nonzero coefficient corresponding to the point in the set $A^{T_1}(w)$ which is marked with a \otimes in Fig. 2 and with the number 2 here. (We assume $w_1 = u$ and $w_4 = z$.) Assuming both $\langle w, w_2 \rangle$ and $\langle w, w_3 \rangle$ are θ_Δ -near-degenerate, we get further propagation to a point in the set $A^{T_3}(w)$ marked with the number 4. This set lies in $\text{star}^3(v)$ but outside of $\text{star}^2(v)$. \square

§10. A stable basis for $\mathcal{S}_\mu^{r, \rho_v}(\Delta_v)$ on a cell Δ_v

Before constructing stable local bases for $\mathcal{S}_d^r(\Delta)$ and for general superspline spaces, we need to examine the superspline space $\mathcal{S}_\mu^{r, \rho_v}(\Delta_v)$ with $r \leq \rho_v < \mu$ on an arbitrary

cell Δ_v . Suppose that v is a vertex which is connected to the vertices v_1, \dots, v_n in counterclockwise order, and let $v_{n+1} = v_1$. Let

$$\Delta_v := \{T_i := \langle v, v_i, v_{i+1} \rangle, \quad i = 1, \dots, n\}$$

form a triangulation of the set

$$\Omega_v := \bigcup_{i=1}^n T_i.$$

In this case Δ_v is called an interior cell. We now construct a stable basis for $\mathcal{S}_\mu^{r, \rho_v}(\Delta_v)$.

Let e be the number of edges attached to v with different slopes. Then by Theorem 2.2 of [30],

$$m := \dim \mathcal{S}_\mu^{r, \rho_v}(\Delta_v) = \binom{\rho_v + 2}{2} + n \left[\binom{\mu - r + 1}{2} - \binom{\rho_v - r + 1}{2} \right] + \sigma, \quad (10.1)$$

where

$$\sigma := \sum_{j=\rho_v-r+1}^{\mu-r} (r + j + 1 - je)_+. \quad (10.2)$$

Suppose $\{\xi_i\}_{i=1}^{n_c}$ are the domain points associated with the cell Δ_v . It is easy to see that

$$n_c = n \left[\binom{\mu - 1}{2} + 2\mu - 1 \right] + 1 = n \left[\frac{\mu^2 + \mu}{2} \right] + 1. \quad (10.3)$$

Given $s \in \mathcal{S}_\mu^{r, \rho_v}(\Delta_v)$, we denote the B-coefficient associated with ξ_i by c_i for $i = 1, \dots, n_c$. Associated with each interior edge of Δ_v , there are $\mu - j + 1$ smoothness conditions to insure C^j continuity across that edge, $j = 1, \dots, r$, and $\rho - r - k + 1$ smoothness conditions to insure C^{ρ_v} continuity at v , $k = 1, \dots, \rho_v - r$. This gives a total of

$$\begin{aligned} n_s &:= n \left[\binom{\mu + 1}{2} - \binom{\mu - r + 1}{2} + \binom{\rho_v - r + 1}{2} \right] \\ &= nr \left[\frac{2\mu - r + 1}{2} \right] + n \binom{\rho_v - r + 1}{2} \end{aligned} \quad (10.4)$$

smoothness conditions to insure that s lies in $\mathcal{S}_\mu^{r, \rho_v}(\Delta_v)$. Note that $n_s < n_c$. These conditions can be written in matrix form

$$Ac = 0, \quad (10.5)$$

where $c = (c_1, \dots, c_{n_c})^T$, and A is an appropriate $n_s \times n_c$ matrix.

In general, the system (10.5) includes some redundant smoothness conditions, and so $n_r := \text{rank}(A) < n_s$. Indeed, since $\dim \mathcal{S}_\mu^{r,\rho_v}(\Delta_v) = n_c - n_r$, it follows that

$$\begin{aligned} n_r &= n \left[\frac{\mu^2 + \mu}{2} \right] + 1 - \binom{\rho_v + 2}{2} - n \left[\binom{\mu - r + 1}{2} - \binom{\rho_v - r + 1}{2} \right] - \sigma \\ &= nr \left[\frac{2\mu - r + 1}{2} \right] + 1 - \binom{\rho_v + 2}{2} + n \binom{\rho_v - r + 1}{2} - \sigma. \end{aligned} \quad (10.6)$$

This implies that the number of redundant equations in (10.5) is

$$n_{red} := \binom{\rho_v + 2}{2} - 1 + \sigma. \quad (10.7)$$

Without loss of generality, we may assume that redundant equations have been dropped, and that (10.5) is written in the equivalent form

$$[A_1 \ A_2] c = 0,$$

where A_1 is an $n_r \times m$ matrix and A_2 is an $n_r \times n_r$ matrix. We may also assume that the columns of A (and the corresponding components of c) have been numbered so that the determinant of A_2 has the maximal absolute value over all $n_r \times n_r$ subdeterminants of A .

Algorithm 10.1. For each $i = 1, \dots, m$, let s_i be the spline in $\mathcal{S}_\mu^{r,\rho_v}(\Delta_v)$ with B-coefficients $c = (c_1, \dots, c_{n_c})^T$ chosen so that $c_i = 1$, $c_j = 0$ for $j = 1, \dots, m$ with $j \neq i$, and c_{m+1}, \dots, c_{n_c} are determined from the linear system

$$A_2 \begin{pmatrix} c_{m+1} \\ \vdots \\ c_{n_c} \end{pmatrix} = -A_1(i), \quad (10.8)$$

where $A_1(i)$ is the i -th column of the matrix A_1 .

The splines $\{s_i\}_{i=1}^m$ are clearly linearly independent since

$$\lambda_j s_i = \delta_{i,j}, \quad j = 1, \dots, m, \quad (10.9)$$

where λ_j is the linear functional which picks off the j -th B-coefficient. It follows that they form a basis for $\mathcal{S}_\mu^{r,\rho_v}(\Delta_v)$. We now show that their construction is a stable process, *i.e.*, for each i , all of the coefficients of s_i are uniformly bounded.

Theorem 10.2. Suppose s_i is a basis spline constructed by Algorithm 10.1. Then its B-coefficients satisfy

$$|c_j| \leq 1, \quad j = 1, \dots, n_c. \quad (10.10)$$

Proof: Fix $1 \leq i \leq m$, and let $c = (c_1, \dots, c_{n_c})$ be the vector of coefficients of s_i as computed from Algorithm 10.1. Then (10.10) clearly holds for $j = 1, \dots, m$. Let $m + 1 \leq j \leq n_c$. Then by Cramer's rule,

$$c_j = \frac{\det(\tilde{A}_2)}{\det(A_2)},$$

where \tilde{A}_2 is the matrix obtained from A_2 by replacing the j -th column by $-A_1(i)$. But then $|c_j| \leq 1$ follows by the choice of A_2 . \square

Note that this is a constructive algorithm for building dual basis splines. Indeed, if we take \mathcal{M}_v to be the set of domain points corresponding to the m coefficients c_1, \dots, c_m which are set (as opposed to calculated) in Algorithm 10.1, then obviously \mathcal{M}_v is a minimal determining set for $\mathcal{S}_\mu^{r,\rho_v}(\Delta_v)$.

A completely analogous algorithm can be used to create stable dual basis splines for $\mathcal{S}_\mu^{r,\rho_v}(\Delta_v)$ in the case where Δ_v is a boundary cell.

§11. A stable basis for $\mathcal{S}_d^{r,\rho}(\Delta)$

In this section we combine the constructions of the two previous sections to create stable local bases for the spaces of supersplines $\mathcal{S}_d^{r,\rho}(\Delta)$ defined in (1.1) for all $d \geq 3r + 2$. As in [24], we assume that

$$k_v + k_u < d \quad \text{for each pair of neighboring vertices } v, u \in \mathcal{V},$$

where

$$k_v := \max\{\rho_v, \mu\}, \quad v \in \mathcal{V},$$

with μ as in (1.3).

Given a triangle $T = \langle u, v, w \rangle$, let

$$\tilde{C}^T := C^T \setminus [D_{k_u}^T(u) \cup D_{k_v}^T(v) \cup D_{k_w}^T(w)].$$

Associated with u , let

$$\tilde{A}^T(u) := A^T(u) \setminus D_{k_u}^T(u),$$

with similar definitions for the other two vertices of T . Associated with the edge $e := \langle u, v \rangle$, we define

$$\begin{aligned} \tilde{G}_L^T(e) &:= G_L^T(e) \setminus D_{k_u}^T(u) \\ \tilde{G}_R^T(e) &:= G_R^T(e) \setminus D_{k_v}^T(v), \\ \tilde{E}^T(e) &:= E^T(e) \setminus [D_{k_u}^T(u) \cup D_{k_v}^T(v)], \end{aligned}$$

with similar definitions for the other edges of T .

Theorem 11.1. *Let \mathcal{M} be the following set of domain points:*

- 1) *for each triangle T , include the set \tilde{C}^T ,*
- 2) *for each edge e , include the set $\tilde{E}^T(e)$, where T is some triangle sharing e ,*
- 3) *for each edge of a triangle T such that e lies on the boundary of Ω , include the sets $\tilde{G}_L^T(e)$ and $\tilde{G}_R^T(e)$,*
- 4) *for each vertex $v \in \mathcal{V}$,*
 - a) *include the set $D_{\rho_v}^T(v)$ for some triangle attached to v if $\rho_v \geq \mu$,*
 - b) *include the domain points in $D_\mu(v)$ corresponding to the stable minimal determining set \mathcal{M}_v of Section 10 for $S_\mu^{r,\rho_v}(\Delta_v)$ if $\rho_v < \mu$,*
- 5) *suppose the vertex $v \notin \mathcal{V}_{NS}$ is connected to v_1, \dots, v_n in counterclockwise order. Let $T_i := \langle v, v_i, v_{i+1} \rangle$ and set $T_0 := T_n = \langle v, v_n, v_1 \rangle$ if v is an interior vertex. Let $1 \leq i_1 < \dots < i_k < n$ be such that e_{i_j} is θ_Δ -near-degenerate at either end, where $e_i := \langle v, v_i \rangle$ for $i = 1, \dots, n$. Let $J_v := \{i_1, \dots, i_k\}$. Then*
 - a) *include $\tilde{G}_L^{T_i}(e_i)$ for all $i \in J_v$,*
 - b) *include $\tilde{A}^{T_i}(v)$ for all $1 \leq i \leq n-1$ such that $i \notin J_v$,*
 - c) *include $\tilde{A}^{T_n}(v)$ if v is an interior vertex,*
- 6) *for each vertex $v \in \mathcal{V}_S$, include the sets $\mathcal{M}_{v,k_v+1}, \dots, \mathcal{M}_{v,2r}$ constructed in Theorem 6.1,*
- 7) *for each $v \in \mathcal{V}_{NS} \setminus \mathcal{V}_S$ include the sets $\mathcal{M}_{v,k_v+1}, \dots, \mathcal{M}_{v,2r}$ constructed in Theorem 6.2.*

Then \mathcal{M} is a stable local minimal determining set for $\mathcal{S}_d^{r,\rho}(\Delta)$.

Proof: It is straightforward to check that \mathcal{M} is a determining set for $\mathcal{S}_d^{r,\rho}(\Delta)$. To see that it is minimal, we check that its cardinality is equal to the formula (2.16) in [24] for the dimension of $\mathcal{S}_d^{r,\rho}(\Delta)$. Then locally supported dual basis splines B_ξ can be stably constructed in the same way as in the proof of Theorem 9.1. \square

§12. Stability and local linear independence

We recall (cf. [13,16,17,19,20,21]) that a set $\mathcal{B} = \{B_\nu\}_{\nu \in \mathcal{I}}$ of basis splines for a spline space \mathcal{S} is called locally linearly independent (LLI) provided that for every $T \in \Delta$, the splines $\{B_\nu\}_{\nu \in \Sigma_T}$ are linearly independent on T , where

$$\Sigma_T := \{\nu : T \subseteq \text{supp } B_\nu\}. \quad (12.1)$$

Since the classical univariate B -splines are both stable and locally linearly independent (cf. Theorems 4.18 and 4.41 in [29]), it seems natural to expect that there also exist bases for bivariate spline spaces which possess both of these properties simultaneously. Here we have constructed stable local bases for the spline spaces $\mathcal{S}_d^r(\Delta)$ and their superspline subspaces, while star-supported LLI bases for the same spaces were recently constructed in [19]. But these bases are different, and in fact we have the following surprising result.

Theorem 12.1. *Given $r \geq 1$ and $d \geq 3r + 2$, there are triangulations such that no basis for $\mathcal{S}_d^r(\Delta)$ is simultaneously stable and locally linearly independent.*

Proof: Suppose $\mathcal{B} := \{B_\nu\}_{\nu \in \mathcal{I}}$ is a stable LLI basis for $\mathcal{S}_d^r(\Delta)$ on a triangulation which contains an interior near-singular vertex v . Following the notation of Sect. 6, suppose v is connected to v_1, v_2, v_3, v_4 in counterclockwise order. For each $1 \leq i \leq 4$, let $e_i := \langle v, v_i \rangle$, and $T_i := \langle v, v_i, v_{i+1} \rangle$. Suppose

$$v_{i-1} = r_i v_{i+1} + s_i v + t_i v_i,$$

and suppose that none of the e_i is degenerate at v , *i.e.*, $t_i \neq 0$. For convenience, we define $\alpha_i, \beta_i, \gamma_i, \mu_i$ to be the linear functionals picking off B-coefficients corresponding to the domain points $\xi_{d-2r, r, r}^{T_i}, \xi_{d-2r, r-1, r+1}^{T_i}, \xi_{d-2r, r+1, r-1}^{T_i}, \xi_{d-2r-1, r, r+1}^{T_i}$, respectively.

For each $1 \leq j \leq 4$, we claim that there is a unique spline $S_j \in \mathcal{S}_d^r(\Delta)$ whose only nonzero coefficients are

$$\begin{aligned} \alpha_j S_j &:= 1, & \gamma_j S_j &:= -r_j / (rt_j), & \gamma_{j+1} S_j &:= r_{j+1}^{1-r} / (rt_{j+1}), \\ \beta_{j-1} S_j &:= r_j^{r-1} \gamma_j S_j, & \beta_j S_j &:= r_{j+1}^{r-1} \gamma_{j+1} S_j, \\ \mu_{j-1} S_j &:= r r_j^{r-1} s_j \gamma_j S_j, & \mu_j S_j &:= r r_{j+1}^{r-1} s_{j+1} \gamma_{j+1} S_j. \end{aligned}$$

It can be verified directly that S_j satisfies all C^r smoothness conditions, and thus belongs to $\mathcal{S}_d^r(\Delta)$. It is also easy to see that

$$\text{supp } S_j = T_{j-1} \cup T_j \cup T_{j+1},$$

and by a property of LLI bases (see [13,21]),

$$S_j = \sum_{\nu \in I_j} c_\nu^{[j]} B_\nu, \tag{12.2}$$

where $I_j := \{\nu : \text{supp } B_\nu \subseteq T_{j-1} \cup T_j \cup T_{j+1}\}$ for $j = 1, 2, 3, 4$. We now define

$$\hat{S} = r_2^r S_1 + S_2 + r_3^{-r} S_3 + (r_3 r_4)^{-r} S_4.$$

The definition of barycentric coordinates implies that $r_1 r_2 r_3 r_4 = 1$. Using this fact, it is easy to check that all of the coefficients of \hat{S} are zero except for

$$\alpha_1 \hat{S} := r_2^r, \quad \alpha_2 \hat{S} := 1, \quad \alpha_3 \hat{S} := r_3^{-r}, \quad \alpha_4 \hat{S} := (r_3 r_4)^{-r}.$$

(For example, $\gamma_1 \hat{S} = r_2^r \gamma_1 S_1 + (r_3 r_4)^{-r} \gamma_1 S_4 = r_1 (-r_2^r + (r_1 r_3 r_4)^{-r}) / (rt_1) = 0$.) By (6.13) and Lemma 4.1 of [25], this immediately implies

$$\|\hat{S}\|_\infty \leq K_3,$$

where K_3 depends only on d and θ_Δ .

In view of (12.2), we can write

$$\hat{S} = \sum_{\nu \in I_1 \cup I_2 \cup I_3 \cup I_4} a_\nu B_\nu.$$

By the assumption that the basis \mathcal{B} is stable, we have

$$\|a\|_\infty \leq K_1^{-1} \|\hat{S}\|_\infty \leq K_3/K_1,$$

where K_1 is the constant in (1.2).

For each ν , let $\tilde{B}_\nu = B_\nu - \mathcal{T}_{2r-1} B_\nu$, where $\mathcal{T}_{2r-1} B_\nu \in \mathcal{S}_{2r-1}^r(\Delta_v) \subseteq \mathcal{S}_d^r(\Delta_v)$ is the spline constructed in Lemma 6.4 which interpolates the derivatives of B_ν at v up to order $2r - 1$. Then the B-coefficients of \tilde{B}_ν corresponding to domain points in the disk $D_{2r-1}(v)$ are zero. Moreover, since the basis \mathcal{B} is stable, it follows from Lemma 6.4 that the B-coefficients of \tilde{B}_ν corresponding to domain points on the ring $R_{2r}(v)$ are bounded in absolute value by a constant K_4 depending only on d and θ_Δ .

Since all of the derivatives of \hat{S} up to order $2r - 1$ at v are zero, $\mathcal{T}_{2r-1} \hat{S} = 0$, and we have (on Δ_v)

$$\hat{S} = \sum_{\nu \in I_1 \cup I_2 \cup I_3 \cup I_4} a_\nu \tilde{B}_\nu.$$

Since the support of \tilde{B}_ν is a subset of the support of B_ν (on Δ_v), it follows that $\alpha_2 \tilde{B}_\nu \neq 0$ only if ν lies in the set

$$\tilde{I}_2 := \{\nu : \text{supp } B_\nu = T_1 \cup T_2 \cup T_3\}.$$

This implies

$$1 = \alpha_2 \hat{S} = \sum_{\nu \in \tilde{I}_2} a_\nu \alpha_2 \tilde{B}_\nu \leq \#\tilde{I}_2 \|a\|_\infty \max_{\nu \in \tilde{I}_2} |\alpha_2 \tilde{B}_\nu|,$$

Now clearly $\#\tilde{I}_2 \leq 3 \binom{d+2}{2}$, and hence there exists $\nu_0 \in \tilde{I}_2$ such that

$$|\alpha_2 \tilde{B}_{\nu_0}| \geq K_5 > 0,$$

where K_5 depends only on d and θ_Δ .

Now consider the following C^r smoothness condition across the edge e_2 :

$$\alpha_1 \tilde{B}_{\nu_0} = r_2^r \alpha_2 \tilde{B}_{\nu_0} + r r_2^{r-1} t_2 \gamma_2 \tilde{B}_{\nu_0} + \sum_{k=1}^{r-1} \binom{r}{r-k-1} r_2^{r-k-1} t_2^{k+1} \eta_{2,k} \tilde{B}_{\nu_0},$$

where $\eta_{2,k}$ is the linear functional which picks off the B-coefficient corresponding to $\xi_{d-2r, r+k+1, r-k-1}^{T_i}$ for $k = 1, \dots, r-1$. Since $\alpha_1 \tilde{B}_{\nu_0} = 0$, this implies

$$|\gamma_2 \tilde{B}_{\nu_0}| + \frac{1}{r} \sum_{k=1}^{r-1} \binom{r}{r-k-1} \left| \frac{t_2}{r_2} \right|^k |\eta_{2,k} \tilde{B}_{\nu_0}| \geq \left| \frac{r_2}{t_2} \right| \frac{K_5}{r},$$

which is unbounded as $t_2 \rightarrow 0$. On the other hand, since the B-coefficients $\gamma_2 \tilde{B}_{\nu_0}$, $\eta_{2,k} \tilde{B}_{\nu_0}$, $k = 1, \dots, r-1$, correspond to domain points on the ring $R_{2r}(v)$, they cannot exceed K_4 in absolute value, which leads to a contradiction and completes the proof. \square

Note that the above proof also applies to the superspline spaces $\mathcal{S}_d^{r,\rho}(\Delta)$ whenever $\rho_v < 2r$ for some near-singular vertex v . On the other hand, if $d \geq 4r+1$ and $\rho_v \geq 2r$ for *all* vertices, then the basis constructed in [24] is both stable and LLI.

§13. Remarks

Remark 13.1. It is well known [5–8,23,24] that the dimension of spline spaces and superspline spaces (when $\rho_v < 2r$) generally depends on the exact geometry of the triangulation, and in particular may change as certain near-singular vertices become singular. Thus, it may seem surprising that it is possible to construct stable bases even though the dimension is not stable. This fact was realized already in [1] for $\mathcal{S}_2^1(\Delta_v)$, where Δ_v is a four-cell associated with a near-singular vertex. The spaces considered in [14] are also examples where stable bases were constructed even though the dimensions were not stable.

Remark 13.2. For $d \geq 3r+2$, algorithms for constructing star-supported bases for $\mathcal{S}_d^r(\Delta)$ were presented in [23], and for general superspline subspaces in [24]. The constructions there produce stable bases for $r=0$, and for $d \geq 4r+1$, $\rho_v \geq 2r$ for $r > 0$. However, they are not generally stable if $\rho_v < 2r$, since some of the basis functions do not remain bounded for sequences of triangulations containing vertices which become singular, even if the smallest angle in the triangulations is bounded away from zero. If $\rho_v < \mu$, then many other sequences of triangulations lead to unbounded basis functions when two edges attached to the same vertex become collinear.

Remark 13.3. Stable bases were constructed in [14] for the superspline space $\mathcal{S}_d^{r,\mu}(\Delta)$, and in [25] for a certain special subspace \mathcal{SS} of $\mathcal{S}_d^{r,\mu}(\Delta)$, as a first step in constructing quasi-interpolation operators with optimal approximation order. Note that these constructions differ from our algorithm for $\mathcal{S}_d^{r,\mu}(\Delta)$ described in Section 11. Compared to the construction in [14], our basis splines have substantially smaller supports in general (see also Remark 13.9). For the space \mathcal{SS} the algorithm in [25] produces basis splines with similar small supports, but does not appear to extend to the full spaces $\mathcal{S}_d^r(\Delta)$ and $\mathcal{S}_d^{r,\mu}(\Delta)$.

Remark 13.4. Well-known finite element results, see e.g. [31], imply that the classical superspline subspaces of $\mathcal{S}_d^1(\Delta)$ have stable local bases. In [18] we have recently extended this construction to the full spline spaces $\mathcal{S}_d^1(\Delta)$ with $d \geq 5$. The construction there uses nodal functionals (point evaluation of certain derivatives). Here we have used the linear functionals λ_ξ which pick off the coefficients of the Bernstein-Bézier form.

Remark 13.5. No constructions of stable bases for spline spaces with $d < 3r + 2$ are known for general triangulations. It is also known [9] that for $d < 3r + 2$, there are triangulations for which $\mathcal{S}_d^r(\Delta)$ has no star-supported basis. However, it is possible to construct stable star-supported bases for some values of $d < 3r + 2$ for classes of splines defined on *special* triangulations using macro-element techniques. These include Clough-Tocher and Powell-Sabin refinements, for example. See [10,11,26,27].

Remark 13.6. For multiresolution applications, it is important to work with sequences of triangulations which are nested. In such cases, the corresponding spline spaces $\mathcal{S}_d^r(\Delta)$ are also nested, but in general the various superspline subspaces are not. (See the discussion of this “super-spline effect” in [15,28].)

Remark 13.7. Following the proof of Theorem 5.3 (see also the proof of Theorem 9.2 in [25]), it is not hard to establish

Theorem 13.8. *Suppose \mathcal{M} is a stable local MDS for a spline space \mathcal{S} , and let $\{B_\xi\}_{\xi \in \mathcal{M}}$ be the corresponding dual basis splines. Given $1 \leq p < \infty$, let $B_{\xi,p} := A_\xi^{-1/p} B_\xi$, where A_ξ denotes the area of the support of B_ξ . Then $\{B_{\xi,p}\}_{\xi \in \mathcal{M}}$ is an L_p -stable basis for \mathcal{S} , i.e., there exist constants K_1 and K_2 depending only on d and θ_Δ such that*

$$K_1 \|c\|_p \leq \left\| \sum_{\xi \in \mathcal{M}} c_\xi B_{\xi,p} \right\|_{p,\Omega} \leq K_2 \|c\|_p \quad (13.1)$$

for all choices of the coefficient vector $c = (c_\xi)_{\xi \in \mathcal{M}}$.

Remark 13.9. Our construction guarantees that the supports of the basis splines are at most $\text{star}^3(v)$ for some vertex v in general. In some cases the supports can be made smaller. An explicit construction of stable star-supported bases for C^1 splines can be found in [18]. A careful examination of the construction here shows that for $r = 2$, we also get star-supported stable bases. Moreover, the same holds for general $r > 2$ provided $d \geq 3r + \lfloor (r + 1)/2 \rfloor + 1$, since in this case $D_{2r}(u) \cap D_\mu(w) = \emptyset$ for any two vertices u, w connected by an edge. For $d = 3r + \lfloor (r + 1)/2 \rfloor$, it can be shown that our construction leads to star^2 -supported stable bases. Consequently, star^3 -supported bases only appear for $r \geq 5$.

Remark 13.10. The proofs of Theorems 6.2 and 10.2 are based on Cramer’s rule. This idea of choosing a submatrix with the greatest determinant was used already in [14].

Remark 13.11. As in [25], the stable local bases constructed here can be used to build quasi-interpolation operators with optimal approximation order.

Remark 13.12. The fact that stability and local linear independence are mutually exclusive for spline spaces $\mathcal{S}_d^r(\Delta)$ with $r \geq 1$ was first established for $r = 1$ in [18]. The proof was based on nodal determining sets. Here we have used the Bernstein-Bézier form to establish the same result for general $r \geq 1$.

Remark 13.13. Our construction of a stable local basis can be easily adapted to the spaces of splines and supersplines on a triangulation on the sphere or a sphere-like surface introduced in [2], see also [3,4]. Indeed, since we are using exclusively Bernstein-Bézier techniques, our construction and the entire argument can be carried over in the same way as was done in [3] for the standard local bases of $\mathcal{S}_d^{r,\rho}(\Delta)$.

References

1. Alfeld, P., Private communication, 1995.
2. Alfeld, P., M. Neamtu, and L. L. Schumaker, Bernstein-Bézier polynomials on spheres and sphere-like surfaces, *Comput. Aided Geom. Design* **13** (1996), 333–349.
3. Alfeld, P., M. Neamtu, and L. L. Schumaker, Dimension and local bases of homogeneous spline spaces, *SIAM J. Math. Anal.* **27(5)** (1996), 1482–1501.
4. Alfeld, P., M. Neamtu, and L. L. Schumaker, Fitting scattered data on sphere-like surfaces using spherical splines, *J. Comput. Appl. Math.* **73** (1996), 5–43.
5. Alfeld, P. and L. L. Schumaker, The dimension of bivariate spline spaces of smoothness r for degree $d \geq 4r + 1$, *Constr. Approx.* **3** (1987), 189–197.
6. Alfeld, P., B. Piper, and L. L. Schumaker, Minimally supported bases for spaces of bivariate piecewise polynomials of smoothness r and degree $d \geq 4r + 1$, *Comput. Aided Geom. Design* **4** (1987), 105–123.
7. Alfeld, P., B. Piper, and L. L. Schumaker, An explicit basis for C^1 quartic bivariate splines, *SIAM J. Numer. Anal.* **24** (1987), 891–911.
8. Alfeld, P. and L. L. Schumaker, On the dimension of bivariate splines spaces of smoothness r and degree $d = 3r + 1$, *Numer. Math.* **57** (1990), 651–661.
9. Alfeld, P. and L. L. Schumaker, Non-existence of star-supported spline bases, *SIAM J. Numer. Anal.* **31** (2000), 1482–1501.
10. Alfeld, P. and L. L. Schumaker, Smooth finite elements based on Clough-Tocher triangle splits, *Numer. Math.*, submitted.
11. Alfeld, P. and L. L. Schumaker, Smooth finite elements based on Powell-Sabin triangle splits, manuscript, 2000.
12. Boor, C. de, B -form basics, in *Geometric Modeling: Algorithms and New Trends*, G. E. Farin (ed), SIAM Publications, Philadelphia, 1987, 131–148.
13. Carnicer, J. M. and J. M. Peña, Least supported bases and local linear independence, *Numer. Math.* **67** (1994), 289–301.
14. Chui, C. K., D. Hong, and R.-Q. Jia, Stability of optimal order approximation by bivariate splines over arbitrary triangulations, *Trans. Amer. Math. Soc.* **347** (1995), 3301–3318.
15. Dahmen, W., P. Oswald, and X.-Q. Shi, C^1 -hierarchical bases, *J. Comput. Appl. Math.* **51** (1994), 37–56.

16. Davydov, O., Locally linearly independent basis for C^1 bivariate splines, in *Mathematical Methods for Curves and Surfaces II*, Morten Dæhlen, Tom Lyche, Larry L. Schumaker (eds), Vanderbilt University Press, Nashville & London, 1998, 71–78.
17. Davydov, O., G. Nürnberger, and F. Zeilfelder, Bivariate spline interpolation with optimal approximation order, *Constr. Approx.*, to appear.
18. Davydov, O. and L. L. Schumaker, Stable local nodal bases for C^1 bivariate polynomial splines, in *Curve and Surface Fitting: Saint-Malo 99*, A. Cohen, C. Rabut, and L. L. Schumaker (eds), Vanderbilt University Press, Nashville TN, 2000, 171–180.
19. Davydov, O. and L. L. Schumaker, Locally linearly independent bases for bivariate polynomial splines, *Advances in Comp. Math.*, to appear.
20. Davydov, O., M. Sommer, and H. Strauss, Locally linearly independent systems and almost interpolation, in *Multivariate Approximation and Splines, ISNM 125*, G. Nürnberger, J. W. Schmidt and G. Walz (eds), Birkhäuser, Basel, 1997, 59–72.
21. Davydov, O., M. Sommer, and H. Strauss, On almost interpolation and locally linearly independent bases, *East J. Approx.* **5** (1999), 67–88.
22. Farin, G., Triangular Bernstein-Bézier patches, *Comput. Aided Geom. Design* **3** (1986), 83–127.
23. Hong, D., Spaces of bivariate spline functions over triangulation, *Approx. Theory Appl.* **7** (1991), 56–75.
24. Ibrahim, A. and L. L. Schumaker, Super spline spaces of smoothness r and degree $d \geq 3r + 2$, *Constr. Approx.* **7** (1991), 401–423.
25. Lai, M. J. and L. L. Schumaker, On the approximation power of bivariate splines, *Advances in Comp. Math.* **9** (1998), 251–279.
26. Lai, M.-J. and L. L. Schumaker, Macro-elements and stable local bases for spaces of splines on Clough-Tocher triangulations, *Advances in Comp. Math.*, to appear.
27. Lai, M.-J. and L. L. Schumaker, Macro-elements and stable local bases for splines on Powell-Sabin triangulations, manuscript, 1999.
28. Le Méhauté, A., Nested sequences of triangular finite element spaces, in *Multivariate Approximation: Recent Trends and Results*, W. Haussman, K. Jetter and M. Reimer (eds), Akademie-Verlag, 1997, 133–145.
29. Schumaker, L. L., *Spline Functions: Basic Theory*, Wiley-Interscience, New York, 1981.
30. Schumaker, L. L., Dual bases for spline spaces on cells, *Comput. Aided Geom. Design* **5** (1988), 277–284.
31. Schumaker, L. L., On super splines and finite elements, *SIAM J. Numer. Anal.* **26** (1989), 997–1005.